

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09350 A2

(51) International Patent Classification⁷: **C12N 15/74**,
A61K 39/02, C12N 15/31, 15/67, A61K 39/095, 39/116,
39/118, 39/09, 39/104, 39/102, 39/295, C07K 14/195,
C12N 1/21

(21) International Application Number: PCT/EP00/07424

(22) International Filing Date: 31 July 2000 (31.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9918319.6 3 August 1999 (03.08.1999) GB

(71) Applicant (for all designated States except US):
SMITHKLINE BEECHAM BIOLOGICALS S.A.
[BE/BE]; 89, rue de l'Institut, B-1330 Rixensart (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BERTHET, François-Xavier, Jacques** [FR/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **DALEMANS, Wilfried, L., J.** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **DENOEL, Philippe** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **DEQUESNE, Guy** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **FERON, Christiane** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **LOBET, Yves** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **POOLMAN, Jan** [NL/NL]; SmithKline

Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **THIRY, Georges** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **THONNARD, Joelle** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE). **VOET, Pierre** [BE/BE]; SmithKline Beecham Biologicals s.a., 89, rue de l'Institut, B-1330 Rixensart (BE).

(74) Agent: **DALTON, Marcus, Jonathan, William**; Corporate Intellectual Property, SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VACCINE COMPOSITION

(57) Abstract: The present invention relates to an immuno-protective and non-toxic Gram-negative bleb vaccine suitable for paediatric use. Examples of the Gram-negative strains from which the blebs are made are *N. meningitidis*, *M. catarrhalis* and *H. influenzae*. The blebs of the invention are improved by one or more genetic changes to the chromosome of the bacterium, including up-regulation of protective antigens, down-regulation of immunodominant non-protective antigens, and detoxification of the Lipid A moiety of LPS.

WO 01/09350 A2

VACCINE COMPOSITION

FIELD OF THE INVENTION

The present invention relates to the field of Gram-negative bacterial vaccine
5 compositions, their manufacture, and the use of such compositions in medicine. More particularly it relates to the field of novel outer-membrane vesicle (or bleb) vaccines, and advantageous methods of rendering these vaccines more effective and safer.

BACKGROUND OF THE INVENTION

10 Gram-negative bacteria are separated from the external medium by two successive layers of membrane structures. These structures, referred to as the cytoplasmic membrane and the outer membrane (OM), differ both structurally and functionally. The outer membrane plays an important role in the interaction of pathogenic bacteria with their respective hosts. Consequently, the surface exposed
15 bacterial molecules represent important targets for the host immune response, making outer-membrane components attractive candidates in providing vaccine, diagnostic and therapeutics reagents.

Whole cell bacterial vaccines (killed or attenuated) have the advantage of supplying multiple antigens in their natural micro-environment. Drawbacks around
20 this approach are the side effects induced by bacterial components such as endotoxin and peptidoglycan fragments. On the other hand, acellular subunit vaccines containing purified components from the outer membrane may supply only limited protection and may not present the antigens properly to the immune system of the host.

Proteins, phospholipids and lipopolysaccharides are the three major
25 constituents found in the outer-membrane of all Gram-negative bacteria. These molecules are distributed asymmetrically: membrane phospholipids (mostly in the inner leaflet), lipooligosaccharides (exclusively in the outer leaflet) and proteins (inner and outer leaflet lipoproteins, integral or polytopic membrane proteins). For many bacterial pathogens which impact on human health, lipopolysaccharide and outer-
30 membrane proteins have been shown to be immunogenic and amenable to confer protection against the corresponding disease by way of immunization.

The OM of Gram-negative bacteria is dynamic and, depending on the environmental conditions, can undergo drastic morphological transformations. Among these manifestations, the formation of outer-membrane vesicles or “blebs” has been studied and documented in many Gram-negative bacteria (Zhou, L *et al.* 1998. *FEMS Microbiol. Lett.* 163: 223-228). Among these, a non-exhaustive list of bacterial pathogens reported to produce blebs include: *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella melitensis*, *Brucella ovis*, *Chlamydia psittaci*, *Chlamydia trachomatis*, *Escherichia coli*, *Haemophilus influenzae*, *Legionella pneumophila*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*. Although the biochemical mechanism responsible for the production of OM blebs is not fully understood, these outer membrane vesicles have been extensively studied as they represent a powerful methodology in order to isolate outer-membrane protein preparations in their native conformation. In that context, the use of outer-membrane preparations is of particular interest to develop vaccines against *Neisseria*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa* and *Chlamydia*. Moreover, outer membrane blebs combine multiple proteinaceous and non-proteinaceous antigens that are likely to confer extended protection against intra-species variants.

In comparison with the other, more widely used, types of bacterial vaccine (whole cell bacterial and purified subunit vaccines), the inventors will show that outer membrane bleb vaccines (if modified in certain ways) may represent the ideal compromise.

The wide-spread use of bacterial subunit vaccines has been due to the intensive study of bacterial surface proteins that have been found to be useful in vaccine applications [for instance *B. pertussis* pertactin]. These proteins are loosely associated with the bacterial outer membrane and can be purified from culture supernatant or easily extracted from the bacterial cells. However it has also been shown that structural, integral outer membrane proteins are also protective antigens. Examples are PorA for *N. meningitidis* serogroup B; D15 for *H. influenzae*; OMP CD for *M. catarrhalis*; OMP F for *P. Aeruginosa*. Such proteins however have rather specific structural features, particularly multiple amphipathic β -sheets, which complicates their straightforward use as purified (recombinant) subunit vaccines.

In addition, it has become clear that multiple component vaccines are needed (in terms of bacterial surface proteins and integral membrane proteins) to supply a reasonable level of protection. For instance, in the case of *B. pertussis* subunit vaccines multicomponent vaccines are superior to mono or bicomponent products.

5 In order to incorporate integral outer-membrane proteins into such a subunit product, native (or near-native) conformational folding of the proteins must be present in the product in order to have a useful immunological effect. The use of excreted outer membrane vesicles or blebs may be an elegant solution to the problem of including protective integral membrane proteins into a subunit vaccine whilst still
10 ensuring that they fold properly.

N. meningitidis serogroup B (menB) excretes outer membrane blebs in sufficient quantities to allow their manufacture on an industrial scale. Such multicomponent outer-membrane protein vaccines from naturally-occurring menB strains have been found to be efficacious in protecting teenagers from menB disease
15 and have become registered in Latin America. An alternative method of preparing outer-membrane vesicles is via the process of detergent extraction of the bacterial cells (EP 11243).

Examples of bacterial species from which bleb vaccines can be made are the following.

20

Neisseria meningitidis:

Neisseria meningitidis (meningococcus) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It occasionally causes invasive bacterial diseases such as bacteremia and meningitis. The incidence of
25 meningococcal disease shows geographical seasonal and annual differences (Schwartz, B., Moore, P.S., Broome, C.V.; Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Most disease in temperate countries is due to strains of serogroup B and varies in incidence from 1-10/100,000/year total population sometimes reaching higher values (Kaczmarek, E.B. (1997), Commun. Dis. Rep. Rev. 7: R55-9, 1995;
30 Scholten, R.J.P.M., Bijlmer, H.A., Poolman, J.T. et al. Clin. Infect. Dis. 16: 237-246, 1993; Cruz, C., Pavez, G., Aguilar, E., et al. Epidemiol. Infect. 105: 119-126, 1990).

Age-specific incidences in the two high risk-groups, infants and teenagers, reach higher levels.

Epidemics dominated by serogroup A meningococci occur, mostly in central Africa, sometimes reaching levels up to 1000/100,000/year (Schwartz, B., Moore, P.S., Broome, C.V. Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Nearly all cases of meningococcal disease as a whole are caused by serogroup A, B, C, W-135 and Y meningococci. A tetravalent A, C, W-135, Y capsular polysaccharide vaccine is available (Armand, J., Arminjon, F., Mynard, M.C., Lafaix, C., J. Biol. Stand. 10: 335-339, 1982).

The polysaccharide vaccines are currently being improved by way of chemically conjugating them to carrier proteins (Lieberman, J.M., Chiu, S.S., Wong, V.K., et al. JAMA 275 : 1499-1503, 1996). A serogroup B vaccine is not available, since the B capsular polysaccharide is non-immunogenic, most likely because it shares structural similarity to host components (Wyle, F.A., Artenstein, M.S., Brandt, M.L. et al. J. Infect. Dis. 126: 514-522, 1972; Finne, J.M., Leinonen, M., Mäkelä, P.M. Lancet ii.: 355-357, 1983).

For many years efforts have been focused on developing meningococcal outer membrane based vaccines (de Moraes, J.C., Perkins, B., Camargo, M.C. et al. Lancet 340: 1074-1078, 1992; Bjune, G., Hoiby, E.A. Gronnesby, J.K. et al. 338: 1093-1096, 1991). Such vaccines have demonstrated efficacies from 57% - 85% in older children (>4 years) and adolescents. Most of these efficacy trials were performed with OMV (outer membrane vesicles, derived by LPS depletion from blebs) vaccines derived from wild-type *N. meningitidis* B strains.

Many bacterial outer membrane components are present in these vaccines, such as PorA, PorB, Rmp, Opc, Opa, FrpB and the contribution of these components to the observed protection still needs further definition. Other bacterial outer membrane components have been defined (using animal or human antibodies) as potentially being relevant to the induction of protective immunity, such as TbpB, NspA (Martin, D., Cadieux, N., Hamel, J., Brodeux, B.R., J. Exp. Med. 185: 1173-1183, 1997; Lissolo, L., Maître-Wilmotte, C., Dumas, p. et al., Inf. Immun. 63: 884-890, 1995). The mechanism of protective immunity will involve antibody mediated bactericidal activity and opsonophagocytosis.

The frequency of *Neisseria meningitidis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiple antibiotic resistant strains, and increased exposure due to an increase in social activities (for instance swimming pools or theatres). It is no longer uncommon to isolate *Neisseria meningitidis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

Moraxella catarrhalis

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It is responsible for several pathologies, the main ones being otitis media in infants and children, and pneumonia the elderly. It is also responsible for sinusitis, nosocomial infections and, less frequently, for invasive diseases.

Bactericidal antibodies have been identified in most adults tested (Chapman, AJ et al. (1985) J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist serum bactericidal activity: in general, isolates from diseased individuals are more resistant than those who are simply colonized (Hol, C et al. (1993) Lancet 341:1281, Jordan, KL et al. (1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be considered as a virulence factor of the bacteria. An opsonizing activity has been observed in the sera of children recovering from otitis media.

The antigens targetted by these different immune responses in humans have not been identified, with the exception of OMP B1, a 84 kDa protein, the expression of which is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S, et al. (1995) Infect.Immun. 63:1516), and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical methods for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD) is highly conserved among *M. catarrhalis* strains, and presents homologies with a

porin of *Pseudomonas aeruginosa*, which has been demonstrated to be efficacious against this bacterium in animal models.

M. catarrhalis produces outer membrane vesicles (Blebs). These Blebs have been isolated or extracted by using different methods (Murphy T.F., Loeb M.R. 1989. Microb. Pathog. 6: 159-174; Unhanand M., Maciver, I., Ramillo O., Arencibia-Mireles O., Argyle J.C., McCracken G.H. Jr., Hansen E.J. 1992. J. Infect. Dis. 165:644-650). The protective capacity of such Bleb preparations has been tested in a murine model for pulmonary clearance of *M. catarrhalis*. It has been shown that active immunization with Bleb vaccine or passive transfer of anti-Blebs antibody induces significant protection in this model (Maciver I., Unhanand M., McCracken G.H. Jr., Hansen, E.J. 1993. J. Infect. Dis. 168: 469-472).

Haemophilus influenzae

Haemophilus influenzae is a non-motile Gram-negative bacterium. Man is its only natural host. *H. influenzae* isolates are usually classified according to their polysaccharide capsule. Six different capsular types designated 'a' through 'f' have been identified. Isolates that fail to agglutinate with antisera raised against one of these six serotypes are classified as nontypeable, and do not express a capsule.

H. influenzae type b (Hib) is clearly different from the other types in that it is a major cause of bacterial meningitis and systemic diseases. Nontypeable strains of *H. influenzae* (NTHi) are only occasionally isolated from the blood of patients with systemic disease. NTHi is a common cause of pneumonia, exacerbation of chronic bronchitis, sinusitis and otitis media. NTHi strains demonstrate a large variability as identified by clonal analysis, whilst Hib strains as a whole are more homogeneous.

Various proteins of *H. influenzae* have been shown to be involved in pathogenesis or have been shown to confer protection upon vaccination in animal models.

Adherence of NTHi to human nasopharyngeal epithelial cells has been reported (Read RC. et al. 1991. J. Infect. Dis. 163:549). Apart from fimbriae and pili (Brinton CC. et al. 1989. Pediatr. Infect. Dis. J. 8:S54; Kar S. et al. 1990. Infect. Immun. 58:903; Gildorf JR. et al. 1992. Infect. Immun. 60:374; St. Geme JW et al. 1991. Infect. Immun. 59:3366; St. Geme JW et al. 1993. Infect. Immun. 61: 2233), many adhesins have been

identified in NTHi. Among them, two surface exposed high-molecular-weight proteins designated HMW1 and HMW2 have been shown to mediate adhesion of NTHi to epithelial cells (St. Geme JW. et al. 1993. Proc. Natl. Acad. Sci. USA 90:2875). Another family of high-molecular-weight proteins has been identified in NTHi strains that lack
5 proteins belonging to HMW1/HMW2 family. The NTHi 115-kDa Hia protein (Barenkamp SJ., St Geme S.W. 1996. Mol. Microbiol. In press) is highly similar to the Hsf adhesin expressed by *H. influenzae* type b strains (St. Geme JW. et al. 1996. J. Bact. 178:6281). Another protein, the Hap protein shows similarity to IgA1 serine proteases and has been shown to be involved in both adhesion and cell entry (St. Geme JW. et al. 1994.
10 Mol. Microbiol. 14:217).

Five major outer membrane proteins (OMP) have been identified and numerically numbered. Original studies using *H. influenzae* type b strains showed that antibodies specific for P1 and P2 OMPs protected infant rats from subsequent challenge (Loeb MR. et al. 1987. Infect. Immun. 55:2612; Musson RS. Jr. et al.
15 1983. J. Clin. Invest. 72:677). P2 was found to be able to induce bactericidal and opsonic antibodies, which are directed against the variable regions present within surface exposed loop structures of this integral OMP (Haase EM. et al. 1994 Infect. Immun. 62:3712; Troelstra A. et al. 1994 Infect. Immun. 62:779). The lipoprotein P4 also may induce bactericidal antibodies (Green BA. et al. 1991.
20 Infect. Immun. 59:3191).

OMP P6 is a conserved peptidoglycan associated lipoprotein making up 1-5 % of the outer membrane (Nelson MB. et al. 1991. Infect. Immun. 59:2658). Later a lipoprotein of about the same molecular weight was recognized called PCP (P6 cross-reactive protein) (Deich RM. et al. 1990. Infect. Immun. 58:3388). A mixture of the
25 conserved lipoproteins P4, P6 and PCP did not reveal protection as measured in a chinchilla otitis-media model (Green BA. et al. 1993. Infect. immun. 61:1950). P6 alone appears to induce protection in the chinchilla model (Demaria TF. et al. 1996. Infect. Immun. 64:5187).

A fimbrin protein (Miyamoto N., Bakaletz, LO. 1996. Microb. Pathog. 21:343)
30 has also been described with homology to OMP P5, which itself has sequence homology to the integral *Escherichia coli* OmpA (Miyamoto N., Bakaletz, LO. 1996. Microb.

Pathog. 21:343; Munson RS. Jr. et al. 1993. Infect. Immun. 61:1017). NTHi seem to adhere to mucus by way of fimbriae.

In line with the observations made with gonococci and meningococci, NTHi expresses a dual human transferrin receptor composed of TbpA and TbpB when grown
5 under iron limitation. Anti-TbpB antibody protected infant rats (Loosmore SM. et al. 1996. Mol. Microbiol. 19:575). Hemoglobin / haptoglobin receptor also have been described for NTHi (Maciver I. et al. 1996. Infect. Immun. 64:3703). A receptor for Haem:Hemopexin has also been identified (Cope LD. et al. 1994. Mol. Microbiol. 13:868). A lactoferrin receptor is also present amongst NTHi, but is not yet
10 characterized (Schryvers AB. et al. 1989. J. Med. Microbiol. 29:121). A protein similar to neisserial FrpB-protein has not been described amongst NTHi.

An 80kDa OMP, the D15 surface antigen, provides protection against NTHi in a mouse challenge model. (Flack FS. et al. 1995. Gene 156:97). A 42kDa outer membrane lipoprotein, LPD is conserved amongst *Haemophilus influenzae* and
15 induces bactericidal antibodies (Akkoyunlu M. et al. 1996. Infect. Immun. 64:4586). A minor 98kDa OMP (Kimura A. et al. 1985. Infect. Immun. 47:253), was found to be a protective antigen, this OMP may very well be one of the Fe-limitation inducible OMPs or high molecular weight adhesins that have been characterized thereafter. *H. Influenzae* produces IgA1-protease activity (Mulks MH., Shoberg RJ. 1994. Meth.
20 Enzymol. 235:543). IgA1-proteases of NTHi have a high degree of antigenic variability (Lomholt H., van Alphen L., Kilian, M. 1993. Infect. Immun. 61:4575).

Another OMP of NTHi, OMP26, a 26-kDa protein has been shown to enhance pulmonary clearance in a rat model (Kyd, J.M., Cripps, A.W. 1998. Infect. Immun. 66:2272). The NTHi HtrA protein has also been shown to be a protective antigen.
25 Indeed, this protein protected Chinchilla against otitis media and protected infant rats against *H. influenzae* type b bacteremia (Loosmore S.M. et al. 1998. Infect. Immun. 66:899).

Outer membrane vesicles (or blebs) have been isolated from *H. influenzae* (Loeb M.R., Zachary A.L., Smith D.H. 1981. J. Bacteriol. 145:569-604; Stull T.L.,
30 Mack K., Haas J.E., Smit J., Smith A.L. 1985. Anal. Biochem. 150: 471-480). The vesicles have been associated with the induction of blood-brain barrier permeability (Wiwpelwey B., Hansen E.J., Scheld W.M. 1989 Infect. Immun. 57: 2559-2560), the

induction of meningeal inflammation (Mustafa M.M., Ramilo O., Syrogiannopoulos G.A., Olsen K.D., McCracken G.H. Jr., Hansen, E.J. 1989. J. Infect. Dis. 159: 917-922) and to DNA uptake (Concino M.F., Goodgal S.H. 1982 J. Bacteriol. 152: 441-450). These vesicles are able to bind and be absorbed by the nasal mucosal epithelium
5 (Harada T., Shimuzu T., Nishimoto K., Sakakura Y. 1989. Acta Otorhinolarygol. 246: 218-221) showing that adhesins and/or colonization factors could be present in Blebs. Immune response to proteins present in outer membrane vesicles has been observed in patients with various *H. influenzae* diseases (Sakakura Y., Harada T., Hamaguchi Y., Jin C.S. 1988. Acta Otorhinolarygol. Suppl. (Stockh.) 454: 222-226; Harada T.,
10 Sakakura Y., Miyoshi Y. 1986. Rhinology 24: 61-66).

Pseudomonas aeruginosa:

The genus *Pseudomonas* consists of Gram-negative, polarly flagellated, straight and slightly curved rods that grow aerobically and do not forms spores.
15 Because of their limited metabolic requirements, *Pseudomonas spp.* are ubiquitous and are widely distributed in the soil, the air, sewage water and in plants. Numerous species of *Pseudomonas* such as *P. aeruginosa*, *P. pseudomallei*, *P. mallei*, *P. maltophilia* and *P. cepacia* have also been shown to be pathogenic for humans. Among this list, *P. aeruginosa* is considered as an important human pathogen since it
20 is associated with opportunistic infection of immuno-compromised host and is responsible for high morbidity in hospitalized patients. Nosocomial infection with *P. aeruginosa* afflicts primarily patients submitted for prolonged treatment and receiving immuno-suppressive agents, corticosteroids, antimetabolites antibiotics or radiation.

The *Pseudomonas*, and particularly *P. aeruginosa*, produces a variety of toxins
25 (such as hemolysins, fibrinolysins, esterases, coagulases, phospholipases, endo- and exo-toxins) that contribute to the pathogenicity of these bacteria. Moreover, these organisms have high intrinsic resistance to antibiotics due to the presence of multiple drug efflux pumps. This latter characteristic often complicates the outcome of the disease.

30 Due to the uncontrolled use of antibacterial chemotherapeutics the frequency of nosocomial infection caused by *P. aeruginosa* has increased considerably over the last 30 years. In the US, for example, the economic burden of *P. aeruginosa*

nosocomial infection is estimated to 4.5 billion US\$ annually. Therefore, the development of a vaccine for active or passive immunization against *P. aeruginosa* is actively needed (for review see Stanislavsky et al. 1997. FEMS Microbiol. Lett. 21: 243-277).

5 Various cell-associated and secreted antigens of *P. aeruginosa* have been the subject of vaccine development. Among *Pseudomonas* antigens, the mucoid substance, which is an extracellular slime consisting predominantly of alginate, was found to be heterogenous in terms of size and immunogenicity. High molecular mass alginate components (30-300 kDa) appear to contain conserved epitopes while lower
10 molecular mass alginate components (10-30 kDa) possess conserved epitopes in addition to unique epitopes. Among surface-associated proteins, PcrV, which is part of the type III secretion-translocation apparatus, has also been shown to be an interesting target for vaccination (Sawa et al. 1999. Nature Medicine 5:392-398).

Surface-exposed antigens including O-antigens (O-specific polysaccharide of
15 LPS) or H-antigens (flagellar antigens) have been used for serotyping due to their highly immunogenic nature. Chemical structures of repeating units of O-specific polysaccharides have been elucidated and these data allowed the identification of 31 chemotypes of *P. aeruginosa*. Conserved epitopes among all serotypes of *P. aeruginosa* are located in the core oligosaccharide and the lipid A region of LPS and
20 immunogens containing these epitopes induce cross-protective immunity in mice against different *P. aeruginosa* immunotypes. The outer core of LPS was implicated to be a ligand for binding of *P. aeruginosa* to airway and ocular epithelial cells of animals. However, heterogeneity exists in this outer core region among different serotypes. Epitopes in the inner core are highly conserved and have been
25 demonstrated to be surface-accessible, and not masked by O-specific polysaccharide.

To examine the protective properties of OM proteins, a vaccine containing *P. aeruginosa* OM proteins of molecular masses ranging from 20 to 100 kDa has been used in pre-clinical and clinical trials. This vaccine was efficacious in animal models against *P. aeruginosa* challenge and induced high levels of specific antibodies in
30 human volunteers. Plasma from human volunteers containing anti-*P. aeruginosa* antibodies provided passive protection and helped the recovery of 87% of patients with severe forms of *P. aeruginosa* infection. More recently, a hybrid protein

containing parts of the outer membrane proteins OprF (amino acids 190-342) and OprI (amino acids 21-83) from *Pseudomonas aeruginosa* fused to the glutathione-S-transferase was shown to protect mice against a 975-fold 50% lethal dose of *P. aeruginosa* (Knapp et al. 1999. Vaccine. 17:1663-1669).

5

The present inventors have realised a number of drawbacks associated with the above wild-type bleb vaccines (either naturally occurring or chemically made).

Examples of such problems are the following:

- 10 - the presence of immunodominant but variable proteins on the bleb (PorA; TbpB, Opa [*N. meningitidis* B]; P2, P5 [non-typeable *H. influenzae*]) – such blebs being effective only against a restricted selection of bacterial species. Type-specificity of the bactericidal antibody response may preclude the use of such vaccines in infancy.
- 15 - the presence of unprotective (non relevant) antigens (Rmp, H8, ...) on the bleb – antigens that are decoys for the immune system
- the lack of presence of important molecules which are produced conditionally (for instance iron-regulated outer membrane proteins, IROMP's, in vivo regulated expression mechanisms) – such conditions are hard to control in bleb production in order to optimise the amount of antigen on the surface
- 20 - the low level of expression of protective, (particularly conserved) antigens (NspA, P6)
- the toxicity of the LPS remaining on the surface of the bleb
- 25 - the potential induction of an autoimmune response because of host-identical structures (for example the capsular polysaccharide in *Neisseria meningitidis* serogroup B, the lacto-N-neotetraose in *Neisseria* LPS, saccharide structure within ntHi LPS, saccharide structures within Pili).

Such problems may prevent the use of bleb vaccines as human vaccine reagents. This is particularly so for paediatric use (<4 years) where reactogenicity against bleb vaccines is particularly important, and where bleb vaccines (for instance the previously mentioned marketed MenB bleb vaccine) have been shown to be

ineffective at immuno-protecting. Accordingly, the present invention provides methods of alleviating the above problems using genetically engineered bacterial strains, which result in improved bleb vaccines. Such methods will be especially useful in the generation of new vaccines against bacterial pathogens such as *Neisseria meningitidis*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, and others.

The bleb vaccines of the invention are designed to focus the immune response on a few protective (preferably conserved) antigens or epitopes - formulated in a multiple component vaccine. Where such antigens are integral OMPs, the outer membrane vesicles of bleb vaccines will ensure their proper folding. This invention provides methods to optimize the OMP and LPS composition of OMV (bleb) vaccines by deleting immunodominant variable as well as non protective OMPs, by creating conserved OMPs by deletion of variable regions, by upregulating expression of protective OMPs, and by eliminating control mechanisms for expression (such as iron restriction) of protective OMPs. In addition the invention provides for the reduction in toxicity of lipid A by modification of the lipid portion or by changing the phosphoryl composition whilst retaining its adjuvant activity or by masking it. Each of these new methods of improvement individually improve the bleb vaccine, however a combination of one or more of these methods work in conjunction so as to produce an optimised engineered bleb vaccine which is immuno-protective and non-toxic - particularly suitable for paediatric use.

SUMMARY OF THE INVENTION

The present invention provides a genetically-engineered bleb preparation from a Gram-negative bacterial strain characterized in that said preparation is obtainable by employing one or more processes selected from the following group:

- a) a process of reducing immunodominant variable or non-protective antigens within the bleb preparation comprising the steps of determining the identity of such antigen, engineering a bacterial strain to produce less or none of said antigen, and making blebs from said strain;
- b) a process of upregulating expression of protective, endogenous (and preferably conserved) OMP antigens within the bleb preparation

- comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of a gene encoding said antigen such that said gene is expressed at a level higher than in the non-modified bleb, and making blebs from said strain;
- 5 c) a process of upregulating expression of conditionally-expressed, protective (and preferably conserved) OMP antigens within the bleb preparation comprising the steps of identifying such an antigen, engineering a bacterial strain so as to remove the repressive control mechanisms of its expression (such as iron restriction), and making blebs from said strain;
- 10 d) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS toxic, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain;
- 15 e) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS less toxic, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of said gene such that said gene is expressed at a level higher than in the non-
- 20 modified bleb, and making blebs from said strain;
- f) a process of reducing lipid A toxicity within the bleb preparation and increasing the levels of protective antigens, comprising the steps of engineering the chromosome of a bacterial strain to incorporate a gene encoding a Polymyxin A peptide, or a derivative or analogue thereof, fused
- 25 to a protective antigen, and making blebs from said strain;
- g) a process of creating conserved OMP antigens on the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to delete variable regions of a gene encoding said antigen, and making blebs from said strain;
- 30 h) a process of reducing expression within the bleb preparation of an antigen which shares a structural similarity with a human structure and may be capable of inducing an auto-immune response in humans (such as the

capsular polysaccharide of *N. meningitidis* B), comprising the steps of identifying a gene involved in the biosynthesis of the antigen, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain; or

- 5 i) a process of upregulating expression of protective, endogenous (and preferably conserved) OMP antigens within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce into the chromosome one or more further copies of a gene encoding said antigen controlled by a heterologous, stronger
10 promoter sequence, and making blebs from said strain.

Further aspects of the invention include, preferential processes for obtaining the above bleb preparation, including optimal positioning of strong promoters for the upregulation of expression of antigens within blebs, preferential antigens for
15 upregulation and downregulation for various bacterial strains in order to obtain bleb preparations particularly suitable for vaccine use. Preferential formulations comprising the blebs of the invention are also provided which are particularly suitable for global vaccines against certain disease states. Vectors for producing the blebs of the invention, and modified bacterial strains from which the blebs of the invention are
20 produced are still further aspects of the invention.

The present invention provides for the first time a bleb vaccine which is immuno-protective and non-toxic when used with children under 4 years of age.

BRIEF DESCRIPTION OF THE DRAWINGS

25

Figure 1: Reactivity of the 735 mAb on different colonies.

Figure 2: Reactivities of specific monoclonal antibodies by whole cell Elisa.

30 **Figure 3:** Schematic representation of the pCMK vectors used to deliver genes, operons and/or expression cassettes in the genome of *Neisseria meningitidis*.

Figure 4: Analysis of PorA expression in total protein extracts of recombinant *N. meningitidis* serogroupB (H44/76 derivatives). Total proteins were recovered from *cps*- (lanes 3 and 4), *cps*- *porA*::*pCMK*+ (lanes 2 and 5) and *cps*- *porA*::*nspA* (lanes 1 and 6) recombinant *N. meningitidis* serogroupB strains and were analyzed under SDS-PAGE conditions in a 12% polyacrylamide gel. Gels were stained with Coomassie blue (lanes 1 to 3) or transferred to a nitrocellulose membrane and immuno-stained with an anti-PorA monoclonal antibody.

Figure 5: Analysis of NspA expression in protein extracts of recombinant *N. meningitidis* serogroupB strains (H44/76 derivatives). Proteins were extracted from whole bacteria (lanes 1 to 3) or outer-membrane blebs preparations (lanes 4 to 6) separated by SDS-PAGE on a 12% acrylamide gel and analyzed by immuno-blotting using an anti-NspA polyclonal serum. Samples corresponding to *cps*- (lanes 1 and 6), *cps*- *porA*::*pCMK*+ (lanes 3 and 4) and *cps*- *porA*::*nspA* (lanes 2 and 5) were analyzed. Two forms of NspA were detected: a mature form (18kDa) co-migrating with the recombinant purified NspA, and a shorter form (15kDa).

Figure 6: Analysis of D15/omp85 expression in protein extracts of recombinant *N. meningitidis* serogroupB strains (H44/76 derivatives). Proteins were extracted from outer-membrane blebs preparations and were separated by SDS-PAGE on a 12% acrylamide gel and analyzed by immuno-blotting using an anti-omp85 polyclonal serum. Samples corresponding to *cps*- (lane 2), and *cps*-, *PorA*+, *pCMK*+*Omp85/D15* (lane 1) recombinant *N. meningitidis* serogroupB strains were analyzed.

Figure 7: General strategy for modulating gene expression by promoter delivery (RS stands for restriction site).

Figure 8: Analysis of outer-membrane blebs produced by recombinant *N. meningitidis* serogroupB *cps*- strains (H44/76 derivatives). Proteins were extracted from outer-membrane bleb preparations and were separated by SDS-PAGE under reducing conditions on a 4-20% gradient polyacrylamide gel. The gel was stained

with Coomassie brilliant blue R250. Lanes 2, 4, 6 corresponded to 5µg of total proteins whereas lanes 3, 5 and 7 were loaded with 10µg proteins.

5 **Figure 9:** Construction of a promoter replacement plasmid used to up-regulate the expression/production of Omp85/D15 in *Neisseria meningitidis* H44/76.

Figure 10: Analysis of OMP85 expression in total protein extracts of recombinant *NmB* (H44/76 derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with rabbit anti-OMP85 (*N.gono*) monoclonal antibody (B).
10

Figure 11: Analysis of OMP85 expression in OMV preparations from recombinant *NmB* (H44/76 derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with rabbit anti-OMP85 polyclonal antibody (B).
15

Figure 12: Schematic representation of the recombinant PCR strategy used to delete the *lacO* in the chimeric *porA/lacO* promoter.

20 **Figure 13:** Analysis of Hsf expression in total protein extracts of recombinant *N. meningitidis* serogroup B (H44/76 derivatives). Total proteins were recovered from Cps-PorA+(lanes 1), and Cps-PorA+/Hsf (lanes 2) recombinant *N. meningitidis* serogroup B strains and were analyzed under SDS-PAGE conditions in a 12% polyacrylamide gel. Gels were stained with Coomassie blue .

25 **Figure 14:** Analysis of GFP expression in total protein extracts of recombinant *N. meningitidis* (H44/76 derivative). Total protein were recovered from Cps-, PorA+ (lane1), Cps-, PorA- GFP+ (lane2 & 3) recombinant strains. Proteins were separated by PAGE-SDS in a 12% polyacrylamide gel and then stained with Coomassie blue.

30 **Figure 15:** Illustration of the pattern of major proteins on the surface of various recombinant bleb preparations as analysed by SDS-PAGE (Coomassie staining).

Figure 16: Specific anti-Hsf response for various bleb and recombinant bleb preparations using purified recombinant Hsf protein.

- 5 **Figure 17:** Analysis of NspA expression in total protein extracts of recombinant *NmB* (serogroup B derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with mouse anti-PorA monoclonal antibody (B) or mouse anti-NspA polyclonal antibody (C).

10

DESCRIPTION OF THE INVENTION

The present invention relates to a general set of tools and methods capable of being used for manufacturing improved, genetically engineered blebs from Gram-negative bacterial strains. The invention includes methods used to make recombinant
15 blebs more immunogenic, less toxic and safer for their use in a human and/or animal vaccine. Moreover, the present invention also describes specific methods necessary for constructing, producing, obtaining and using recombinant, engineered blebs from various Gram-negative bacteria, for vaccine, therapeutic and/or diagnostic purposes. By the methods of the invention, the biochemical composition of bacterial blebs can
20 be manipulated by acting upon/altering the expression of bacterial genes encoding products present in or associated with bacterial outer-membrane blebs (outer membrane proteins or OMPs). The production of blebs using a method of genetic modification to increase, decrease or render conditional the expression of one or more genes encoding outer-membrane components are also included in the scope of this
25 invention.

For clarity, the term "expression cassette" will refer herein to all the genetic elements necessary to express a gene or an operon and to produce and target the corresponding protein(s) of interest to outer-membrane blebs, derived from a given bacterial host. A non-exhaustive list of these features includes control elements
30 (transcriptional and/or translational), protein coding regions and targeting signals, with appropriate spacing between them. Reference to the insertion of promoter sequences means, for the purposes of this invention, the insertion of a sequence with

at least a promoter function, and preferably one or more other genetic regulatory elements comprised within an expression cassette. Moreover, the term "integrative cassette" will refer herein to all the genetic elements required to integrate a DNA segment in given bacterial host. A non-exhaustive list of these features includes a delivery vehicle (or vector), with recombinogenic regions, and selectable and counter selectable markers.

Again for the purpose of clarity, the terms 'engineering a bacterial strain to produce less of said antigen' refers to any means to reduce the expression of an antigen of interest, relative to that of the non-modified (i.e., naturally occurring) bleb such that expression is at least 10% lower than that of the non-modified bleb. Preferably it is at least 50% lower. "Stronger promoter sequence" refers to a regulatory control element that increases transcription for a gene encoding antigen of interest. "Upregulating expression" refers to any means to enhance the expression of an antigen of interest, relative to that of the non-modified (i.e., naturally occurring) bleb. It is understood that the amount of 'upregulation' will vary depending on the particular antigen of interest but will not exceed an amount that will disrupt the membrane integrity of the bleb. Upregulation of an antigen refers to expression that is at least 10% higher than that of the non-modified bleb. Preferably it is at least 50% higher. More preferably it is at least 100% (2 fold) higher.

Aspects of the invention relate to individual methods for making improved engineered blebs, to a combination of such methods, and to the bleb compositions made as a result of these methods. Another aspect of the invention relates to the genetic tools used in order to genetically modify a chosen bacterial strain in order to extract improved engineered blebs from said strain.

The engineering steps of the processes of the invention can be carried out in a variety of ways known to the skilled person. For instance, sequences (e.g. promoters or open reading frames) can be inserted, and promoters/genes can be disrupted by the technique of transposon insertion. For instance, for upregulating a gene's expression, a strong promoter could be inserted via a transposon up to 2 kb upstream of the gene's initiation codon (more preferably 200-600 bp upstream, most preferably approximately 400 bp upstream). Point mutation or deletion may also be used (particularly for down-regulating expression of a gene).

Such methods, however, may be quite unstable or uncertain, and therefore it is preferred that the engineering step [particularly for processes a), b), c), d), e), h) and i) as described below] is performed via a homologous recombination event. Preferably, the event takes place between a sequence (a recombinogenic region) of at least 30 nucleotides on the bacterial chromosome, and a sequence (a second recombinogenic region) of at least 30 nucleotides on a vector transformed within the strain. Preferably the regions are 40-1000 nucleotides, more preferably 100-800 nucleotides, most preferably 500 nucleotides). These recombinogenic regions should be sufficiently similar that they are capable of hybridising to one another under highly stringent conditions (as defined later).

Recombination events may take place using a single recombinogenic region on chromosome and vector, or via a double cross-over event (with 2 regions on chromosome and vector). In order to perform a single recombination event, the vector should be a circular DNA molecule. In order to perform a double recombination event, the vector could be a circular or linear DNA molecule (see Figure 7). It is preferable that a double recombination event is employed and that the vector used is linear, as the modified bacterium so produced will be more stable in terms of reversion events. Preferably the two recombinogenic regions on the chromosome (and on the vector) are of similar (most preferably the same) length so as to promote double cross-overs. The double cross-over functions such that the two recombinogenic regions on the chromosome (separated by nucleotide sequence 'X') and the two recombinogenic regions on the vector (separated by nucleotide sequence 'Y') recombine to leave a chromosome unaltered except that X and Y have interchanged. The position of the recombinogenic regions can both be positioned upstream or downstream of, or may flank, an open reading frame of interest. These regions can consist of coding, non-coding, or a mixture of coding and non-coding sequence. The identity of X and Y will depend on the effect desired. X may be all or part of an open reading frame, and Y no nucleotides at all, which would result in sequence X being deleted from the chromosome. Alternatively Y may be a strong promoter region for insertion upstream of an open reading frame, and therefore X may be no nucleotides at all.

Suitable vectors will vary in composition depending what type of recombination event is to be performed, and what the ultimate purpose of the

recombination event is. Integrative vectors used to deliver region Y can be conditionally replicative or suicide plasmids, bacteriophages, transposons or linear DNA fragments obtained by restriction hydrolysis or PCR amplification. Selection of the recombination event is selected by means of selectable genetic marker such as
5 genes conferring resistance to antibiotics (for instance kanamycin, erythromycin, chloramphenicol, or gentamycin), genes conferring resistance to heavy metals and/or toxic compounds or genes complementing auxotrophic mutations (for instance *pur*, *leu*, *met*, *aro*).

10

Process a) and f) – Down regulation/Removal of Variable and non-protective immunodominant antigens

Many surface antigens are variable among bacterial strains and as a consequence are protective only against a limited set of closely related strains. An
15 aspect of this invention covers the reduction in expression, or, preferably, the deletion of the gene(s) encoding variable surface protein(s) which results in a bacterial strain producing blebs which, when administered in a vaccine, have a stronger potential for cross-reactivity against various strains due to a higher influence exerted by conserved proteins (retained on the outer membranes) on the vaccinee's immune system.
20 Examples of such variable antigens include: for *Neisseria* - pili (PilC) which undergoes antigenic variations, PorA, Opa, TbpB, FrpB; for *H. influenzae* - P2, P5, pilin, IgA1-protease; and for *Moraxella* - CopB, OMP106.

Other types of gene that could be down-regulated or switched off are genes which, *in vivo*, can easily be switched on (expressed) or off by the bacterium. As outer
25 membrane proteins encoded by such genes are not always present on the bacteria, the presence of such proteins in the bleb preparations can also be detrimental to the effectiveness of the vaccine for the reasons stated above. A preferred example to down-regulate or delete is *Neisseria* Opc protein. Anti-Opc immunity induced by an Opc containing bleb vaccine would only have limited protective capacity as the
30 infecting organism could easily become Opc⁺. *H. influenzae* HgpA and HgpB are other examples of such proteins.

In process a), these variable or non-protective genes are down-regulated in expression, or terminally switched off. This has the above-mentioned surprising advantage of concentrating the immune system on better antigens that are present in low amounts on the outer surface of blebs.

5 The strain can be engineered in this way by a number of strategies including transposon insertion to disrupt the coding region or promoter region of the gene, or point mutations or deletions to achieve a similar result. Homologous recombination may also be used to delete a gene from a chromosome (where sequence X comprises part (preferably all) of the coding sequence of the gene of interest). It may additionally
10 be used to change its strong promoter for a weaker (or no) promoter (where nucleotide sequence X comprises part (preferably all) of the promoter region of the gene, and nucleotide sequence Y comprises either a weaker promoter region [resulting in a decreased expression of the gene(s)/operon(s) of interest], or no promoter region). In this case it is preferable for the recombination event to occur within the region of the
15 chromosome 1000 bp upstream of the gene of interest.

Alternatively, Y may confer a conditional transcriptional activity, resulting in a conditional expression of the gene(s)/operon(s) of interest (down-regulation). This is useful in the expression of molecules that are toxic to or not well supported by the bacterial host.

20 Most of the above-exemplified proteins are integral OMPs and their variability may be limited only to one or few of their surface exposed loops. Another aspect of this invention [process g)] covers the deletion of DNA regions coding for these surface exposed loops which leads to the expression of an integral OMP containing conserved surface exposed loops. Surface exposed loops of *H. influenzae* P2 and P5
25 are preferred examples of proteins that could be transformed into cross-reactive antigens by using such a method. Again, homologous recombination is a preferred method of performing this engineering process.

Process b) - Promoter delivery and modulation:

30 A further aspect of the invention relates to modifying the composition of blebs by altering *in situ* the regulatory region controlling the expression of gene(s) and/or operon(s) of interest. This alteration may include partial or total replacement of the

endogenous promoter controlling the expression of a gene of interest, with one conferring a distinct transcriptional activity. This distinct transcriptional activity may be conferred by variants (point mutations, deletions and/or insertions) of the endogenous control regions, by naturally occurring or modified heterologous promoters or by a combination of both. Such alterations will preferably confer a transcriptional activity stronger than the endogenous one (introduction of a strong promoter), resulting in an enhanced expression of the gene(s)/operon(s) of interest (up-regulation). Such a method is particularly useful for enhancing the production of immunologically relevant Bleb components such as outer-membrane proteins and lipoproteins (preferably conserved OMPs, usually present in blebs at low concentrations).

Typical strong promoters that may be integrated in *Neisseria* are *porA* [SEQ ID NO: 24], *porB* [SEQ ID NO:26], *lgtF*, *Opa*, *p110*, *lst*, and *hpuAB*. *PorA* and *PorB* are preferred as constitutive, strong promoters. It has been established (Example 9) that the *PorB* promoter activity is contained in a fragment corresponding to nucleotides -1 to -250 upstream of the initiation codon of *porB*. In *Moraxella*, it is preferred to use the *ompH*, *ompG*, *ompE*, *OmpB1*, *ompB2*, *ompA*, *OMPCD* and *Omp106* promoters, and in *H. influenzae*, it is preferred to integrate the *P2*, *P4*, *P1*, *P5* and *P6* promoters.

Using the preferred double cross-over homologous recombination technology to introduce the promoter in the 1000 bp upstream region, promoters can be placed anywhere from 30-970 bp upstream of the initiation codon of the gene to be up-regulated. Although conventionally it is thought the promoter region should be relatively close to the open reading frame in order to obtain optimal expression of the gene, the present inventors have surprisingly found that placement of the promoter further away from the initiation codon results in large increases in expression levels. Thus it is preferred if the promoter is inserted 200-600 bp from the initiation codon of the gene, more preferably 300-500 bp, and most preferably approximately 400 bp from the initiation ATG.

Process c) - Bleb components produced conditionally

The expression of some genes coding for certain bleb components is carefully regulated. The production of the components is conditionally modulated and depends upon various metabolic and/or environmental signals. Such signals include, for example, iron-limitation, modulation of the redox potential, pH and temperature variations, nutritional changes. Some examples of bleb components known to be produced conditionally include iron-regulated outer-membrane proteins from *Neisseiria* and *Moraxella* (for instance TbpB, LbpB), and substrate-inducible outer-membrane porins. The present invention covers the use of the genetic methods described previously (process a) or b)) to render constitutive the expression of such molecules. In this way, the influence of environmental signal upon the expression of gene(s) of interest can be overcome by modifying/replacing the gene's corresponding control region so that it becomes constitutively active (for instance by deleting part [preferably all] or the repressive control sequence – e.g. the operator region), or inserting a constitutive strong promoter. For iron regulated genes the *fur* operator may be removed. Alternatively, process i) may be used to deliver an additional copy of the gene/operon of interest in the chromosome which is placed artificially under the control of a constitutive promoter.

20

Processes d), and e) - Detoxification of LPS

The toxicity of bleb vaccines presents one of the largest problems in the use of blebs in vaccines. A further aspect of the invention relates to methods of genetically detoxifying the LPS present in Blebs. Lipid A is the primary component of LPS responsible for cell activation. Many mutations in genes involved in this pathway lead to essential phenotypes. However, mutations in the genes responsible for the terminal modifications steps lead to temperature-sensitive (*htrB*) or permissive (*msbB*) phenotypes. Mutations resulting in a decreased (or no) expression of these genes result in altered toxic activity of lipid A. Indeed, the non-lauroylated (*htrB* mutant) or non-myristoylated (*msbB* mutant) lipid A are less toxic than the wild-type lipid A. Mutations in the lipid A 4'-kinase encoding gene (*lpxK*) also decreases the toxic activity of lipid A.

30

Process d) thus involves either the deletion of part (or preferably all) of one or more of the above open reading frames or promoters. Alternatively, the promoters could be replaced with weaker promoters. Preferably the homologous recombination techniques described above are used to carry out the process.

5 The sequences of the *htrB* and *msbB* genes from *Neisseria meningitidis* B, *Moraxella catarrhalis*, and *Haemophilus influenzae* are additionally provided for this purpose.

LPS toxic activity could also be altered by introducing mutations in genes/loci involved in polymyxin B resistance (such resistance has been correlated with addition
10 of aminoarabinose on the 4' phosphate of lipid A). These genes/loci could be *pmrE* that encodes a UDP-glucose dehydrogenase, or a region of antimicrobial peptide-resistance genes common to many enterobacteriaceae which could be involved in aminoarabinose synthesis and transfer. The gene *pmrF* that is present in this region encodes a dolicol-phosphate manosyl transferase (Gunn J.S., Kheng, B.L., Krueger J.,
15 Kim K., Guo L., Hackett M., Miller S.I. 1998. *Mol. Microbiol.* 27: 1171-1182).

Mutations in the PhoP-PhoQ regulatory system, which is a phospho-relay two component regulatory system (f. i. PhoP constitutive phenotype, PhoP^c), or low Mg⁺⁺ environmental or culture conditions (that activate the PhoP-PhoQ regulatory system) lead to the addition of aminoarabinose on the 4'-phosphate and 2-hydroxymyristate
20 replacing myristate (hydroxylation of myristate). This modified lipid A displays reduced ability to stimulate E-selectin expression by human endothelial cells and TNF- α secretion from human monocytes.

Process e) involves the upregulation of these genes using a strategy as described above (strong promoters being incorporated, preferably using homologous
25 recombination techniques to carry out the process).

Alternatively, rather than performing any such mutation, a polymyxin B resistant strain could be used as a vaccine production strain (in conjunction with one or more of the other processes of the invention), as blebs from such strains also have reduced LPS toxicity (for instance as shown for meningococcus - van der Ley, P,
30 Hamstra, HJ, Kramer, M, Steeghs, L, Petrov, A and Poolman, JT. 1994. *In*: Proceedings of the ninth international pathogenic *Neisseria* conference. The Guildhall, Winchester, England).

As a further alternative (and further aspect of the invention) the inventors provide a method of detoxifying a Gram-negative bacterial strain comprising the step of culturing the strain in a growth medium containing 0.1mg-100g of aminoarabinose per litre medium.

5 As a further still alternative, synthetic peptides that mimic the binding activity of polymyxin B (described below) may be added to the Bleb preparation in order to reduce LPS toxic activity (Rustici, A, Velucchi, M, Faggioni, R, Sironi, M, Ghezzi, P, Quataert, S, Green, B and Porro M 1993. *Science* 259: 361-365; Velucchi, M, Rustici, A, Meazza, C, Villa, P, Ghezzi, P and Porro, M. 1997. *J. Endotox. Res.* 4:).

10 Process f) - Anchoring homologous or heterologous proteins to outer-membrane blebs whilst reducing the toxicity of LPS

A further aspect of this invention covers the use of genetic sequences encoding polymyxin B peptides (or analogues thereof) as a means to target fusion proteins to the outer-membrane. Polymyxin B is a cyclic peptide composed of non tRNA-
15 encoded amino acids (produced by Gram-positive actinomycetal organisms) that binds very strongly to the Lipid A part of LPS present in the outer-membrane. This binding decreases the intrinsic toxicity of LPS (endotoxin activity). Peptides mimicking the structure of Polymyxin B and composed of canonical (tRNA encoded) amino acids have been developed and also bind lipid A with a strong affinity. These
20 peptides have been used for detoxifying LPS. One of these peptides known as SAEP-2 (Nterminus-Lys-Thr-Lys-Cys-Lys-Phe-Leu-Lys-Lys-Cys-Cterminus) was shown to be very promising in that respect (Molecular Mapping and detoxifying of the Lipid A binding site by synthetic peptides (1993). Rustici, A., Velucchi, M., Faggioni, R., Sironi, M., Ghezzi, P., Quataert, S., Green, B. and M. Porro. *Science* 259, 361-365).
25

The present process f) of the invention provides an improvement of this use. It has been found that the use of DNA sequences coding for the SEAP-2 peptide (or derivatives thereof), fused genetically to a gene of interest (encoding for instance a T cell antigen or a protective antigen that is usually secreted such as a toxin, or a
30 cytosolic or periplasmic protein) is a means for targeting the corresponding recombinant protein to the outer-membrane of a preferred bacterial host (whilst at the same time reducing the toxicity of the LPS).

This system is suitable for labile proteins which would not be directly exposed to the outside of the bleb. The bleb would therefore act as a delivery vehicle which would expose the protein to the immune system once the blebs had been engulfed by T-cells. Alternatively, the genetic fusion should also comprise a signal peptide or transmembrane domain such that the recombinant protein may cross the outer membrane for exposure to the host's immune system.

This targeting strategy might be of particular interest in the case of genes encoding proteins that are not normally targeted to the outer-membrane. This methodology also allows the isolation of recombinant blebs enriched in the protein of interest. Preferably, such a peptide targeting signal allows the enrichment of outer membrane blebs in one or several proteins of interest, which are naturally not found in that given subcellular localization. A non exhaustive list of bacteria that can be used as a recipient host for such a production of recombinant blebs includes *Neisseria meningitidis*, *Neisseiria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, and *Chlamydia pneumoniae*.

Although it is preferred that the gene for the construct is engineered into the chromosome of the bacterium [using process i)], an alternative preferred embodiment is for SAEP-2-tagged recombinant proteins to be made independently, and attached at a later stage to a bleb preparation.

A further embodiment is the use of such constructs in a method of protein purification. The system could be used as part of an expression system for producing recombinant proteins in general. The SAEP-2 peptide tag can be used for affinity purification of the protein to which it is attached using a column containing immobilised lipid A molecules.

25

Process h) - Cross-reactive polysaccharides

The isolation of bacterial outer-membrane blebs from encapsulated Gram-negative bacteria often results in the co-purification of capsular polysaccharide. In some cases, this "contaminant" material may prove useful since polysaccharide may enhance the immune response conferred by other bleb components. In other cases however, the presence of contaminating polysaccharide material in bacterial bleb preparations may prove detrimental to the use of the blebs in a vaccine. For instance,

30

it has been shown at least in the case of *N. meningitidis* that the serogroup B capsular polysaccharide does not confer protective immunity and is susceptible to induce an adverse auto-immune response in humans. Consequently, process h) of the invention is the engineering of the bacterial strain for bleb production such that it is free of capsular polysaccharide. The blebs will then be suitable for use in humans. A particularly preferred example of such a bleb preparation is one from *N. meningitidis* serogroup B devoid of capsular polysaccharide.

This may be achieved by using modified bleb production strains in which the genes necessary for capsular biosynthesis and/or export have been impaired. Inactivation of the gene coding for capsular polysaccharide biosynthesis or export can be achieved by mutating (point mutation, deletion or insertion) either the control region, the coding region or both (preferably using the homologous recombination techniques described above). Moreover, inactivation of capsular biosynthesis genes may also be achieved by antisense over-expression or transposon mutagenesis. A preferred method is the deletion of some or all of the *Neisseria meningitidis* *cps* genes required for polysaccharide biosynthesis and export. For this purpose, the replacement plasmid pMF121 (described in Frosh et al.1990, *Mol. Microbiol.* 4:1215-1218) can be used to deliver a mutation deleting the *cpsCAD* (+ *galE*) gene cluster. Alternatively the *siaD* gene could be deleted, or down-regulated in expression (the meningococcal *siaD* gene encodes alpha-2,3-sialyltransferase, an enzyme required for capsular polysaccharide and LOS synthesis). Such mutations may also remove host-similar structures on the saccharide portion of the LPS of the bacteria.

Process i) - Delivery of one or more further copies of a Gene and/or operon in a host chromosome, or delivery of a heterologous gene and/or operon in a host chromosome.

An efficient strategy to modulate the composition of a Bleb preparation is to deliver one or more copies of a DNA segment containing an expression cassette into the genome of a Gram-negative bacterium. A non exhaustive list of preferred bacterial species that could be used as a recipient for such a cassette includes *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia pneumoniae*. The gene(s) contained in the expression cassette may be homologous (or

endogenous) (i.e. exist naturally in the genome of the manipulated bacterium) or heterologous (i.e. do not exist naturally in the genome of the manipulated bacterium). The reintroduced expression cassette may consist of unmodified, "natural" promoter/gene/operon sequences or engineered expression cassettes in which the promoter region and/or the coding region or both have been altered. A non-exhaustive list of preferred promoters that could be used for expression includes the promoters *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *lst*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*, the promoters *p2*, *p5*, *p4*, *ompF*, *p1*, *ompH*, *p6*, *hin47* from *H. influenzae*, the promoters *ompH*, *ompG*, *ompCD*, *ompE*, *ompB1*, *ompB2*, *ompA* of *M. catarrhalis*, the promoter λ pL, *lac*, *tac*, *araB* of *Escherichia coli* or promoters recognized specifically by bacteriophage RNA polymerase such as the *E. coli* bacteriophage T7. A non-exhaustive list of preferred genes that could be expressed in such a system includes *Neisseria* NspA, Omp85, PilQ, TbpA/B complex, Hsf, PldA, HasR; *Chlamydia* MOMP, HMWP; *Moraxella* OMP106, HasR, PilQ, OMP85, PldA; *Bordetella pertussis* FHA, PRN, PT.

In a preferred embodiment of the invention the expression cassette is delivered and integrated in the bacterial chromosome by means of homologous and/or site specific recombination. Integrative vectors used to deliver such genes and/or operons can be conditionally replicative or suicide plasmids, bacteriophages, transposons or linear DNA fragments obtained by restriction hydrolysis or PCR amplification. Integration is preferably targeted to chromosomal regions dispensable for growth *in vitro*. A non exhaustive list of preferred loci that can be used to target DNA integration includes the *porA*, *porB*, *opa*, *opc*, *rmp*, *omp26*, *lecA*, *cps*, *lgtB* genes of *Neisseria meningitidis* and *Neisseria gonorrhoeae*, the *P1*, *P5*, *hmw1/2*, *IgA-protease*, *fimE* genes of NTHi; the *lecA1*, *lecA2*, *omp106*, *uspA1*, *uspA2* genes of *Moraxella catarrhalis*. Alternatively, the expression cassette used to modulate the expression of bleb component(s) can be delivered into a bacterium of choice by means of episomal vectors such as circular/linear replicative plasmids, cosmids, phasmids, lysogenic bacteriophages or bacterial artificial chromosomes. Selection of the recombination event can be selected by means of selectable genetic marker such as genes conferring resistance to antibiotics (for instance kanamycin, erythromycin, chloramphenicol, or gentamycin), genes conferring resistance to heavy metals and/or

toxic compounds or genes complementing auxotrophic mutations (for instance *pur*, *leu*, *met*, *aro*).

Heterologous Genes - Expression of foreign proteins in outer-membrane blebs

5 Outer-membrane bacterial blebs represent a very attractive system to produce, isolate and deliver recombinant proteins for vaccine, therapeutic and/or diagnostic uses. A further aspect of this invention is in respect of the expression, production and targeting of foreign, heterologous proteins to the outer-membrane, and the use of the bacteria to produce recombinant blebs.

10 A preferred method of achieving this is via a process comprising the steps of: introducing a heterologous gene, optionally controlled by a strong promoter sequence, into the chromosome of a Gram-negative strain by homologous recombination. Blebs may be made from the resulting modified strain.

 A non-exhaustive list of bacteria that can be used as a recipient host for
15 production of recombinant blebs includes *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia pneumoniae*. The gene expressed in such a system can be of viral, bacterial, fungal, parasitic or higher eukaryotic origin.

 A preferred application of the invention includes a process for the expression
20 of *Moraxella*, *Haemophilus* and/or *Pseudomonas* outer-membrane proteins (integral, polytopic and/or lipoproteins) in *Neisseria meningitidis* recombinant blebs. The preferable integration loci are stated above, and genes that are preferably introduced are those that provide protection against the bacterium from which they were isolated. Preferred protective genes for each bacterium are described below.

25 Further preferred applications are: blebs produced from a modified *Haemophilus influenzae* strain where the heterologous gene is a protective OMP from *Moraxella catarrhalis*; and blebs produced from a modified *Moraxella catarrhalis* strain where the heterologous gene is a protective OMP from *Haemophilus influenzae* (preferred loci for gene insertion are given above, and preferred protective antigens
30 are described below).

 A particularly preferred application of this aspect is in the field of the prophylaxis or treatment of sexually-transmitted diseases (STDs). It is often

difficult for practitioners to determine whether the principal cause of a STD is due to gonococcus or *Chlamydia trachomatis* infection. These two organisms are the main causes of salpingitis – a disease which can lead to sterility in the host. It would therefore be useful if a STD could be vaccinated against or treated with a combined vaccine effective against disease caused by both organisms. The Major Outer Membrane Protein (MOMP) of *C. trachomatis* has been shown to be the target of protective antibodies. However, the structural integrity of this integral membrane protein is important for inducing such antibodies. In addition, the epitopes recognised by these antibodies are variable and define more than 10 serovars. The previously described aspect of this invention allows the proper folding of one or more membrane proteins within a bleb outer membrane preparation. The engineering of a gonococcal strain expressing multiple *C. trachomatis* MOMP serovars in the outer membrane, and the production of blebs therefrom, produces a single solution to the multiple problems of correctly folded membrane proteins, the presentation of sufficient MOMP serovars to protect against a wide spectrum of serovars, and the simultaneous prophylaxis/treatment of gonococcal infection (and consequently the non-requirement of practitioners to initially decide which organism is causing particular clinical symptoms – both organisms can be vaccinated against simultaneously thus allowing the treatment of the STD at a very early stage). Preferred loci for gene insertion in the gonococcal chromosome are give above. Other preferred, protective *C. trachomatis* genes that could be incorporated are HMWP, PmpG and those OMPs disclosed in WO 99/28475.

Targeting of heterologous proteins to outer-membrane blebs:

The expression of some heterologous proteins in bacterial blebs may require the addition of outer-membrane targeting signal(s). The preferred method to solve this problem is by creating a genetic fusion between a heterologous gene and a gene coding for a resident OMP as a specific approach to target recombinant proteins to blebs. Most preferably, the heterologous gene is fused to the signal peptides sequences of such an OMP.

Neisserial bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i) when carried out on a Neisserial strain, including gonococcus, and meningococcus (particularly *N. meningitidis* B):

5 NspA (WO 96/29412), Hsf-like (WO 99/31132), Hap (PCT/EP99/02766), PorA, PorB, OMP85 (WO 00/23595), PilQ (PCT/EP99/03603), PldA (PCT/EP99/06718), FrpB (WO 96/31618), TbpA (US 5,912,336), TbpB, FrpA/FrpC (WO 92/01460), LbpA/LbpB (PCT/EP98/05117), FhaB (WO 98/02547), HasR (PCT/EP99/05989), lipo02 (PCT/EP99/08315), Tbp2 (WO 99/57280), MltA (WO 99/57280), and ctrA

10 (PCT/EP00/00135). They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via process a): PorA, PorB, PilC, TbpA, TbpB, LbpA, LbpB, Opa, and Opc.

One or more of the following genes are preferred for downregulation via

15 process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process e): pmrA, pmrB, pmrE, and pmrF.

Preferred repressive control sequences for process c) are: the *fur* operator region (particularly for either or both of the TbpB or LbpB genes); and the DtxR

20 operator region.

One or more of the following genes are preferred for downregulation via process h): galE, siaA, siaB, siaC, siaD, ctrA, ctrB, ctrC, and ctrD.

Pseudomonas aeruginosa bleb preparations

25 One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): PcrV, OprF, OprI. They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

30 *Moraxella catarrhalis* bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): OMP106 (WO 97/41731 & WO

96/34960), HasR (PCT/EP99/03824), PilQ (PCT/EP99/03823), OMP85 (PCT/EP00/01468), lipo06 (GB 9917977.2), lipo10 (GB 9918208.1), lipo11 (GB 9918302.2), lipo18 (GB 9918038.2), P6 (PCT/EP99/03038), ompCD, CopB (Helminen ME, et al (1993) Infect. Immun. 61:2003-2010), D15 (PCT/EP99/03822),
 5 OmplA1 (PCT/EP99/06781), Hly3 (PCT/EP99/03257), LbpA and LbpB (WO 98/55606), TbpA and TbpB (WO 97/13785 & WO 97/32980), OmpE, UspA1 and UspA2 (WO 93/03761), and Omp21. They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via
 10 process a): CopB, OMP106, OmpB1, TbpA, TbpB, LbpA, and LbpB.

One or more of the following genes are preferred for downregulation via process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process
 15 e): pmrA, pmrB, pmrE, and pmrF.

Haemophilus influenzae bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): D15 (WO 94/12641), P6 (EP 281673), TbpA, TbpB, P2, P5 (WO 94/26304), OMP26 (WO 97/01638), HMW1,
 20 HMW2, HMW3, HMW4, Hia, Hsf, Hap, Hin47, and Hif (all genes in this operon should be upregulated in order to upregulate pilin). They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via process a): P2, P5, Hif, IgA1-protease, HgpA, HgpB, HMW1, HMW2, Hxu, TbpA,
 25 and TbpB.

One or more of the following genes are preferred for downregulation via process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process
 30 e): pmrA, pmrB, pmrE, and pmrF.

Vaccine Formulations

A preferred embodiment of the invention is the formulation of the bleb preparations of the invention in a vaccine which may also comprise a pharmaceutically acceptable excipient.

5 The manufacture of bleb preparations from any of the aforementioned modified strains may be achieved by any of the methods well known to a skilled person. Preferably the methods disclosed in EP 301992, US 5,597,572, EP 11243 or US 4,271,147 are used. Most preferably, the method described in Example 8 is used.

Vaccine preparation is generally described in Vaccine Design ("The subunit and adjuvant approach" (eds Powell M.F. & Newman M.J.) (1995) Plenum Press New York).

The bleb preparations of the present invention may be adjuvanted in the vaccine formulation of the invention. Suitable adjuvants include an aluminium salt such as aluminum hydroxide gel (alum) or aluminium phosphate, but may also be a
15 salt of calcium (particularly calcium carbonate), iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatised polysaccharides, or polyphosphazenes.

Suitable Th1 adjuvant systems that may be used include, Monophosphoryl lipid A, particularly 3-de-O-acylated monophosphoryl lipid A, and a combination of
20 monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL) together with an aluminium salt. An enhanced system involves the combination of a monophosphoryl lipid A and a saponin derivative particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in WO96/33739. A
25 particularly potent adjuvant formulation involving QS21 3D-MPL and tocopherol in an oil in water emulsion is described in WO95/17210 and is a preferred formulation.

The vaccine may comprise a saponin, more preferably QS21. It may also comprise an oil in water emulsion and tocopherol. Unmethylated CpG containing oligo nucleotides (WO 96/02555) are also preferential inducers of a TH1 response and
30 are suitable for use in the present invention.

The vaccine preparation of the present invention may be used to protect or treat a mammal susceptible to infection, by means of administering said vaccine via

systemic or mucosal route. These administrations may include injection *via* the intramuscular, intraperitoneal, intradermal or subcutaneous routes; or *via* mucosal administration to the oral/alimentary, respiratory, genitourinary tracts. Thus one aspect of the present invention is a method of immunizing a human host against a disease caused by infection of a gram-negative bacteria, which method comprises administering to the host an immunoprotective dose of the bleb preparation of the present invention.

The amount of antigen in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each dose will comprise 1-100 μ g of protein antigen, preferably 5-50 μ g, and most typically in the range 5 - 25 μ g.

An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects may receive one or several booster immunisations adequately spaced.

Ghost or Killed Whole cell vaccines

The inventors envisage that the above improvements to bleb preparations and vaccines can be easily extended to ghost or killed whole cell preparations and vaccines (with identical advantages). The modified Gram-negative strains of the invention from which the bleb preparations are made can also be used to make ghost and killed whole cell preparations. Methods of making ghost preparations (empty cells with intact envelopes) from Gram-negative strains are well known in the art (see for example WO 92/01791). Methods of killing whole cells to make inactivated cell preparations for use in vaccines are also well known. The terms 'bleb preparations' and 'bleb vaccines' as well as the processes described throughout this document are therefore applicable to the terms 'ghost preparation' and 'ghost vaccine', and 'killed whole cell preparation' and 'killed whole cell vaccine', respectively, for the purposes of this invention.

Combinations of methods a) – i)

It may be appreciated that one or more of the above processes may be used to produce a modified strain from which to make improved bleb preparations of the invention. Preferably one such process is used, more preferably two or more (2, 3, 4, 5, 6, 7, 8 or 9) of the processes are used in order to manufacture the bleb vaccine. As each additional method is used in the manufacture of the bleb vaccine, each improvement works in conjunction with the other methods used in order to make an optimised engineered bleb preparation.

A preferred meningococcal (particularly *N. meningitidis* B) bleb preparation comprises the use of processes a), b), d) and/or e), and h). Such bleb preparations are safe (no structures similar to host structures), non-toxic, and structured such that the host immune response will be focused on high levels of protective (and preferably conserved) antigens. All the above elements work together in order to provide an optimised bleb vaccine.

Similarly for *M. catarrhalis* and non-typeable *H. influenzae*, preferred bleb preparations comprise the use of processes a), b), and d) and/or e).

A further aspect of the invention is thus an immuno-protective and non-toxic Gram-negative bleb, ghost, or killed whole cell vaccine suitable for paediatric use.

By paediatric use it is meant use in infants less than 4 years old.

By immunoprotective it is meant that at least 40% (and preferably 50, 60, 70, 80, 90 and 100%) of infants seroconvert (4-fold increase in bactericidal activity [the dilution of antisera at which 50% of bacteria die – see for example PCT/EP98/05117]) against a set of heterologous strains to be selected from the major clonal groups known. For meningococcus B these strains should have a different PorA type from the bleb production strain, and should preferably be 2, 3, 4 or, most preferably, all 5 of strains H44/76, M97/252078, BZ10, NGP165 and CU385. For non-typeable *H. influenzae*, the strains should preferably be 2, 3, 4 or, most preferably, all 5 of strains 3224A, 3219C, 3241A, 640645, and A840177. For *M. catarrhalis*, the strains should preferably be 2, 3, 4 or, most preferably, all 5 of strains ATCC 43617, 14, 358, 216 and 2926.

By non-toxic it is meant that there is a significant (2-4 fold, preferably 10 fold) decrease of endotoxin activity as measured by the well-known LAL and pyrogenicity assays.

5 Vaccine Combinations

A further aspect of the invention are vaccine combinations comprising the bleb preparations of the invention with other antigens which are advantageously used against certain disease states. It has been found that blebs are particularly suitable for formulating with other antigens, as they advantageously have an adjuvant effect on the
10 antigens they are mixed with.

In one preferred combination, the meningococcus B bleb preparations of the invention are formulated with 1, 2, 3 or preferably all 4 of the following meningococcal capsular polysaccharides which may be plain or conjugated to a protein carrier: A, C, Y or W. Such a vaccine may be advantageously used as a global
15 meningococcus vaccine. Rather than use the meningococcus B bleb preparations of the invention, it is also envisaged that the formulation could alternatively contain wild-type meningococcus B bleb preparations from 2 or more (preferably several) strains belonging to several subtype/serotypes (for instance chosen from P1.15, P1.7,16, P1.4, and P1.2).

20 In a further preferred embodiment, the meningococcus B bleb preparations of the invention [or the aforementioned mix of 2 or more wild-type meningococcus B bleb preparations], preferably formulated with 1, 2, 3 or all 4 of the plain or conjugated meningococcal capsular polysaccharides A, C, Y or W, are formulated with a conjugated *H. influenzae* b capsular polysaccharide, and one or more plain or
25 conjugated pneumococcal capsular polysaccharides. Optionally, the vaccine may also comprises one or more protein antigens that can protect a host against *Streptococcus pneumoniae* infection. Such a vaccine may be advantageously used as a global meningitis vaccine.

The pneumococcal capsular polysaccharide antigens are preferably selected
30 from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F and 33F (most preferably from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F).

Preferred pneumococcal proteins antigens are those pneumococcal proteins which are exposed on the outer surface of the pneumococcus (capable of being recognised by a host's immune system during at least part of the life cycle of the pneumococcus), or are proteins which are secreted or released by the pneumococcus.

5 Most preferably, the protein is a toxin, adhesin, 2-component signal transducer, or lipoprotein of *Streptococcus pneumoniae*, or fragments thereof. Particularly preferred proteins include, but are not limited to: pneumolysin (preferably detoxified by chemical treatment or mutation) [Mitchell *et al.* Nucleic Acids Res. 1990 Jul 11; 18(13): 4010 "Comparison of pneumolysin genes and proteins from *Streptococcus*

10 *pneumoniae* types 1 and 2.", Mitchell *et al.* Biochim Biophys Acta 1989 Jan 23; 1007(1): 67-72 "Expression of the pneumolysin gene in *Escherichia coli*: rapid purification and biological properties.", WO 96/05859 (A. Cyanamid), WO 90/06951 (Paton *et al.*), WO 99/03884 (NAVA)]; PspA and transmembrane deletion variants thereof (US 5804193 - Briles *et al.*); PspC and transmembrane deletion variants

15 thereof (WO 97/09994 - Briles *et al.*); PsaA and transmembrane deletion variants thereof (Berry & Paton, Infect Immun 1996 Dec;64(12):5255-62 "Sequence heterogeneity of PsaA, a 37-kilodalton putative adhesin essential for virulence of *Streptococcus pneumoniae*"); pneumococcal choline binding proteins and transmembrane deletion variants thereof; CbpA and transmembrane deletion variants

20 thereof (WO 97/41151; WO 99/51266); Glyceraldehyde-3-phosphate - dehydrogenase (Infect. Immun. 1996 64:3544); HSP70 (WO 96/40928); PcpA (Sanchez-Beato *et al.* FEMS Microbiol Lett 1998, 164:207-14); M like protein, SB patent application No. EP 0837130; and adhesin 18627, SB Patent application No. EP 0834568. Further preferred pneumococcal protein antigens are those disclosed in WO

25 98/18931, particularly those selected in WO 98/18930 and PCT/US99/30390.

In a further preferred embodiment, the *Moraxella catarrhalis* bleb preparations of the invention are formulated with one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against non-typeable *H. influenzae* infection. Optionally, the vaccine may also comprise one

30 or more protein antigens that can protect a host against *Streptococcus pneumoniae* infection. The vaccine may also optionally comprise one or more antigens that can protect a host against RSV and/or one or more antigens that can protect a host against

influenza virus. Such a vaccine may be advantageously used as a global otitis media vaccine.

Preferred non-typeable *H. influenzae* protein antigens include Fimbrin protein (US 5766608) and fusions comprising peptides therefrom (eg LB1 Fusion) (US 5843464 - Ohio State Research Foundation), OMP26, P6, protein D, TbpA, TbpB, Hia, Hmw1, Hmw2, Hap, and D15.

Preferred influenza virus antigens include whole, live or inactivated virus, split influenza virus, grown in eggs or MDCK cells, or Vero cells or whole flu virosomes (as described by R. Gluck, Vaccine, 1992, 10, 915-920) or purified or recombinant proteins thereof, such as HA, NP, NA, or M proteins, or combinations thereof.

Preferred RSV (Respiratory Syncytial Virus) antigens include the F glycoprotein, the G glycoprotein, the HN protein, or derivatives thereof.

In a still further preferred embodiment, the non-typeable *H. influenzae* bleb preparations of the invention are formulated with one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against *M. catarrhalis* infection. Optionally, the vaccine may also comprise one or more protein antigens that can protect a host against *Streptococcus pneumoniae* infection. The vaccine may also optionally comprise one or more antigens that can protect a host against RSV and/or one or more antigens that can protect a host against influenza virus. Such a vaccine may be advantageously used as a global otitis media vaccine.

Nucleotide sequences of the invention

A further aspect of the invention relates to the provision of new nucleotide sequences which may be used in the processes of the invention. Specific upstream regions from various genes from various strains are provided which can be used in, for instance, processes a), b), d) and h). In addition, coding regions are provided for performing process d).

General method for the analysis of the non-coding flanking region of a bacterial gene, and its exploitation for modulated expression of the gene in blebs

The non-coding flanking regions of a specific gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the Shine-Dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences.

This sequence information allows the modulation of the natural expression of the gene in question. The upregulation of the gene expression may be accomplished by altering the promoter, the Shine-Dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modifications. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation includes, but is not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be done *in vivo* by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random mutagenesis, or site-directed replacement, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the

wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome.

In process b), for example, the expression of a gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream
5 sequence of the gene, in vitro modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.

In other preferred examples, the described approaches can be used to generate
10 recombinant bacterial strains with improved characteristics for vaccine applications, as described above. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, and strains with modulated expression of immunodominant proteins.

15
SEQ ID NO:2-23, 25, 27-38 are all Neisserial upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each. SEQ ID NO: 39-62 are all *M. catarrhalis* upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each.
20 SEQ ID NO: 63-75 are all *H. influenzae* upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each. All of these can be used in genetic methods (particularly homologous recombination) for up-regulating, or down-regulating the open reading frames to which they are associated (as described before). SEQ ID NO: 76-81 are the coding regions for the
25 HtrB and MsbB genes from Neisseria, *M. catarrhalis*, and *Haemophilus influenzae*. These can be used in genetic methods (particularly homologous recombination) for down-regulating (in particular deleting) part (preferably all) of these genes [process d)].

Another aspect of the invention is thus an isolated polynucleotide sequence
30 which hybridises under highly stringent conditions to at least a 30 nucleotide portion of the nucleotides in SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof. Preferably the isolated sequence should be long enough to perform homologous

recombination with the chromosomal sequence if it is part of a suitable vector – namely at least 30 nucleotides (preferably at least 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 nucleotides). More preferably the isolated polynucleotide should comprise at least 30 nucleotides (preferably at least 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 nucleotides) of SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof.

As used herein, highly stringent hybridization conditions include, for example, 6X SSC, 5X Denhardt, 0.5% SDS, and 100 µg/mL fragmented and denatured salmon sperm DNA hybridized overnight at 65 °C and washed in 2X SSC, 0.1% SDS one time at room temperature for about 10 minutes followed by one time at 65 °C for about 15 minutes followed by at least one wash in 0.2X SCC, 0.1% SDS at room temperature for at least 3-5 minutes.

A further aspect is the use of the isolated polynucleotide sequences of the invention in performing a genetic engineering event (such as transposon insertion, or site specific mutation or deletion, but preferably a homologous recombination event) within 1000 bp upstream of a Gram-negative bacterial chromosomal gene in order to either increase or decrease expression of the gene. Preferably the strain in which the recombination event is to take place is the same as the strain from which the upstream sequences of the invention were obtained. However, the meningococcus A, B, C, Y and W and gonococcus genomes are sufficiently similar that upstream sequence from any of these strains may be suitable for designing vectors for performing such events in the other strains. This is may also be the case for *Haemophilus influenzae* and non-typeable *Haemophilus influenzae*.

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The
5 examples are illustrative, but do not limit the invention.

Example 1: Construction of a *Neisseria meningitidis* serogroup B strain lacking capsular polysaccharides.

10 The plasmid pMF121 (Frosch et al., 1990) has been used to construct a *Neisseria meningitidis* B strain lacking the capsular polysaccharide. This plasmid contains the flanking regions of the gene locus coding for the biosynthesis pathway of the group B polysaccharide (B PS), and the erythromycin resistance gene. Deletion of the B PS resulted in loss of expression of the group B capsular polysaccharide as well as a
15 deletion in the active copy of *galE* leading to the synthesis of galactose deficient LPS.

Strain transformation:

Neisseria meningitidis B H44/76 strain (B:15:P1.7, 16;Los 3,7,9) was selected for transformation. After an overnight CO₂ incubation on MH plate (without
20 erythromycin), cells were collected in liquid MH containing 10 mM MgCl₂ (2 ml were used per MH plate) and diluted up to an OD of 0.1 (550 nm). To this 2 ml solution, 4 µl of the plasmid pMF121 stock solution (0.5 µg/ml) were added for a 6 hours incubation period at 37°C (with shaking). A control group was done with the same amount of *Neisseria meningitidis* B bacteria, but without addition of plasmid.
25 After the incubation period, 100 µl of culture, as such, at 1/10, 1/100 and 1/1000 dilutions, were put in MH plates containing 5, 10, 20, 40 or 80 µg erythromycin /ml before incubation for 48 hours at 37°C.

Colony blotting:

30 After plate incubation, 20 colonies were grown and selected from the 10 and 20 µg erythromycin/ml MH plates, while there was no colony growth in the control group without plasmid transformation. The H44/76 wild type strain was unable to grow in

the selected erythromycin plates (10 to 80 µg erythromycin/ml). The day after, all the visible colonies were placed on new MH plates without erythromycin in order to let them grow. Afterwards, they were transferred onto nitrocellulose sheets (colony blotting) for presence of B polysaccharide. Briefly, colonies were blotted onto a
5 nitrocellulose sheet and rinsed directly in PBS-0.05 % Tween 20 before cell inactivation for 1 hour at 56°C in PBS-0.05% Tween 20 (diluant buffer). Afterwards, the membrane was overlaid for one hour in the diluant buffer at room temperature (RT). Then, sheets were washed again for three times 5 minutes in the diluant buffer before incubation with the anti-B PS 735 Mab (Boehringer) diluted at 1/3000 in the
10 diluant buffer for 2 hours at RT. After a new washing step (3 times 5 minutes), the monoclonal antibody was detected with a biotinylated anti-mouse Ig from Amersham (RPN 1001) diluted 500 times in the diluant buffer (one hour at RT) before the next washing step (as described above). Afterwards, sheets were incubated for one hour at RT with a solution of streptavidin-peroxidase complex diluted 1/1000 in the diluant
15 buffer. After the last three washing steps using the same method, nitrocellulose sheets were incubated for 15 min in the dark using the revelation solution (30 mg of 4-chloro-1-naphtol solution in 10 ml methanol plus 40 ml PBS and 30 mcl of H₂O₂ 37% from Merck). The reaction was stopped with a distilled water-washing step.

20 Whole cell Elisas:

Whole cell Elisas were also done using the two transformed colonies ("D" and "R") and the wild type strain (H44/76) as coated bacteria (20 µg protein/ml), and a set of different monoclonal antibodies used to characterize *Neisseria meningitidis* strains. The following Mabs were tested: anti-B PS (735 from Dr Frosch), and the other Mabs
25 from NIBSC: anti-B PS (Ref 95/750) anti-P1.7 (A-PorA, Ref 4025), anti-P1.16 (A-PorA, Ref 95/720), anti-Los 3,7,9 (A-LPS, Ref 4047), anti-Los 8 (A-LPS, Ref 4048), and anti-P1.2 (A-PorA Ref 95/696).

Microtiter plates (Maxisorp, Nunc) were coated with 100 µl of the recombinant meningococcal B cells solution overnight (ON) at 37°C at around 20
30 µg/ml in PBS. Afterwards, plates are washed three times with 300 µl of 150 mM NaCl - 0.05 % Tween 20, and were overlaid with 100 µl of PBS-0.3 % Casein and incubated for 30 min at room temperature with shaking. Plates were washed again

using the same procedure before incubation with antibodies. Monoclonal antibodies (100 µl) were used at different dilutions (as shown in Figure 2) in PBS-0.3 % Casein 0.05 % Tween 20 and put onto the microplates before incubation at room temperature for 30 min with shaking, before the next identical washing step. 100 µl of the anti-
5 mouse Ig (from rabbit, Dakopatts E0413) conjugated to biotin and diluted at 1/2000 in PBS-0.3 % Casein - 0.05 % Tween 20 were added to the wells to detect bound monoclonal antibodies. After the washing step (as before), plates were incubated with a streptavidin-peroxidase complex solution (100 µl of the Amersham RPN 1051) diluted at 1/4000 in the same working solution for 30 min at room temperature under
10 shaking conditions. After this incubation and the last washing step, plates are incubated with 100 µl of the chromogen solution (4 mg orthophenylenediamine (OPD) in 10 ml 0.1 M citrate buffer pH4.5 with 5 µl H₂O₂) for 15 min in the dark. Plates are then read at 490/620 nm using a spectrophotometer.

15 Results:

Figure 1 shows that from the 20 isolated colonies, which were able to growth on the selected medium with erythromycin, only two (the "D" and the "R") colonies were shown negative for presence of B polysaccharide. Among the others, 16 were clearly positive for B PS and still resistant to erythromycin. This indicated that they
20 integrated the plasmid into their genome, but in the wrong orientation, and keeping intact the B PS and LPS gene (no double crossing-over). Positive and negative controls were also tested on the plates, and showed that the H44/76 wild type NmB strain was clearly positive for the B polysaccharide, while meningococcus A (A1) and meningococcus C (C11) strains were clearly negative with this anti-B PS 735 Mab.
25 These results indicate that around 10 % of the selected colonies correctly integrated the plasmid in their genome by making a double crossing-over, while the other strains/colonies were obtained after a simple crossing-over, keeping the B PS and LPS genes intact and expressed.

Using whole cell Elisa, results (Figure 2 and the Table below) clearly indicate
30 that the two "D" and "R" transformants (derived from D and R colonies) can not be recognized anymore by the anti-B PS Mabs (735 and 95/750), nor by the anti-Los 3,7,9 and anti-Los 8 Mabs. However, when using specific anti-PorA Mabs, there is a

clear reaction with the anti-P1.7 and anti-P1.16 Mabs on the cells, as also observed in the wild-type strain. No reaction was observed with a non-specific anti-PorA Mab (anti-P1.2 mab). These results confirm that the PorA protein, and specifically P1.7 and P1.16 epitopes are still present after transformation, while B polysaccharide and
 5 Los 3,7,9 and Los 8 epitopes (LPS) were not.

Table : Specificities of the monoclonal antibodies tested

Mabs Tested	Directed against	Result
Anti-B PS 735	B polysaccharide	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-B PS 95/750 from NIBSC	B PS	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-P1.7 (NIBSC)	Loop 1 of Porin A	++ on all wild type and mutants strains
Anti-P1.16 (NIBSC)	Loop 4 of Porin A	++ on all wild type and mutants strains
Anti-Los 3,7,9	LPS	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-Los 8 (NIBSC)	LPS	+/- on the wild type strain (-) on the "D" and "R" mutants
Anti-P1.2 (NIBSC)	Anti-Porin A Sero-subtype 1.2	(-) on all wild type and mutants strains

10 **Example 2: Construction of versatile gene delivery vectors (the pCMK series) targeting integration in the *porA* locus of *Neisseria meningitidis*.**

A plasmid allowing homologous recombination and stable integration of foreign DNA in the *porA* locus of *Neisseria meningitidis* was constructed. This
 15 delivery vector (genes, operons and/or expression cassettes) is useful for constructing *Neisseria meningitidis* strains producing recombinant, improved blebs. Typically, such a vector contains at least: (1) a plasmid backbone replicative in *E. coli* but not in *Neisseria meningitidis* (a suicide plasmid), (2) at least one, but preferably two regions

of homology for targeting the integration in a chromosomal locus such as *porA*, (3) Efficient transcriptional (promoter, regulatory region and terminator) and translational (optimised ribosome binding site and initiation codon) signals functional in *Neisseria meningitidis*, (4) a multiple cloning site and (5) selectable gene(s) allowing the
 5 maintenance of the plasmid in *E. coli* and the selection of integrants in *Neisseria meningitidis*. Additional elements include, for example, uptake sequences to facilitate the entry of foreign DNA in *Neisseria meningitidis*, and counter selectable markers such as *sacB*, *rpsL*, *gltS* to enhance the frequency of double cross-over events.

A schematic drawing of the vector constructed in this example and designated
 10 pCMK is represented in Figure 3. Its corresponding complete nucleotide sequence is shown in SEQ. ID NO:1. pCMK derives from a pSL1180 backbone (PharmaciaBiotech, Sweeden), a high copy-number plasmid replicative in *E. coli*, harbouring the *bla* gene (and thereby conferring resistance to ampicillin). In addition to this, pCMK functionally contains two *porA* flanking regions (*porA*5' and *porA*3' containing a transcription terminator) necessary for homologous recombination, a
 15 selectable marker conferring resistance to kanamycin, two uptake sequences, a *porA/lacO* chimeric promoter repressed in the *E.coli* host expressing *lacIq* but transcriptionally active in *Neisseria meningitidis*, and a multiple cloning site (5 sites present: *NdeI*, *KpnI*, *NheI*, *PstI* and *SphI*) necessary for the insertion of foreign
 20 DNA in pCMK.

pCMK was constructed as follows. The *porA*5' and *porA*3' recombinogenic regions, the *porA/lacO* promoter were PCR amplified using the oligonucleotides listed in the table below, cloned in pTOPO and sequenced. These DNA fragments were successively excised from pTOPO and recloned in pSL1180. The kanamycin
 25 resistance cassette was excised from pUC4K (PharmaciaBiotech, Sweeden) and was introduced between the *porA*5' flanking region and the *porA/lacO* promoter region.

Table: Oligonucleotides used in this work

Oligonucleotides	Sequence	Remark(s)
PorA5' Fwd	5'-CCC AAG CTT GCC GTC TGA ATA CAT CCC GTC ATT CCT CA-3'	<i>Hind</i> III cloning site Uptake sequence (_)
PorA5' Rev	5'-CGA TGC TCG CGA CTC CAG AGA CCT CGT GCG GGC C-3'	<i>Nru</i> I cloning site
PorA3' Fwd	5'-GGA AGA TCT GAT TAA ATA GGC GAA AAT ACC AGC TAC GA-3'	<i>Bgl</i> II cloning site Stop codons (_)
PorA3' Rev	5'-GCC GAA TTC TTC AGA CGG C GC AGC AGG AAT TTA TCG G-3'	<i>Eco</i> RI cloning site Uptake sequence (_)
PoLa Rev1	5'- GAA TTG TTA TCC GCT CAC AAT TCC GGG CAA ACA CCC GAT AC-3'	
PoLa Rev2	5'-GAA TTC CAT ATG ATC GGC TTC CTT TTG TAA ATT TGA TAA AAA CCT AAA AAC ATC GAA TTG TTA TCC GCT C-3'	<i>Nde</i> I cloning site
PorAlacO Fwd	5'-AAG CTC TGC AGG AGG TCT GCG CTT GAA TTG-3'	<i>Pst</i> I cloning site
PorAlacO Rev	5'-CTT AAG GCA TAT GGG CTT CCT TTT GTA A- 3'	<i>Nde</i> I cloning site
PPA1	5'- GCG GCC GTT GCC GAT GTC AGC C-3'	
PPA2	5'-GGC ATA GCT GAT GCG TGG AAC TGC-3'	
N-full-01:	5'-GGG AAT TCC ATA TGA AAA AAG CAC TTG CCA CAC-3'	<i>Nde</i> I cloning site
Nde-NspA-3:	5'- GGA ATT CCA TAT GTC AGA ATT TGA CGC GCA C -3'	<i>Nde</i> I cloning site
PNS1	5'- CCG CGA ATT CGG AAC CGA ACA CGC CGT TCG-3'	<i>Eco</i> RI cloning site
PNS1	5'- CGT CTA GAC GTA GCG GTA TCC GGC TGC -3'	<i>Xba</i> I cloning site
PromD15-51X	5'- GGG CGA ATT CGC GGC CGC CGT CAA CGG CAC ACC CGT TG-3'	<i>Eco</i> RI and <i>Not</i> I cloning sites
PromD15-S2	5'- GCT CTA GAG CGG AAT GCG GTT TCA GAC G- 3'	<i>Xba</i> I cloning site
PNS4	5'- AGC TTT ATT TAA ATC CTT AAT TAA CGC GTC CGG AAA ATA TGC TTA TC_34	<i>Swa</i> I and <i>Pac</i> I cloning sites
PNS5	5'- AGC TTT GTT TAA ACC CTG TTC CGC TGC TTC GGC-3'	<i>Pme</i> I cloning site
D15-S4	5'- GTC CGC ATT TAA ATC CTT AAT TAA GCA GCC GGA CAG GGC GTG G-3'	<i>Swa</i> I and <i>Pac</i> I cloning sites
D15-S5	5'- AGC TTT GTT TAA AGG ATC AGG GTG TGG TCG GGC-3'	<i>Pme</i> I cloning site

Example 3: Construction of a *Neisseria meningitidis* serogroup B strain lacking both capsular polysaccharides and the major immunodominant antigen PorA.

Modulating the antigenic content of outer membrane blebs may be
5 advantageous in improving their safety and efficacy in their use in vaccines, or
diagnostic or therapeutic uses. Components such as the *Neisseria meningitidis*
serogroup B capsular polysaccharides should be removed to exclude the risk of
inducing autoimmunity (see example 1). Similarly, it is beneficial to suppress the
immunodominance of major outer-membrane antigens such as PorA, which induce
10 strain-specific bactericidal antibodies but fail to confer cross-protection. To illustrate
such an approach, we used the pCMK(+) vector to construct a *Neisseria meningitidis*
serogroup B strain lacking both capsular polysaccharides and the immunodominant
PorA outer membrane protein antigen. For this purpose, a deletion of the *porA* gene
was introduced in the H44/76 *cps*- strain, described in example 1 by means of
15 homologous recombination.

The H44/76 *cps*- strain was prepared competent and transformed with two 2µg
of supercoiled pCMK(+) plasmid DNA as described previously. Aliquot fractions of
the transformation mixture (100µl) were plated on Mueller-Hinton plates
supplemented with Kanamycin (200µg/ml) and incubated at 37°C for 24 to 48 hours.
20 Kanamycin-resistant colonies were selected, restreaked on MH-Kn and grown for an
additional 24 hours at 37°C. At that stage half of the bacterial culture was used to
prepare glycerol stocks (15 % vol./vol.) and was kept frozen at -70°C. Another
fraction (estimated to be 10⁸ bacteria) was resuspended in 15 µl of distilled water,
boiled ten minutes and used as a template for PCR screening. Two *porA* internal
25 primers named , PPA1 and PPA2, were synthesized and used to perform PCR
amplification on boiled bacterial lysates in the conditions described by the supplier
(HiFi DNA polymerase, Boehringer Mannheim, GmbH). The thermal cycling used
was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C
10min., 4°C up to recovery). Since a double crossing-over between pCMK DNA and
30 the chromosomal *porA* locus deletes the region required for #1 and #2 annealing,
clones lacking a 1170bp PCR amplification fragment were selected as *porA* deletion
mutants. These PCR results were further confirmed by analyzing in parallel, the

presence of PorA in the corresponding bacterial protein extracts. For that purpose, another aliquot of bacteria (estimated to be $5 \cdot 10^8$ bacteria) was re-suspended in 50 μ l of PAGE-SDS buffer (SDS 5%, Glycerol 30%, Beta-mercaptoethanol 15%, Bromophenol blue 0.3mg/ml, Tris-HCl 250 mM pH6.8), boiled (100°C) frozen (-20°C) / boiled (100°C) three times and was separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-PorA monoclonal antibody as described in Maniatis *et al.* As represented in Figure 4, both Coomassie and immunoblot staining confirmed that *porA* PCR negative clones do not produce detectable levels of PorA. This result confirm that the pCMK vector is functional and can be used successfully to target DNA insertion in the *porA* gene, abolishing concomitantly the production of the PorA outer membrane protein antigen.

Example 4: Up-regulation of the NspA outer membrane protein production in blebs derived from a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *porA* and *cps* genes.

Enriching bleb vesicles with protective antigens is advantageous for improving the efficiency and the coverage of outer membrane protein-based vaccines. In that context, recombinant *Neisseria meningitidis* strains lacking functional *cps* and *porA* genes were engineered so that the expressions level of the outer-membrane protein NspA was up-regulated. For that purpose, the gene coding for NspA was PCR amplified using the N01-full-*NdeI* and *NdeI*-3' oligonucleotide primers (see table in example 2). The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling done was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was digested with *NdeI* and inserted in the *NdeI* restriction site of the pCMK(+) delivery vector. Insert orientation was checked and recombinant plasmids, designed pCMK(+)-NspA, were purified at a large scale using the QIAGEN maxiprep kit and 2 μ g of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps*

genes (strain described in example 1). Integration resulting from a double crossing-over between the pCMK(+)-NspA vector and the chromosomal *porA* locus were selected using a combination of PCR and Western blot screening procedures presented in example 3.

5 Bacteria (corresponding to about $5 \cdot 10^8$ bacteria) were re-suspended in 50 μ l of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-NspA polyclonal serum. Both Coomassie (data not shown) and
10 immunoblot staining (see figure 4) confirmed that *porA* PCR negative clones do not produce detectable levels of PorA. The expression of NspA was examined in Whole-cell bacterial lysates (WCBL) or outer-membrane bleb preparations derived from NmB [*cps*-, *porA*-] or NmB [*cps*-, *porA*-, *Nspa*+]. Although no difference was observable by Coomassie staining, immunoblotting with the anti-NspA polyclonal
15 serum detected a 3-5 fold increased in the expression of NspA (with respect to the endogenous NspA level), both in WCBL and outer-membrane bleb preparations (see figure 5). This result confirm that the pCMK(+)-NspA vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins such as NspA, abolishing concomitantly the production of the PorA outer membrane protein
20 antigen.

Example 5: Up-regulation of the D15/Omp85 outer membrane protein antigen in blebs derived from a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing PorA.

25 Certain geographically isolated human populations (such as Cuba) are infected by a limited number of *Neisseria meningitidis* isolates belonging largely to one or few outer membrane protein serotypes. Since PorA is a major outer-membrane protein antigen inducing protective and strain-specific bactericidal antibodies, it is then
30 possible to confer vaccine protection using a limited number of *porA* serotypes in a vaccine. In such a context, the presence of PorA in outer membrane vesicles may be advantageous, strengthening the vaccine efficacy of such recombinant improved

blebs. Such PorA containing vaccines, however, can be improved still further by increasing the level of other cross-reactive OMPs such as omp85/D15.

In the following example, the pCMK(+) vector was used to up-regulate the expression of the Omp85/D15 outer membrane protein antigen in a strain lacking functional *cps* genes but expressing *porA*. For that purpose, the gene coding for Omp85/D15 was PCR amplified using the D15-*NdeI* and D15-*NotI* oligonucleotide primers. The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling done was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was inserted in the pTOPO cloning vector according to the manufacturer's specifications and confirmatory sequencing was performed. This Omp85/D15 DNA fragment was excised from pTOPO by restriction hydrolysis using *NdeI/NsiI* and subsequently cloned in the corresponding restriction sites of the pCMK(+) delivery vector. Recombinant plasmids, designed pCMK(+)-D15 were purified on a large scale using the QIAGEN maxiprep kit and 2 µg of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes (strain described in example 1). In order to preserve the expression of *porA*, integration resulting from a single crossing-over (either in Omp85/D15 or in *porA*) were selected by a combination of PCR and Western blot screening procedures. Kanamycin resistant clones testing positive by *porA*-specific PCR and western blot were stored at -70°C as glycerol stocks and used for further studies.

Bacteria (corresponding to about 5.10^8 bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-*porA* monoclonal antibody. As represented in Figure 6, both Coomassie and immunoblot staining confirmed that *porA* PCR positive clones produce PorA.

The expression of D15 was examined using outer-membrane bleb preparations derived from NmB [*cps*-, *porA*-] or NmB [*cps*-, *porA*+, D15+]. Coomassie detected a significant increase in the expression of D15 (with respect to the endogenous D15 level), preparations (see Figure 6). This result confirmed that the pCMK(+)-D15

vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins such as D15, without abolishing the production of the major PorA outer membrane protein antigen.

5 **Example 6: Construction of versatile promoter delivery vectors**

Rational: The rational of this approach is represented in Figure 7 and can be summarized in 7 essential steps. Some of these steps are illustrated below with the construction of Vector for up-regulating the expression of NspA and D15/Omp85.

10

Vector for up-regulating the expression of the NspA gene.

Step 1. A DNA region (997bp) located upstream from the NspA coding gene was discovered (SEQ. ID NO:2) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC
15 13090. Two oligonucleotide primers referred to as PNS1 and PNS2 (see table in example 2) were designed using this sequence and synthesized. These primers were used for PCR amplification using genomic DNA extracted from the H44/76 strain.
Step 2. The corresponding amplicons were cleaned-up using the Wizard PCR kit (Promega, USA) and submitted to digestion with the *EcoRI/XbaI* restriction enzymes
20 for 24 hours using the conditions described by the supplier (Boehringer Mannheim, Germany). The corresponding DNA fragments were gel purified and inserted in the corresponding sites of the pUC18 cloning vector. **Step 3.** Recombinant plasmids were prepared on a large scale and an aliquot fraction was used as a template for inverse PCR amplification. Inverse PCR was performed using the PNS4 and PNS5
25 oligonucleotides using the following thermal cycling conditions: 25 times (94°C 1min., 50°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). Linearized pUC 18 vectors harbouring a deletion in the NspA upstream region insert were obtained.

30 Vector for up-regulating the expression of the D15/omp85 gene.

Step 1. A DNA region (1000 bp) located upstream from the D15/omp85 coding gene was discovered (SEQ. ID NO:3) in the private Incyte PathoSeq database containing

unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. Two oligonucleotide primers referred to as PromD15-51X and PromD15-S2 (see table in example 2) were designed using this sequence and synthesized. These primers were used for PCR amplification using genomic DNA extracted from the H44/76 strain. **Step 2.** The corresponding amplicons were cleaned-up using the Wizard PCR kit (Promega, USA) and submitted to digestion with the *EcoRI/XbaI* restriction enzymes for 24 hours in the conditions described by the supplier (Boehringer Mannheim, Germany). The corresponding DNA fragments were gel purified and inserted in the corresponding sites of the pUC18 cloning vector. **Step 3.** Recombinant plasmids were prepared on a large scale and an aliquot fraction was used as a template for inverse PCR amplification. Inverse PCR was performed using the D15-S4 and D15-S5 oligonucleotides using the following thermal cycling conditions: 25 times (94°C 1min., 50°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). Linearized pUC 18 vectors harbouring a deletion in the D15/omp85 upstream region insert were obtained.

Example 7: Fermentation processes for producing recombinant blebs .

The examples listed below describe methods for producing recombinant blebs lacking either capsular polysaccharides or capsular polysaccharides and PorA. Such a procedure may be used for a wide range of *Neisseria meningitidis* recombinant strains and may be adapted over an extended scale range.

Culture media: *Neisseria meningitidis* serogroup B strains were propagated in solid (FNE 004 AA, FNE 010 AA) or liquid (FNE 008 AA) culture media. These new media for growing meningococcus are advantageously free of animal products, and are considered a further aspect of the invention.

Components	FNE 004 AA	FNE 008 AA	FNE 010 AA
Agar	18 g/L	-	18 g/L
NaCl	6 g/L	6 g/L	6 g/L
Na-Glutamate	-	1.52 g/L	-
NaH ₂ PO ₄ ·2H ₂ O	2.2 g/L	2.2 g/L	2.2 g/L
KCl	0.09 g/L	0.09 g/L	0.09 g/L
NH ₄ Cl	1.25 g/L	1.25 g/L	1.25 g/L
Glucose	5 g/L	20 g/L	5 g/L
Yeast Extract UF	-	2.5 g/L	-
Soy Pepton	5 g/L	30 g/L	5 g/L
CaCl ₂ ·2H ₂ O	0.015 g/L	-	0.015 g/L
MgSO ₄ ·7H ₂ O	0.6 g/L	0.6 g/L	0.6 g/L
Erythromycine :	0.015 g/L	-	-
Kanamycine	-	-	0.2 g/L

5 Flask cultivation of *Neisseria meningitidis* serogroup B cps- recombinant blebs: This was performed in two steps comprising preculture on solid medium followed by liquid cultivation. **Solid pre-culture** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE004AA (see above). The Petri dish was incubated at 37°C for 18 ± 2 hours. The surface growth was resuspended in 8 mL of FNE008AA (see above) supplemented with 15 mg/L of erythromycin. **Flask culture.** 2 mL of resuspended solid pre-culture were added to a 2 litre flask containing 400 mL of FNE008AA supplemented with 15 mg/L of erythromycin. The flask was placed on a shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

15

20 Batch mode cultivation of *Neisseria meningitidis* serogroup B cps- recombinant blebs: This was performed in three steps comprising preculture on solid medium, liquid cultivation and batch mode cultivation. **Solid pre-culture.** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE004AA (see above). The Petri dish was

incubated at 37°C for 18 ± 2 hours. The surface growth was resuspended in 8 mL of FNE008AA (see above) supplemented with 15 mg/L of erythromycin. **Liquid pre-culture.** 2 mL of resuspended solid pre-culture were added to one 2 liters flask containing 400 mL of FNE008AA supplemented with 15 mg/L of erythromycin. The flask was placed on a shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours. The content of the flask was used to inoculate the 20 liters fermenter. **Batch mode culture in fermenter.** The inoculum (400 mL) was added to a pre-sterilized 20 liters (total volume) fermenter containing 10 L of FNE008AA supplemented with 15 mg/L of erythromycin. The pH was adjusted to and maintained at 7.0 by the automated addition of NaOH (25% w/v) and H₃PO₄ (25% v/v). The temperature was regulated at 37°C. The aeration rate was maintained at 20 L of air / min and the dissolved oxygen concentration was maintained at 20% of saturation by the agitation speed control. The overpressure in the fermenter was maintained at 300 g/cm². After 9 ± 1 hours, the culture was in stationary phase. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

Flask cultivation of *Neisseria meningitidis* serogroup B cps-, PorA- recombinant blebs: This was performed in two steps comprising preculture on solid medium followed by liquid cultivation. **Solid pre-culture.** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE010AA (see above). The Petri dish was incubated at 37°C for 18 ± 2 hours. The surface growth was resuspended in 8 mL of FNE008AA (see above) supplemented with 200 mg/L of kanamycin. **Flask culture.** 2 mL of resuspended solid pre-culture were added to a 2 litre flask containing 400 mL of FNE008AA supplemented with 200 mg/L of kanamycin. The flask was placed on a shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

Example 8: Isolation and purification of blebs from meningococci devoid of capsular polysaccharide

5 Recombinant blebs were purified as described below. The cell paste (42gr) was suspended in 211 ml of 0.1M Tris-Cl buffer pH 8.6 containing 10 mM EDTA and 0.5% Sodium Deoxycholate (DOC). The ratio of buffer to biomass was 5/1 (V/W). The biomass was extracted by magnetic stirring for 30 minutes at room temperature. Total extract was then centrifuged at 20,000g for 30 minutes at 4°C (13,000 rpm in a
10 JA-20 rotor, Beckman J2-HS centrifuge). The pellet was discarded. The supernatant was ultracentrifuged at 125,000g for 2 hours at 4°C (40,000 rpm in a 50.2Ti rotor, Beckman L8-70M ultracentrifuge) in order to concentrate vesicles. The supernatant was discarded. The pellet was gently suspended in 25 ml of 50 mM Tris-Cl buffer pH 8.6 containing 2 mM EDTA, 1.2% DOC and 20% sucrose. After a second
15 ultracentrifugation step at 125,000g for 2 hours at 4°C, vesicles were gently suspended in 44 ml of 3% sucrose and stored at 4°C. All solutions used for bleb extraction and purification contained 0.01% thiomersalate. As illustrated in Figure 8, this procedure yields protein preparations highly enriched in outer-membrane proteins such as PorA and PorB.

20

Example 9: Identification of bacterial promoters suitable for up-regulation antigens-coding genes

The use of strong bacterial promoter elements is essential to obtain up-regulation of genes coding for outer membrane proteins. In that context, we have
25 shown previously that up-regulating the *Neisseria meningitidis* *nspA*, *hsf*, and *omp85* genes using the *porA* promoter has allowed us to isolate recombinant blebs enriched in the corresponding NspA, Hsf and Omp85 proteins. Alternatives to the *porA* promoter may be useful to obtain different levels of up-regulation, to overcome potential *porA* phase variation and/or to achieve conditional gene expression (iron-
30 regulated promoters). Here we describe a method allowing the identification of a precise transcriptional start site of strong promoter elements likely to confer high level of expression in bacteria. Since promoter regulatory elements are classically

encompassed within 200 bp upstream and 50bp downstream from the +1 site (Collado-Vides J, Magasanik B, Gralla JD, 1991, *Microbiol Rev* 55(3):371-94), the result of such an experiment allows us to identify DNA fragments of about 250 bp carrying strong promoter activities. Major outer membrane proteins such as *Neisseria meningitidis* PorA, PorB & Rmp, *Haemophilus influenzae* P1, P2, P5 & P6, *Moraxella catarrhalis* OmpCD, OmpE, as well as some cytoplasmic and/or iron regulated proteins of these bacteria possess strong promoter elements. As a validation of this general methodology, we mapped the transcriptional start site of the strong *Neisseria meningitidis* *porA* and *porB* promoters using rapid amplification of cDNA elements (5' RACE).

The principles of 5' RACE are the following: 1) Total RNA extraction using QIAGEN "RNeasy" Kit. Genomic DNA removing by DNase treatment followed by QIAGEN purification; 2) mRNA reverse transcription with a *porA* specific 3' end primer (named *porA3*). Expected cDNA size: 307 nt. RNA removing by alkaline hydrolysis; 3) Ligation of a single-stranded DNA oligo anchor (named DT88) to the 3' end of the cDNA using T4 RNA ligase. Expected product size: 335 nt. Amplification of the anchor-ligated cDNA using a combination of hemi-nested PCR; 4) PCR amplification of the anchor-ligated cDNA using a complementary-sequence anchor primer as the 5' end primer (named DT89) and a 3'end primer (named p1-2) which is internal to the 3'end RT primer *porA3*. Expected product size: 292 bp; 5) PCR amplification of previous PCR products using DT89 as 5'end primer and p1-1 as 3'end primer (internal to p1-2). Expected product size: 211bp; and 6) Sequencing with p1-1 primer (expected products size can be calculated because *porA* transcription start site is known: 59 nt before the "ATG" translation start site).

Experimental procedure

Total RNA was extracted from approximately 10^9 cells of *Neisseria meningitidis* serogroup B *cps- porA+* strain. Extraction of 1 ml of a liquid culture at appropriate optical density ($OD_{600} = 1$) was performed by the QIAGEN "RNAeasy" kit according to the manufacturer's instructions. Chromosomal DNA was removed by addition of 10U of RNase-free DNase (Roche Diagnostics, Mannheim, Germany) to the 30 μ l of eluted RNA and was

incubated at 37°C for 15 min. The DNA-free RNA was purified with the same QIAGEN kit according to instructions.

Reverse transcription reactions were performed using primer porA3 and 200U of SUPERScript II reverse transcriptase (Life Technologies). The RT reactions were performed in a 50µl volume containing: 5µl of 2mM dNTP, 20 pmol of porA3 primer, 5µl of 10X SUPERScript II buffer, 9µl of 25mM MgCl₂, 4µl of 0.1M DTT, 40U of recombinant ribonuclease inhibitor and 1 µg of total RNA. The porA3 primer was annealed stepwise (70°C for 2 min, 65°C for 1 min, 60°C for 1 min, 55°C for 1 min, 50°C for 1 min, and 45°C for 1 min) before the SUPERScript II was added. The RT reaction was performed at 42°C for 30 min, followed by 5 cycles (50°C for 1 min, 53°C for 1 min and 56°C for 1 min) to destabilize RNA secondary structure. Two parallel reactions were performed with the reverse transcriptase omitted from one reaction as negative control.

The RNA was removed by alkaline hydrolysis cleavage with the addition of 1 µl of 0.5M EDTA followed by 12.5 µl of 0.2 M NaOH before incubation at 68°C for 5 min. The reactions were neutralized by adding 12.5 µl of 1 M Tris-HCl (pH7.4) and precipitated by the addition of 20 µg of glycogen (Roche Molecular Biochemicals, Mannheim, Germany), 5 µl of 3 M sodium acetate and 60 µl of isopropanol. Both samples were resuspended in 20 µl of 10:1 TE (10 mM Tris-HCl, pH 7.4; 1 mM EDTA, pH8).

T4 RNA ligase was used to anchor a 5'-phosphorylated, 3'end ddCTP-blocked anchor oligonucleotide DT88 (see table below). Two parallel ligations were performed overnight at room temperature with each containing: 1.3 µl of 10X RNA ligase buffer (Roche Molecular Biochemicals), 0.4 µM DT88, 10 µl of either cDNA or RT control sample and 3 U of T4 RNA ligase. As negative controls, a second set of ligation reactions was performed, omitting the T4 RNA ligase. The resulting ligation-reaction mixtures were used directly without purification in the subsequent PCR.

The anchor-ligated cDNA was amplified using a combination of hemi-nested and hot-started PCR approaches to increase specificity and product yield. Four separate first-round PCR were performed on the RT/ligase reaction and controls in a 30 µl volume, each containing: 3 µl of 10X Taq Platinum buffer, 3µl of 25 mM MgCl₂, 1 µl of 10mM dNTP, 10 pmol of each primers and 1 µl of corresponding RNA ligation reaction. The PCR were hot started by the use of Taq Platinum (Life Technologies) DNA polymerase (2U added). The first ligation-anchored PCR (LA-PCR) was performed using 10 pmol of both the anchor-specific primer DT89 and the transcript-specific primer p1-2 (see table below) which is internal to the 3' end RT primer porA3. The PCR was performed using an initial 95°C for a 5 min step (for DNA polymerase activation) followed by 10 cycles at 95°C for 10 s and 70°C

for 1 min (reducing one degree per cycle), 15 cycles at 95°C for 10 s and 60°C for 1 min. The second hemi-nested LA-PCR was performed under the same conditions using primer DT89 and the p1-2 internal primer, together with 10 pmol of p1-1 (see table below) and 1 µl of first-round PCR. Amplification products were purified using the QIAGEN "QIAquick PCR purification" kit according to manufacturer instructions before submitted to sequencing.

The CEQ™ Dye Terminator Cycle Sequencing kit (Beckman, France) was used to sequence the RACE PCR products using 10 pmol of primer p1-1. Sequencing reactions were performed according to the provided instructions and sequencing products were analyzed by the Ceq2000 DNA Analysis System (Beckman-Coulter).

DT88	5' GAAGAGAAGGTGGAAATGGCGTTTTGGC 3'
DT89	5' CCAAAACGCCATTTCCACCTTCTCTTC 3'
porA3	5' CCAAATCCTCGCTCCCCTTAAAGCC 3'
p1-2	5' CGCTGATTTTCGTCCTGATGCGGC 3'
p1-1	5' GGTCAATTGCGCCTGGATGTTCTCTG 3'

Results for the *Neisseria meningitidis* porA promoter

The start of transcription for *Neisseria meningitidis* serogroup B (strain H44/76) *porA*-mRNA was mapped 59 bp upstream of the ATG start codon using the described 5'-RACE procedure. This result confirms the mapping performed by primer extension and published by van der Ende *et al* (1995). This result supports that a DNA fragment containing nucleotides -9 to -259 with regard to the *porA* ATG is suitable for driving strong gene expression in *Neisseria meningitidis* and possibly in other bacterial species such as *Haemophilus*, *Moraxella*, *Pseudomonas*.

Results for the *Neisseria meningitidis* porB promoter

The same experimental strategy has been applied for *Neisseria meningitidis* serogroup B (strain H44/76) *porB* transcription start site mapping. Primers listed in the table below correspond to 3' end RT primer (porB3), transcript-specific primer that is internal to the *porB3* (porB2) and internal to the *porB2* (porB1). *porB3*, *porB2* and *porB1* are respectively located 265 bp, 195 bp and 150 bp downstream the ATG start codon.

porB1	5' GGTAGCGGTTGTAACCTTCAGTAACTT 3'
porB2	5' GTCTTCTTGGCCTTTGAAGCCGATT 3'
porB3	5' GGAGTCAGTACCGGCGATAGATGCT 3'

Using *porB1* and DT89 primers a ~200 bp PCR amplicon was obtained by performing 5' - RACE mapping. Since *porB1* is located 150 bp from the *porB* ATG start

codon, this result supports that the *porB* transcriptional start site is located about 50 bp (+/- 30 bp) upstream of the *porB* ATG.

The exact nucleotide corresponding to transcription initiation is presently being determined by DNA sequencing. The above PCR result supports that a DNA fragment containing nucleotides -1 to -250 with regard to the *porB* ATG start codon is suitable for driving strong gene expression in *Neisseria meningitidis* and possibly in other bacterial species such as *Haemophilus*, *Moraxella*, *Pseudomonas*.

Example 10: Up-regulation of the *N. meningitidis* serogroup B *Omp85* gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the D15/*Omp85* gene by the strong *porA* promoter in order to up-regulate the production of the D15/*Omp85* antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (1000 bp) located upstream from the *D15/omp85* coding gene was discovered (SEQ ID NO:3) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. The main steps of this procedure are represented in Figure 9. Briefly, a DNA fragment (1000bp) covering nucleotides -48 to -983 with respect to the *D15/Omp85* gene start codon (ATG) was PCR amplified using oligonucleotides **ProD15-51X** (5'-GGG CGA ATT CGC GGC CGC CGT CAA CGG CAC ACC GTT G-3') and **ProD15-52** (5'-GCT CTA GAG CGG AAT GCG GTT TCA GAC G-3') containing *EcoRI* and *XbaI* restriction sites (underlined) respectively. This fragment was submitted to restriction and inserted in pUC18 plasmid restricted with the same enzymes. The construct obtained was submitted to *in vitro* mutagenesis using the Genome Priming system (using the pGPS2 donor plasmid) commercialized by New England Biolabs (MA, USA). Clones having inserted a mini-transposon (derived from Tn7 and harboring a chloramphenicol resistance gene) were selected. One clone containing a mini-transposon insertion located in the *D15/Omp85* 5' flanking region, 401 bp downstream from the *EcoRI* site was isolated and used for further studies. This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992), Biotechniques12: 528-534) in order to (i) delete a repeated DNA sequence (Tn7R) generated by the transposition process, (ii) insert meningococcal uptake sequences

required for transformation, and (iii) insert suitable restriction sites allowing cloning of foreign DNA material such as promoters. The circle PCR was performed using the **TnRD15-KpnI/XbaI + US** (5'-CGC CGG TAC CTC TAG AGC CGT CTG AAC CAC TCG TGG ACA ACC C-3') & **TnR03Cam(KpnI)** (5'-CGC CGG TAC CGC CGC TAA CTA TAA CGG TC-3') oligonucleotides containing uptake sequences and suitable restriction sites (*KpnI* and *XbaI*) underlined. The resulting PCR fragment was gel-purified, digested with *Asp718* (isoschizomer of *KpnI*) and ligated to a 184bp DNA fragment containing the *porA* promoter and generated by PCR using the **PorA-01** (5'-CGC CGG TAC CGA GGT CTG CGC TTG AAT TGT G-3') and **PorA02** (5'-CGC CGG TAC CTC TAG ACA TCG GGC AAA CAC CCG-3') oligonucleotides containing *KpnI* restriction sites. Recombinant clones carrying a *porA* promoter inserted in the correct orientation (transcription proceeding in the *EcoRI* to *XbaI* direction) were selected and used to transform a strain of *Neisseria meningitidis* serogroup B lacking capsular polysaccharide (*cps*-) and one of the major outer membrane proteins - PorA (*porA*-). Recombinant *Neisseria meningitidis* clones resulting from a double crossing over event (PCR screening using oligonucleotides **Cam-05** (5'-GTA CTG CGA TGA GTG GCA GG-3') & **proD15-52**) were selected on GC medium containing 5µg/ml chloramphenicol and analyzed for D15/Omp85 expression. As represented in Figure 10, the production of D15/Omp85 was significantly increased in the total protein extracts of *Nm* strains resulting from promoter replacement, when compared to parental strain (*cps*-). This result was also observed when analyzing outer-membrane blebs prepared from the same strains (see Figure 17). These results are attributable to the replacement of the endogenous *D15* promoter by the strong *porA* promoter. In addition, it was surprisingly found that expression, where the *porA* promoter was introduced approximately 400 bp upstream of the initiator codon, was approximately 50 times greater than when the promoter was placed approximately 100 bp upstream. Altogether, these experiments support that the promoter replacement strategy works and allows the up-regulation of the synthesis of integral outer-membrane proteins in outer-membrane blebs.

30

Certain geographically isolated human populations (such as Cuba) are infected by a limited number of *Neisseria meningitidis* isolates belonging largely to one or

few outer membrane protein serotypes. Since PorA is a major outer-membrane protein antigen which can induce protective and strain-specific bactericidal antibodies, it may be possible to confer vaccine protection in such a population using a limited number of porA serotypes. Moreover, PorA may interact with or stabilize some other outer
5 membrane proteins. In this context, the presence of PorA in outer membrane vesicles may be advantageous, strengthening the vaccine efficacy of such recombinant improved blebs.

For such a reason, it may be desirable to up-regulate the expression of D15/Omp85 outer membrane protein in a *Neisseria meningitidis* serogroup B strain
10 lacking functional *cps* genes but expressing PorA. Genomic DNA was extracted from the recombinant *Neisseria meningitidis* serogroup B *cps*-, *porA*-, D15/Omp85+ strain using the QIAGEN Genomic Tips 100-G kit. 10µgr of this material was linearized and used to transform *Neisseria meningitidis* serogroup B *cps*- following a classical transformation protocol. Recombinant *Neisseria* were obtained on GC agar plates
15 containing 5µgr/ml chloramphenicol.

Integrations resulting from a double crossing-over upstream of the *D15* gene were screened by PCR as described previously. As homologous recombinations can occur everywhere in the chromosome, a second PCR screening was performed to control the integrity of the *porA* locus in the recombinant strain. For this purpose,
20 internal *porA* primers PPA1 (5- GCG GCC GTT GCC GAT GTC AGC C -3') and PpA2 (5- GGC ATA GCT GAT GCG TGG AAC TGC -3') were used in a PCR screening experiment. The amplification of an 1170bp fragment confirms the presence of the *porA* gene in the recombinant bacteria.

Recombinant bacteria (corresponding to about $5 \cdot 10^8$ bacteria) can be re-suspended in 50 µl of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times
25 and then separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels can then be stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed either with an anti-porA monoclonal antibody or with an anti-D15/Omp85 rabbit polyclonal antibody. Analysis of outer-membrane blebs prepared from the same
30 strains can also be performed.

Example 11: Up-regulation of the Hsf protein antigen in a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing PorA.

As described above, in certain countries, the presence of PorA in outer membrane vesicles may be advantageous, and can strengthen the vaccine efficacy of recombinant improved blebs. In the following example, we have used a modified pCMK(+) vector to up-regulate the expression of the Hsf protein antigen in a strain lacking functional *cps* genes but expressing PorA. The original pCMK(+) vector contains a chimeric *porA/lacO* promoter repressed in *E. coli* host expressing *lacIq* but transcriptionally active in *Neisseria meningitidis*. In the modified pCMK(+), the native *porA* promoter was used to drive the transcription of the *hsf* gene. The gene coding for Hsf was PCR amplified using the HSF 01-*NdeI* and HSF 02-*NheI* oligonucleotide primers, presented in the table below. Because of the sequence of the HSF 01-*NdeI* primer the Hsf protein expressed will contain two methionine residues at the 5' end. The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling was the following: 25 times (94°C 1min., 48°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was subsequently cloned in the corresponding restriction sites of pCMK(+) delivery vector. In this recombinant plasmid, designed pCMK(+)-Hsf, we deleted the *lacO* present in the chimeric *porA/lacO* promoter by a recombinant PCR strategy (See Figure 12). The pCMK(+)-Hsf plasmid was used as a template to PCR amplify 2 separate DNA fragments:

-fragment 1 contains the *porA* 5' recombinogenic region, the Kanamycin resistance gene and the *porA* promoter. Oligonucleotide primers used, RP1(*SacII*) and RP2, are presented in the table below. RP1 primer is homologous to the sequence just upstream of the *lac* operator.

-fragment 2 contains the Shine-Dalgarno sequence from the *porA* gene, the *hsf* gene and the *porA* 3' recombinogenic region. Oligonucleotide primers used, RP3 and RP4(*ApaI*), are presented in the table below. RP3 primer is homologous to the sequence just downstream of the *lac* operator. The 3' end of fragment 1 and the 5' end of fragment 2 have 48 bases overlapping. 500ng of each PCR (1 and 2) were used for a final PCR reaction using primers RP1 and RP4. The final amplicon obtained was subcloned in pSL1180 vector restricted with *SacII* and *ApaI*. The modified plasmid

pCMK(+)-Hsf was purified at a large scale using the QIAGEN maxiprep kit and 2 µg of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes (the strain described in example 1). In order to preserve the expression of *porA*, integration resulting from a single crossing-over was selected by a combination of PCR and Western blot screening procedures. Kanamycin resistant clones testing positive by *porA*-specific PCR and western blot were stored at -70°C as glycerol stocks and used for further studies. Bacteria (corresponding to about 5.10⁸ bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. The expression of Hsf was examined in Whole-cell bacterial lysates (WCBL) derived from *NmB* [Cps-, PorA+] or *NmB* [Cps-, PorA+, Hsf+]. Coomassie staining detected a significant increase in the expression of Hsf (with respect to the endogenous Hsf level) (See in Figure 13). This result confirms that the modified pCMK(+)-Hsf vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins, without abolishing the production of the major PorA outer membrane protein antigen.

20 Oligonucleotides used in this work

Oligonucleotides	Sequence	Remark(s)
Hsf 01-Nde	5'- GGA ATT CCA TAT GAT GAA CAA AAT ATA CCG C-3'	<i>Nde</i> I cloning site
Hsf 02-Nhe	5'-GTA GCT AGC TAG CTT ACC ACT GAT AAC CGA C-3'	<i>Nhe</i> I cloning site
GFP-mut-Asn	5'-AAC TGC AGA ATT AAT ATG AAA GGA GAA GAA CTT TTC-3'	<i>Asn</i> I cloning site Compatible with <i>Nde</i> I
GFP-Spe	5'-GAC ATA CTA GTT TAT TTG TAG AGC TCA TCC ATG -3'	<i>Spe</i> I cloning site Compatible with <i>Nhe</i> I
RP1 (SacII)	5'- TCC CCG CGG GCC GTC TGA ATA CAT CCC GTC-3'	<i>Sac</i> II cloning site
RP2	5'-CAT ATG GGC TTC CTT TTG TAA ATT TGA GGG CAA ACA CCC GAT ACG TCT TCA-3'	

RP3	5'-AGA CGT ATC GGG TGT TTG CCC TCA AAT TTA CAA AAG GAA GCC CAT ATG -3'	
RP4(ApaI)	5'-GGG TAT TCC GGG CCC TTC AGA CGG CGC AGC AGG -3'	<i>ApaI</i> cloning site

Example 12: Expression of the Green Fluorescent Protein in a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing *PorA*.

5 In the following example, the pCMK vector was used to test the expression of a cytoplasmic heterologous protein in *Neisseria meningitidis*. The Green Fluorescent Protein was amplified from the pKen-Gfpmut2 plasmid with the primers GFP-Asn-mut2 and GFP-Spe (see table in Example 11). *AsnI* gives cohesive ends compatible with *NdeI*, *SpeI* gives cohesive ends compatible with *NheI*. The conditions used for
10 PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling was the following: 25 times (94°C 1min., 48°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was subsequently cloned in the pCMK(+) delivery vector digested with *NdeI* and *NheI* restriction enzymes. In this recombinant plasmid,
15 designed pCMK(+)-GFP, we deleted the *lacO* present in the chimeric *porA/lacO* promoter by a recombinant PCR strategy. The pCMK(+)-GFP plasmid was used as template to PCR amplify 2 separate DNA fragments:

-fragment 1 contained the *porA* 5' recombinogenic region, the Kanamycin resistance gene and the *porA* promoter. Oligonucleotide primers used, RP1(*SacII*) and
20 RP2 (see table in example 11). RP1 primer is homologous to the sequence just upstream of the *lac* operator.

-fragment 2 contains the *PorA* Shine-Dalgarno sequence, the *gfp* gene and the *porA* 3' recombinogenic region. Oligonucleotide primers used, RP3 and RP4(*ApaI*), are presented in the table in example 11. RP3 primer is homologous to the sequence
25 just downstream of the *lac* operator.

The 3'end of fragment 1 and the 5'end of fragment 2 have 48 bases overlapping. 500ng of each PCR (1 and 2) were used for a final PCR reaction using

primers RP1 and RP4. Twenty μ g of this PCR fragment were used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes.

Transformation with linear DNA is less efficient than with circular plasmid DNA but all the recombinants obtained performed a double crossing-over (confirmed by a combination of PCR and Western blot screening procedures). Kanamycin resistant clones were stored at -70°C as glycerol stocks and used for further studies. Bacteria (corresponding to about $5 \cdot 10^8$ bacteria) were re-suspended in 50 μ l of PAGE-SDS buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel.

The expression of GFP was examined in Whole-cell bacterial lysates (WCBL) derived from *NmB* [Cps-, PorA+] or *NmB* [Cps-, PorA-, GFP+]. Coomassie staining detected an expression of GFP absent in the recipient *Neisseria meningitidis* strain (see figure 14).

Example 13: Up-regulation of the *N. meningitidis* serogroup B *NspA* gene by promoter replacement

The aim of the experiment was to replace the endogenous promoter region of the *NspA* gene by the strong *porA* promoter, in order to up-regulate the production of the NspA antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (924bp) located upstream from the *NspA* coding gene was discovered (SEQ ID NO: 7) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. A DNA fragment (675bp) covering nucleotides -115 to -790 with respect to the *NspA* gene start codon (ATG) was PCR amplified using oligonucleotides PNS1' (5'-CCG CGA ATT CGA CGA AGC CGC CCT CGA C-3') and PNS2 (5'-CGT CTA GAC GTA GCG GTA TCC GGC TGC -3') containing *EcoRI* and *XbaI* restriction sites (underlined) respectively. The PCR fragment was submitted to restriction with *EcoRI* and *XbaI* and inserted in pUC18. This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992), Biotechniques12: 528-534) in order to insert meningococcal uptake sequences required for transformation, and suitable restriction sites allowing cloning of a *CmR*/*PorA* promoter cassette. The circle PCR was performed using the

BAD01-2 (5'- GGC GCC CGG GCT CGA GCT TAT CGA TGG AAA ACG CAG C-3') & **BAD02-2** (5'-GGC GCC CGG GCT CGA GTT CAG ACG GCG CGC TTA TAT AGT GGA TTA AC -3') oligonucleotides containing uptake sequences and suitable restriction sites (*Xma*I and *Xho*I) underlined. The resulting PCR fragment was gel-purified and digested with *Xho*I. The *CmR*/PorA promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers

BAD 15-2 (5'-GGC GCC CGG GCT CGA GTC TAG ACA TCG GGC AAA CAC CCG-3') & **BAD 03-2** (5'- GGC GCC CGG GCT CGA GCA CTA GTA TTA CCC TGT TAT CCC-3') oligonucleotides containing suitable restriction sites (*Xma*I, *Xba*I, *Spe*I and *Xho*I) underlined. The PCR fragment obtained was submitted to digestion and inserted in the circle PCR plasmid restricted with the corresponding enzymes. 10 µg of the recombinant plasmid were linearized and used to transform a strain of *Neisseria meningitidis* serogroup B lacking capsular polysaccharide (*cps*-) and one of the major outer membrane proteins - PorA (*porA*-). Recombinant *Neisseria meningitidis* clones resulting from a double crossing over event (PCR screening using oligonucleotides **BAD 25** (5'-GAG CGA AGC CGT CGA ACG C -3') & **BAD08** (5'-CTT AAG CGT CGG ACA TTT CC-3')) were selected on GC agar plates containing 5µg/ml chloramphenicol and analyzed for NspA expression. Recombinant bacteria (corresponding to about $5 \cdot 10^8$ bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed either with an anti-PorA monoclonal antibody or with anti-NspA polyclonal antibody (figure 17). As for Omp85, there is a surprising indication that insertion of the promoter approximately 400 bp upstream of the NspA initiation codon expresses more protein than if placed approximately 100 bp upstream.

The same recombinant pUC plasmid can be used to up-regulate the expression of NspA in a *Neisseria meningitidis* serogroup B strain lacking functional *cps* gene but still expressing PorA.

Example 14: Up-regulation of the *N. meningitidis* serogroup B *pldA* (*omplA*) gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the *pldA* (*omplA*) gene by the strong *porA* promoter in order to up-regulate the production of the PldA (OmplA1) antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (373bp) located upstream from the *pldA* coding sequence was discovered (SEQ ID NO: 18) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC 13090. This DNA contains the sequence coding for a putative *rpsT* gene. The stop codon of *rpsT* is located 169bp upstream the *pldA* ATG. To avoid the disruption of this potentially important gene, we decided to insert the *CmR/PorA* promoter cassette just upstream of the ATG of *pldA*. For that purpose, a DNA fragment of 992 bp corresponding to the the *rpsT* gene, the 169 bp intergenic sequence and the 499 first nucleotides of *pldA* gene was PCR amplified from *Neisseria meningitidis* serogroup B genomic DNA using oligonucleotides **PLA1 Amo5** (5'-GCC GTC TGA ATT TAA AAT TGC GCG TTT ACA G-3') and **PLA1 Amo3** (5'-GTA GTC TAG ATT CAG ACG GCG CAA TTT GGT TTC CGC AC -3') containing uptake sequences (underlined). PLA1 Amo3 contains also a *XbaI* restriction site. This PCR fragment was cleaned with a High Pure Kit (Roche, Mannheim, Germany) and directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to insert suitable restriction sites allowing cloning of a *CmR/PorA* promoter cassette. The circle PCR was performed using the **CIRC1-Bgl** (5'CCT AGA TCT CTC CGC CCC CCA TTG TCG -3') & either **CIRC1-XH-RBS/2** (5'-CCG CTC GAG TAC AAA AGG AAG CCG ATA TGA ATA TAC GGA ATA TGC G-3') or **CIRC2-XHO/2** (5'-CCG CTC GAG ATG AAT ATA CGG AAT -3') oligonucleotides containing suitable restriction sites (*Bgl*II and *Xho*I) underlined. The *CmR/PorA* promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') and **CM-PORA-3** (5'- CCG CTC GAG ATA AAA ACC TAA AAA CAT CGG GC-3') containing suitable restriction sites (*Bgl*II and *Xho*I) underlined. This PCR fragment was cloned

in the circle PCR plasmid obtained with primers CIRC1-Bgl and CIRC1-XH-RBS/2. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*- *porA*-) strains. Integration by double crossing-over in the upstream region of *pldA* will direct the insertion of the *porA* promoter directly upstream of the *pldA* ATG.

5 Another cassette was amplified from the genomic DNA of the recombinant *Neisseria meningitidis* serogroup B (*cps*-, *porA*-, D15/Omp85+) over-expressing D15/Omp85 by promoter replacement. This cassette contains the *cmR* gene, the *porA* promoter and 400bp corresponding to the 5' flanking sequence of the *D15/Omp85* gene. This sequence has been proven to be efficacious for up-regulation of the expression of

10 D15/Omp85 in *Neisseria* and will be tested for the up-regulation of the expression of other *Neisseria* antigens. Primers used for the amplification were **BAD 20** and **CM-PORA-D15/3** (5'- CGG CTC GAG TGT CAG TTC CTT GTG GTG C-3') containing *XhoI* restriction sites (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-Bgl and CIRC2-XHO/2. This

15 plasmid will be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*-, *porA*-) strains. Integration by double crossing-over in the upstream region of *pldA* will direct the insertion of the *porA* promoter 400bp upstream the *pldA* ATG.

Example 15: Up-regulation of the *N. meningitidis* serogroup B *tbpA* gene by promoter replacement.

20

The aim of the experiment was to replace the endogenous promoter region of the *tbpA* gene by the strong *porA* promoter, in order to up-regulate the production of the TbpA antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (731bp) located upstream from

25 the *tbpA* coding sequence was discovered (SEQ ID NO: 17) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC 13090. This DNA contains the sequence coding for TbpB antigen. The genes are organized in an operon. The *tbpB* gene will be deleted and replaced by the *CmR/porA* promoter cassette. For that purpose, a DNA fragment of 3218bp corresponding to the 509bp 5' flanking

30 region of *tbpB* gene, the 2139bp *tbpB* coding sequence, the 87bp intergenic sequence and the 483 first nucleotides of *tbpA* coding sequence was PCR amplified from *Neisseria meningitidis* serogroup B genomic DNA using oligonucleotides **BAD16** (5'-

GGC CTA GCT AGC CGT CTG AAG CGA TTA GAG TTT CAA AAT TTA TTC-3') and **BAD17** (5'-GGC CAA GCT TCA GAC GGC GTT CGA CCG AGT TTG AGC CTT TGC-3') containing uptake sequences and *NheI* and *HindIII* restriction sites (underlined). This PCR fragment was cleaned with a High Pure Kit (Boehringer Mannheim, Germany) and directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to (i) insert suitable restriction sites allowing cloning of a *CmR/PorA* promoter cassette and (ii) to delete 209bp of the 5' flanking sequence of *tbpB* and the *tbpB* coding sequence. The circle PCR was performed using the **BAD 18** (5'-TCC CCC GGG AAG ATC TGG ACG AAA AAT CTC AAG AAA CCG-3') & the **BAD 19** (5'-GGA AGA TCT CCG CTC GAG CAA ATT TAC AAA AGG AAG CCG ATA TGC AAC AGC AAC ATT TGT TCC G -3') oligonucleotides containing suitable restriction sites *XmaI*, *BglII* and *XhoI* (underlined). The *CmR/PorA* promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD21** (5'- GGA AGA TCT CCG CTC GAG ACA TCG GGC AAA CAC CCG-3') & **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') containing suitable restriction sites *XmaI*, *SpeI*, *BglII* and *XhoI* (underlined). This PCR fragment was cloned in the circle PCR plasmid. This plasmid will be used to transform *Neisseria meningitidis* serogroup B (*cps-*) and (*cps- porA-*) strains. Integration by double crossing-over in the upstream region of *tbpA* will direct the insertion of the *porA* promoter directly upstream of the *tbpA* ATG.

Example 16: Up-regulation of the *N. meningitidis* serogroup B *pilQ* gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the *pilQ* gene by the strong *porA* promoter, in order to up-regulate the production of the PilQ antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (772bp) located upstream from the *pilQ* coding gene was discovered (SEQ ID NO: 12) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC 13090. This DNA contains the sequence coding for PilP antigen. The *pilQ* gene is part of an operon we do not want to disturb, pilins being essential elements of the bacteria. The *CmR/porA* promoter

cassette was introduced upstream the *pilQ* gene following the same strategy described for the up-regulation of the expression of the *pldA* gene. For that purpose, a DNA fragment of 866 bp corresponding to the 3' part of the *pilP* coding sequence, the 18bp intergenic sequence and the 392 first nucleotides of *pilQ* gene was PCR amplified from *Neisseria* serogroup B genomic DNA using **PQ-rec5-Nhe** (5'-CTA GCT AGC GCC GTC TGA ACG ACG CGA AGC CAA AGC-3') and **PQ-rec3-Hin** (GCC AAG CTT TTC AGA CGG CAC GGT ATC GTC CGA TTC G-3') oligonucleotides containing uptake sequences and *NheI* and *HindIII* restriction sites (underlined). This PCR fragment was directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to insert suitable restriction sites allowing cloning of a *CmR*/*PorA* promoter cassette. The circle PCR was performed using the **CIRC1-PQ-Bgl** (5'-GGA AGA TCT AAT GGA GTA ATC CTC TTC TTA-3') & either **CIRC1-PQ-XHO** (5'-CCG CTC GAG TAC AAA AGG AAG CCG ATA TGA TTA CCA AAC TGA CAA AAA TC-3') or **CIRC2-PQ-X** (5'-CCG CTC GAG ATG AAT ACC AAA CTG ACA AAA ATC -3') oligonucleotides containing suitable restriction sites *Bgl*II and *Xho*I (underlined). The *CmR*/*PorA* promoter cassette was amplified from the pUC D15/*Omp85* plasmid previously described, using primers **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') and **CM-PORA-3** (5'- CCG CTC GAG ATA AAA ACC TAA AAA CAT CGG GCA AAC ACC C-3') containing suitable restriction sites *Bgl*II and *Xho*I (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-PQ-Bgl and CIRC1-PQ-XHO. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*-, *porA*-) strains. Integration by double crossing-over in the upstream region of *pilQ* will direct the insertion of the *porA* promoter directly upstream of the *pilQ* ATG.

Another cassette was amplified from the genomic DNA of the recombinant *Neisseria meningitidis* serogroup B (*cps*-, *porA*-, *D15/Omp85*+) over-expressing *D15/Omp85* by promoter replacement. This cassette contains the *cmR* gene, the *porA* promoter and 400bp corresponding to the 5' flanking sequence of the *D15/Omp85* gene. This sequence has been proven to be efficacious for up-regulation of the expression of *D15/Omp85* in *Neisseria meningitidis* and will be tested for the up-regulation of the expression of other *Neisseria* antigens. Primers used for the

amplification were **BAD 20** and **CM-PORA-D153** (5'- GGG CTC GAG TGT CAG TTC CTT GTG GTG C-3') containing *Xho*I restriction sites (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-PQ-Bgl and CIRC2-PQ-X. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*-, *porA*-) strains. Integration by double crossing-over in the upstream region of *pilQ* will direct the insertion of the *porA* promoter 400bp upstream the *pilQ* ATG.

Example 17: Construction of a *kanR/sacB* cassette for introducing "clean", unmarked mutations in the *N. meningitidis* chromosome.

The aim of the experiment is to construct a versatile DNA cassette containing a selectable marker for the positive screening of recombination in the chromosome of *Neisseria meningitidis* (ie: *kanR* gene), and a counter selectable marker to delete the cassette from the chromosome after recombination (ie: *sacB* gene). By this method, any heterologous DNA introduced during homologous recombination will be removed from the *Neisseria* chromosome.

A DNA fragment containing the *neoR* gene and the *sacB* gene expressed under the control of its own promoter was obtained by restriction of the pIB 279 plasmid (Blomfield IC, Vaughn V, Rest RF, Eisenstein BI (1991), Mol Microbiol 5:1447-57) with *Bam*HI restriction enzyme. The recipient vector was derived from plasmid pCMK, previously described. The *kanR* gene of the pCMK was deleted by restriction with enzymes *Nru*I and *Eco*RV. This plasmid was named pCMKs. The *neoR/sacB* cassette was inserted in the pCMKs at a *Bgl*III restriction site compatible with *Bam*HI ends.

E. coli harboring the plasmid is unable to grow in the presence of 2% sucrose in the culture medium, confirming the functionality of the *sacB* promoter.

This plasmid contains recombinogenic sequences allowing the insertion of the cassette at the *porA* locus in the chromosome of *Neisseria meningitidis* serogroup B. Recombinant *Neisseria* were obtained on GC agar plates containing 200µg/ml of kanamycin. Unfortunately, the *sacB* promoter was not functional in *Neisseria meningitidis*: no growth difference was observed on GC agar plates containing 2% sucrose.

A new cassette was constructed containing the *sacB* gene under the control of the *kanR* promoter. A circle PCR was performed using the plasmid pUC4K ((Amersham Pharmacia Biotech, USA)) as a template with **CIRC-Kan-Nco** (5'-CAT GCC ATG GTT AGA AAA ACT CAT CGA GCA TC-3') & **CIRC-Kan-Xba** (5'-CTA GTC TAG ATC AGA ATT GGT TAA TTG GTT G-3') oligonucleotides containing *NcoI* and *XbaI* restriction sites (underlined). The resulting PCR fragment was gel-purified, digested with *NcoI* and ligated to the *sacB* gene generated by PCR from the pIB279 plasmid with **SAC/NCO/NEW5** (5'-CAT GCC ATG GGA GGA TGA ACG ATG AAC ATC AAA AAG TTT GCA A-3') oligonucleotide containing a *NcoI* restriction site (underlined) and a RBS (bold) & **SAC/NCO/NEW3** (5'-GAT CCC ATG GTT ATT TGT TAA CTG TTA ATT GTC-3') oligonucleotide containing a *NcoI* restriction site (underlined). The recombinant *E. coli* clones can be tested for their sensitivity on agar plates containing 2% sucrose. The new *kanR/sacB* cassette can be subcloned in the pCMKs and used to transform a *Neisseria meningitidis* serogroup B *cps*- strain. The acquired sucrose sensitivity will be confirmed in *Neisseria*. The pCMKs plasmid will be used to transform the recombinant *kanR/SacB Neisseria* to delete the entire cassette inserted in the chromosome at the *porA* locus. Clean recombinant *Neisseria* will be obtained on GC agar plates containing 2% sucrose.

20

Example 18: Use of small recombinogenic sequences (43bp) to allow homologous recombination in the chromosome of *Neisseria meningitidis*.

The aim of the experiment is to use small recombinogenic sequences (43bp) to drive insertions, modifications or deletions in the chromosome of *Neisseria*. The achievement of this experiment will greatly facilitate future work, in terms of avoiding subcloning steps of homologous sequences in *E. coli* (recombinogenic sequences of 43bp can easily be added in the PCR amplification primer). The *kanR* gene was PCR amplified from plasmid pUC4K with oligonucleotides **Kan-PorA-5** (5'-GCC GTC TGA ACC CGT CAT TCC CGC GCA GGC GGG AAT CCA GTC CGT TCA GTT TCG GGA AAG CCA CGT TGT GTC-3') containing 43bp homologous to the 5' flanking sequence of *NmB porA* gene (bold) and an uptake sequence (underlined) & **Kan-PorA-3** (5'-TTC AGA CGG CGC AGC AGG AAT

30

TTA TCG GAA ATA ACT GAA ACC GAA CAG ACT AGG CTG AGG TCT
GCC TCG-3') containing 43bp homologous to the 3' flanking sequence of *NmB porA*
gene (bold) and an uptake sequence (underlined). The 1300bp DNA fragment
obtained was cloned in pGemT vector (Promega, USA). This plasmid can be used to
transform a *Neisseria meningitidis* serogroupB *cps*- strain. Recombinant *Neisseria*
will be obtained on GC plates containing 200µg/ml kanamycin. Integrations resulting
from a double crossing-over at the *porA* locus will be screened by PCR with primers
PPA1 & PPA2 as described previously.

**Example 19: Active protection of mice immunized with WT and recombinant
Neisseria meningitidis blebs**

Animals were immunised three times (IP route) with 5 µg of the different
OMVs adsorbed on Al(OH)₃ on days 0, 14 and 28. Bleedings were done on days 28
(day 14 Post II) and 35 (day 7 post III), and they were challenged on day 35 (IP
route). The challenge dose was 20 x LD₅₀ (~10⁷ CFU/mouse). Mortality rate was
monitored for 7 days after challenge.

OMVs injected were:

Group1: Cps-, PorA+ blebs

Group2: Cps-, PorA- blebs

Group3: Cps-, PorA-, NspA+ blebs

Group4: Cps-, PorA-, Omp85+ blebs

Group5: Cps-, PorA-, Hsf+ blebs

Figure 15 illustrates the pattern of these OMVs by analyzed SDS Page
(Coomassie staining).

24 hours after the challenge, there was 100% mortality (8/8) in the negative
control group (immunised with Al(OH)₃ alone) while mice immunised with the 5
different OMVs preparations were still alive (7 to 8/8 mice survived). Sickness was
also monitored during the 7 days and the mice immunised with the NSPA over-
expressed blebs appeared to be less sick than the other groups. PorA present in PorA+
blebs is likely to confer extensive protection against infection by the homologous
strain. However, protection induced by PorA- up-regulated blebs is likely to be due at
least to some extent, to the presence of increased amount of NspA, Omp85 or Hsf.

Example 20: Immunogenicity of recombinant blebs measured by whole cell & specific ELISA methods

To measure the ability of the antibodies to recognize the antigens present on the MenB cell surface, the pooled mice sera (from Example 19) were tested by whole cell ELISA (using tetracyclin inactivated cells), and titers were expressed as mid-point titers. All types of bleb antibodies induce a high whole cell Ab titer while the negative control group was clearly negative.

Blebs	WCE(H44/76) mid-point titer	
	14P2	14P3
CPS(-) PorA(+)	23849	65539
CPS(-) PorA(-)	20130	40150
CPS(-) PorA(-) NSPA(+)	8435	23846
CPS(-) PorA(-) OMP85(+)	4747	16116
CPS(-) PorA(-) HSF(+)	6964	22504
(-)	51	82

10

The specific Ab response to available recombinant HSF protein was carried out. Microplates were coated with 1 µg/ml full length HSF molecule.

The results illustrated in Figure 16 show that there was a good specific HSF response when HSF over-expressed OMVs were used to immunize mice (using purified recombinant HSF on the plates). The HSF over-expressed blebs induce a good level of specific antibodies.

15

SEQ. ID NO:1

Nucleotide sequence of the pCMK(+) vector

TCTTCCGCTTCCTCGCTCACTGACTCGCTCGCTCGGTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGT
AATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTA
5 AAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCTCGACGAGCATCACAAAATCGACGCTCAAGTCAGAGG
TGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTTCCCTGGAAGCTCCCTCGTGCCTCTCTGTTCGACCCCT
GCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCAGCTGTAGGTATCTCA
GTTCGGTGTAGGTCTGTCCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTTCAGCCGACCGCTGCGCCTTATCCGGT
AACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC
10 GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGTATCT
GCGCTCTGCTGAAGCCAGTTACCTTCGGAAGAGGTTGGTAGCTCTTGATCCGGCAAAACAAACCCGCTGGTAGCGGT
GGTTTTTTTGTGTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTCTACGGGGTC
TGACGCTCAGTGAACGAAACTCAGCTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTT
TAAATTAAGAAATGAAGTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGT
15 GAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCTGACTCCCCGTCGTGTAGATAACTACGATACG
GGAGGGCTTACCATCTGGCCCCAGTCTGCAATGATACCGCGAGACCCAGCTCACCAGCTCCAGATTTATCAGCAATAA
ACCAGCCAGCCGGAAGGGCCGAGCGCAGAAAGTGGTCTCTGCAACTTTATCCGCTCCATCCAGTCTATTAATTGTTGCCGG
GAAGCTAGAGTAAGTAGTTTCGCCAGTTAATAGTTTGCACGACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTC
GTGTTTTGGTATGGCTTCATTAGCTCCGTTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAAG
20 CGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAGTAAAGTTGGCCGAGTGTATCACTCATGTTATGGCAGCACTG
CATAATTCTCTTACTGTCTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAAGTACTCAACCAAGTCATTCTGAGAATA
GTGTATGCGCGACCGAGTTGCTCTTGCCTCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAGTGC
TCATCATTGAAAAACGTTCTTCGGGCGCAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACT
CGTGACCCAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAACAGGAAGGCAAAATGCCGC
25 AAAAAAGGGAATAAGGGCGACACGGAATGTTGAATACTCATACTCTTCTCTTTTCAATATTATTGAAGCATTATCAGG
GTTATTGTCTCATGAGCGGATACATATTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCGCA
AAAGTGCCACCTGACGTCTAAGAAACCATTAATTATCATGACATTAACCTATAAAAAATAGGCGTATCACGAGGCCCTTTCCG
TCTCGCGCGTTTCGGTGATGACGGTGAAAACTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTGTAAGCGG
ATGCCGGGAGCAGACAAGCCCGTCAGGGCGCGTCAGCGGGTGTGCGGGGTGTCGGGGCTGGCTTAACTATGCGGCATCA
30 GAGCAGATTGTACTGAGAGTGACCATATAAATTGTAACGTTAATATTTTGTAAAAATTCGCGTTAAATTTTGTAAAT
CAGCTCATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAAGAATAGCCCGAGATAGGGTTGAGTG
TTGTTCCAGTTTGAACAAGAGTCCACTATTAAAGAAGCTGGACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGC
GATGGCCCACTACGTGAACCATCACCAATCAAGTTTTTGGGGTGGAGGTGCGGTAAAGCACTAAATCGGAACCCCTAA
AGGGAGCCCCGATTTAGAGCTTGACGGGGAAGCCGCGCAACGTGCGGAGAAAGGAAGGAAGAAAGCGAAAGGAGCGG
35 GCGCTAGGGCGCTGGCAAGTGTAGCGGTACGCTGCGCGTAACCACCACACCCGCGCGCTTAATGCGCCGCTACAGGGC
GCGTACTATGGTTGCTTTGACGTATGCGGTGTGAAATACCGCACAGATGCGTAAGGAGAAAAATACCGCATCAGGCGCCAT
TCGCCATTACAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCTCTTCGCTATTACGCCAGCTGGCGAAAGGGG
ATGTGCTGCAAGGCGATTAAGTTGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGACGGCCAGTGCCAAGC
TTGCCGTCTGAATACATCCCGTCATTCCTCAAAAAACAGAAAAACCAAAATCAGAAACCTAAATCCCGTCATTCCCGCGCA
40 GCGGGAATCCAGTCCGTTTCAGTTTCGGTCATTTCGGATAAATTCCTGCTGCTTTTCATTCTAGATTCCCACTTTTCGTG
GGAATGACGGCGGAAGGGTTTTGGTTTTTTCCGATAAATCTTGAGGCATTGAAATTTAGATTCCCGCTGCGCGGGAA
TGACGGCTGTAGATGCCGATGGTCTTTATAGCGGATTACAAAAATCAGGACAAGGCGACGAAGCCGACAGTACAG
ATAGTACGGAACCGATTCACTTGGTGTTCAGCACCTTAGAGAATCGTTCTCTTTGAGCTAAGGCGAGGCAACGCCGTAC
TTGTTTTTGTAAATCCACTATAAAGTGCGCGTGTGTTTTTTATGGCGTTTTAAAAAGCCGAGACTGCATCCGGGCGAGC
45 AGCGCATCGGCCCGCACGAGGTCTCTGGAGTCGCGAGCATCAAGGGCGAATCTGACGGGGGGGGGGGAAAGCCAGTT
GTGTCTCAAAATCTCTGATGTACATTGCACAAGATAAAAAATATCATCATGAACAATAAACTGTCTGCTTACATAAA
CAGTAATACAAGGGGTGTTATGAGCCATATTCAACGGGAAACGTCTTGCTCGAGGCCGCGATTAAATCCCAACATGGATG

CTGATTATATGGGTATAAATGGGCTCGCGATAATGTCTGGGCAATCAGGTGCGACAATCTATCGATTGTATGGGAAGCCC
GATGCGCCAGAGTTGTTTCTGAAACATGGCAAAGGTAGCGTTGCCAATGATGTTACAGATGAGATGGTCAGACTAAACTG
GCTGACGGAATTTATGCTCTTCCGACCATCAAGCATTTTATCCGTACTCTCTGATGATGCATGGTTACTCACCCTGCGA
TCCCCGGGAAAACAGCATTCCAGGTATTAGAAGAATATCCTGATTACAGGTGAAAATATTGTTGATGCGCTGGCAGTGTTC
5 CTGCGCCGGTTGCATTTCGATTCTCTGTTTGTAAATGTCTCTTTTAAACAGCGATCGCGTATTTCTGCTCGCTCAGGCGCAATC
ACGAATGAATAACGGTTTGGTTGATGCGAGTGATTTTGTATGACGAGCGTAATGGCTGGCCTGTTGAACAAGTCTGGAAG
AAATGCATAAGCTTTTGCCATTCTCACCAGATTCACTCGTCACTCATGGTGATTTCTCACTTGATAACCTTATTTTGTAC
GAGGGGAAATTAATAGGTTGTATTGATGTTGGACGAGTCGGAATCGCAGACCGATACCAGGATCTTGCCATCTTATGGAA
CTGCTCTCGGTGAGTTTCTCTTCAATTACAGAAACGGCTTTTTCAAAAATATGGTATTGATAATCCTGATATGAATAAAT
10 TGCAGTTTCATTTGATGCTCGATGAGTTTCTAATCAGAATTGGTTAATTGGTTGTAACACTGGCAGAGCATTACGCTG
ACTTGACGGGACGGCGGCTTTGTGAATAAATCGAATTTTGTCTGAGTTGAAGGATCAGATCACGCATCTTCCCGACAAC
GCAGACCGTTCCGTGGCAAAGCAAAGTTCAAATCACCACCTGGTCCACCTACAACAAAGCTCTCATCAACCGTGCGTC
CCTCACTTTCTGGCTGGATGATGGGGCGATTAGGCTGATGAGTCAGCAACACCTTCTTACGAGGCAGACCTCAGC
GCCCCCCCCCTGACAGGAGTCTGCGCTTGAATTGTGTTGTAGAAACACAACGTTTGTGAAAAATAAGCTATTGTTT
15 TATATCAAAATATAATCATTTTAAAAATAAGGTTGCGGCATTTATCAGATATTTGTTCTGAAAAATGGTTTTTTCGGG
GGGGGGGTATAATTGAAGACGTATCGGGTGTGTCGGGAATTGTGAGCGGATAACAATTCGATGTTTTTAGGTTTTTA
TCAATTTTACAAAGGAAGCCCATATGCATCCTAGGCCTATTAATATTCGGAGTATACGTAGCCGGCTAACGTTAACAA
CCGTACCTCTAGAACTATAGCTAGCATGCGCAAATTTAAAGCGCTGATATCGATCGCGCGCAGATCTGATTAAATAGGC
GAAATACAGCTACGATCAAATCATCGCGCGGTTGATTATGATTTTCCAAACGCACCTCCGCCATCGTGTCTGGCGC
20 TTGGCTGAAACGCAATACCGGCATCGGCACTACACTCAAATTAATGCGCGCTCCGTGCGTTTGGCCACAAATCTAAA
TATCGGGCGGTGAAGCGGATAGCTTTGTTTTGACGGCTTCGCCTTCATTCTTTGATTGCAATCTGACTGCCAATCTGC
TTCAGCCCCAAACAAAAACCGGATACGGAAGAAAAACGGCAATAAGACAGCAAATACCGTCTGAAAGATTTTCAGACG
GTATTTTCGATTTTGGCTTGGTTTGACATATAGTGAGACCTTGGCAAAAATAGTCTGTTAACGAAAATTTGACGCATAA
AAATGCGCAAAAAATTTCAATTGCTTAAACCTTCTTAATATTGAGCAAAAAGTAGGAAAAATCAGAAAAGTTTGTGA
25 TTTTGAAAAATGAGATTGAGCATAAAATTTTAGTAACCTATGTTATTGCAAAGGTCTCGAATTGTCATTCCCACGAGGCG
GGAATCTAGTCTGTTTCGGTTTCAGTTATTTCCGATAAATCCTGCTGCGCCGTCTGAAGAATTCTGAATCATGTCATAG
CTGTTTCTGTGTGAAATTGTTATCCGCTCACAATTCACACACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGG
TGCTTAATGAGTGAGCTAACTCACATTAATGCGTTGCGCTCACTGCCCCGTTTCCAGTCGGGAAACCTGTCGTGCCAGC
TGCATTAATGAATCGGCCAACGCGCGGGAGAGCGGTTTGCATATTGGGCGC

30

SEQ. ID NO:2

Nucleotide sequence of DNA region (997 bp) up-stream from the NspA gene in the *Neisseria meningitidis* serogroup A strain Z2491.

GGAACCGAACACGCGTTTCGGTCATACGCCGCCGAAAGGTTTGGCGCAAGACGAAGCCGCCCTCGACATCGAAGACGCGG
35 TACACGGCGCGCTGGAAGCGCGGTTTGTCCACTACGAAACATCGGCTTTTGGCAAACAGCCATGCAGTGCCGCCAC
AATTTGAACTACTGGCAGTTTCGGCGATTATTTAGGCATAGGCGCGGGCGCGCACGGCAAAATTTCTATCCCGACCGCAT
CGAGCGCACCGTCCGCGCCGCCACCCCAACGACTACCTCGCCTTAATGCAAAACCGACCGAGCGAAGCCGTGCAACGCA
AAACCGTCGCGCGCGAAGATTGCGGTTTGAATTCATGATGAACGCCCTGCGCCTGACCGACGGCGTACCCACCGCGATG
TTGCAGGAGCGCACGGGCGTACCGAGTGCCAAAATCATGGCGCAAATCGAAACGGCAAGGCAAAAAGGCTGCTGGAAC
40 CGACCCCGCGTATTCGCGCCGACCGAAAAAGACGCTTGTTTTAAACGATTTGCTGCAGTGTTTTATAGTGGATTA
ACAAAAACAGTACGGCGTTGCCTCGCCTTAGCTCAAAGAGAACGATTCTCTAAGGTGCTGAAGCACCAAGTGAATCGGT
TCCGTACTATCTGTACTGTCTGCGGCTTCGTGCGCTTGTCTGATTTTGTTAATCCACTATATAAGCGCAAAACAAATCG
GCGGCGCCCGGGAAAAACCCCCGAACGCGTCCGGAATAATGCTTATCGATGGAACCGCAGCCGCATCCCCGCGCG
GCGTTTCAGACGGCACAGCGCGCGCGGAAATGTCGACGCTTAAGGCACAGACGCACAAAAACCGTATGCCTGCAC
45 CTGCAACAATCCGACAGATACCGCTGTTTTTCCAAACCGTTTGAAGTTTCAACCATCCGCGCGTGATGCCGCCACCA

CCATTTAAAGGCAACGCGCGGGTTAACGGCTTTGCCG

SEQ. ID NO:3

Nucleotide sequence of DNA region (1000 bp) up-stream from the D15/Omp85 gene in the *Neisseria meningitidis* serogroup B strain ATCC13090.

5 ACCATTGCCGCCCCGCGCGGCTTCCAAAGCGGCGACAAAATACAATCCGTCAACGGCACACCCGTTGCAGATTGGGGCAG
CGCGCAAAACCGAAATCGTCTCAACCTCGAAGCCGGCAAAGTCGCCGTCGGGTTGACACGGCATCAGGCGCGCAACCCGT
CCGACCATCGATGCCGACGGCAGCCCGAAGCCGGTAAATCGCAAAAAACCAAGGCTACATCGGACTGATGCCCTTTA
10 AAATCACAACCGTTGCCGCTGGCGTGGAAAAAGCAGCCCCGCGCAAAAAGCAGGCCTGAAACCGGGCGACAGGCTGACT
GCCGCCGACGGCAAAACCCATTACCTCATGGCAAGAATGGGCAAACTGACCCGCCAAAGCCCCGGCAAAAAATCACCCCT
GAACTACGAACGCGCCGGACAAACCCATACCGCCGACATCCGCCCGGATCTGTGCAACAGCCCGACACACCCCTGATCG
GGCGCGTCGGCCTCCGTCCGACGCGGACAGGGCGTGGGACGCGCAAATCCGCCGCGAGTACCGTCCGTCTGTTATCCGC
GCATTCCGGCATGGGCTGGGAAAAAACCGTTTCCCACTCGTGGACAACCCCTCAAATTTTTCGGCAAACTAATCAGCGGCAA
CGCCTCCGTGACGCATATTTCCGGGCGCTGACCAATTGCCGACATTGCCGACAGTCCGCCGAACCTCGGCTTGCAAAGTT
15 ATTTGGAATTTTGGCACTGGTCAGCATCAGCCTCGGCGTGTGAACCTGCTGCCCGTCCCCGTTTTGGACGGCGGCCAC
CTCGTGTTTTATACTGCCGAATGGATACGCGGCAAACTTTGGGGCAACGCGTCCAAAACATCGGTTTGCGCTTCGGGCT
TGCCCTCATGATGCTGATGATGGCGTTCGCTTCTTCAACGACGTTACCCGGTGTCTCGGTTAGATTTTACGTTTCGGAA
TGCCGTCTGAAACCGCATTCCGCACCACAAGGAACTGACA

20 SEQ. ID NO:4

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hsf-like gene from *Neisseria meningitidis*

25 ATCCCGCGCAGGCGGGAATCCAGAAACGCAACAGGAATTTATCGGAAAAACAGAAACCTCACCGCCGTCATT
CCGCAAAAGCGGGAATCTAGAAACACAAACGCGCAGGACTTTATCAGAAAAACAGAAACCCACCGCCGTCATTCCCGC
AAAAGCGGGAATCCAGACCCGTCGGCAGCGGAACTTACCGGATAAACAGTTTCTTAGATTCCAGTCTTAGATTCCCG
CTTTCGCGGGAATGACGAGATTTTAGATTATGGGAATTTATCAGGAATGATTGAATCCATAGAAAAACACAGGAATCTA
TCAGAAAAACAGAAACCCACCGCGTCATTCCCGCGCAGGCGGGAATCCAGAAACACAAACGCGGCGAGGACTTTATCGG
AAAAAACCGAAACCCACCGACCGTCATTCCCGCAAAAGTTGGAATCCAAAAACGCAACAGGAATTTATCGGAAAA
AAACAGAAACCCACCGCGTCATTCCCGCGCAGGCGGGAATCCAGAAACACAAACGCAACAGGAATTTATCGGAAAAAC
30 AGAAACCCACCGACCGTCATTCCCGCAAAAGCGGGAATCCAGCAACCGAAACACAGGAATCTATCAGCAAAACAG
AAACCCACCGACCGTCATTCCCGCGCAGGCGGGAATCCAGAAACACAAACGCGGCGAGGACTTTATCGGAAAAACAGAA
ACCCACCGACCGTCATTCCCGCAAAAGCTGGAATCCAAAAACGCAACAGGAATTTATCGGAAAAACAGAAAC
ACCCACCGCGTCATTCCCGCAAAAGCGGGAATCCAGACCCGTCGGCAGCGGAACTTACCGGATAAAACAGTTTCTTAGAT
TCCACGTCCAGATTCCCGCTTCGCGGGAATGACGAGATTTAAGTTGGGGGAATTTATCAGAAAAACCCCAACCCCA
35 AAAACCGGGCGGATGCCGCACCATCCGCCCAAAACCCGATTTAACCATTCAAACAAACCAAGAAAAACAAA

SEQ. ID NO:5

Nucleotide sequence of DNA region (772 bp) up-stream from the PilQ gene from *Neisseria meningitidis*

40 GCGATGTCGGGAAGCCTTCTCCGAATCATTACCCCTTGAGTCGCTGAAAATCGCCCAATCTCCGGAACGGCGGCAAT
CATGACGGCAAGAGCAGCATCCTGAACCTCAGTGCCATTGCCACCACCTACCAAGCAAAATCCGTAGAAGAGCTTGCCGC
AGAAGCGGCACAAAATGCCGAGCAAAATAACTTACGTTAGGGAAACCATGAAACACTATGCCTTACTCATCAGCTTTCT
GGCTCTCTCCGCGTGTTCCTCAAGGTTCTGAGGACCTAAACGAATGGATGGCACAACCGCAGCGGAAGCCAAAGCAGAAA
TCATACCTTTCCAAGCACCTACCTGCGGTTGCGCCGGTATACAGCCCGCCGAGCTTACAGGGCCGAACGCATTTCGAC
45 TTCCCGCGCATGGAACCGACAAAAAGGGGAAATGCCCCGACACCAAGCGTATTAAAGAAACGCTGGAAAAATTCAG
TTTGGAATAATATGCGTTATGTCGGCATTTTGAAGTCTGGACAGAAAGTCTCCGCTTCATCGAGGCTGAAGGTTATGTCT
ACACTGTCGGTGTGCGCAACTATTTGGGACAAACTACGGTAGAATCGAAAGCATTACCGACGACAGCATCGTCTGAAC
GAGCTGATAGAAGACAGCACGGGCAACTGGGTTTCCGTAAAGCAGAACTGCTGTTGAATTCTTCCGACAAAAACCCGA
ACAAGCGGCAGCACCTGCCGAGAACAAATTAAGAAGAGGATTACTCCATT

50

SEQ. ID NO:6

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hap gene from *Neisseria meningitidis*

GTGCGGCAAAAAACAGCAAAAGCCCGCTGTCGATTGCTGACCGTCCGCGTCCGTAAAATCAGCATAGGTTGCCACGCGC
 5 GGCTTGGGCGTTTTCCACACAAGCCTCTGCCATCGGACGACAGGTTTTTCCCGATATGCGTATCAGCCCACGCGCGC
 GCGCCCGGGTGCGGTAGCGACTGCCGAATCGTTGGAACGTTATCCGACATAAAACCCCGAAAATTCAAACAGCGCGC
 ATTATAGCAATGCCGTCTGAAGTCCGACGGTTTGGCTTTTCAGACGGCATAAAACCGCAAAATGCTTGATAAATCCGTC
 CGCTGACCTAATATAACCATATGAAAAACGAACACATACGCCCTTCTGCTCGGTATAGGCTCGCTGCTGGGTCTGTT
 CCATCCCGCAAAAACCGCCATCCGCCCAATCCCGCGACGATCTCAAAAACATCGGCGGCGGATTTCACCGCGCCATAG
 AGAAAGCGCGAAAATGACCGAAAACGCACAGGACAAGGCGCGGAGGCTGTCGAAACCGTCGTAATCCCGGAGCTTG
 10 TCGAGCAATCCTGTCCGACGAGTACGTGCAAAATATGATAGCCCGGCGTTTCCATTCCGGATCGTTGCCGCGCCGCTCG
 GACTTGGCGCAATACAACGACATTATCAGCAACGGGGCAGACCGCATTATGGCAATGGCGGAAAAAGAACAGCCGTCGG
 GCACGAAACCATACGGCAAGACCAAACTTCAACAGGCGCGGGCAACTGTACGGCTTCATCAGCGTCATCTGATACTGC
 TTTTGGCGTCTTCTCGTATGGAGCGGCTACCCCGCAACCGCGCTCCCTTGGCGGCGCACAGTGGTTGCTTGGCG
 GGTGCTTTCTGTGATTGGAAGAAGCCGAGACCAAGCAAAAATTAATTGCAATCCTAGGCGGTGCTTCATATCCGCCGGA
 15 ACGCCGAACCGCACATATAGGCACATCCCGCGCGCCGCGGAAGCGGAGCGCGCTCCCAACAAACCCGAATCCCG
 TCAGATAAGGAAAAATA

SEQ. ID NO:7

Nucleotide sequence of DNA region (924 bp) up-stream from the NspA gene from *Neisseria meningitidis* (serogroup B) (ATCC13090)

GGAAACGAACACGCGCTTCGGTCATACGCCGCCGAAAGGTTTGGCGCAAGACGAAGCGCCCTCGACATCGAAGACGCGG
 TACACGGCGCGCTGGAAGCGCGGGTTTTGTCCACTACGAAACATCGGCTTTTGGCAACAGCCATGCAGTCCCGCCAC
 AATTTGAACTACTGGCAGTTTCGGCGATTATTTAGGCATAGGCGCGGGCGCTCACGGCAAAATTTCTATCCCGACCGCAT
 25 CGAGCGCACCCTCGCGCGCGCCACCCCAACGACTACCTCGCTTAATGCAAGGCCAACCGAGTGAAGCCGTGGAACGCA
 AAACCGTTGCCGCGGAAGATTGCGGTTTGAAGTTCATGATGAACGCCCTGCGCCTGACCGACGCGTACCCGCCGCGATGT
 TGCAGGAGCGCACGGGCTACCGAGTGCCTAAATCATGGCGCAATCGAAACGGCAAGGCAAAAAGGCGCTGCTGGAACCC
 GACCCCGCGCTATTCGCGCCGACCGAAAAGGAGCGCTGTTTTTAAACGATTGCTGCAGTGTTTTTATAGTGGATTAA
 CAAAACCAAGTACGGCGTTGCTCGCCTTAGCTCAAGAGAAGGATTCTCTAAGGTGCTGAAGCACCAGTGAATCGGTT
 30 CCGTACTATTGTACTGTCTGCGGCTTCGTGCGCTTGTCTGATTGTTTGTAACTCACTATATAAGCGCAACAAATCGG
 CGGCGCGCGGGAACCCCGCCGAAACGCGTCCGGAATATGCTTATCGATGGAAACGCAGCCGCATCCCCCGCGG
 GCGTTTCAGACGGCACAGCCGCGCGGAAATGTCGACGCTTAAGGCACAGACGCACAAAAACCGTATGCTGCACCT
 GCAACAATCCGACAGATACCGCTGTTTTTCAAAACCGTTTGCA

SEQ. ID NO:8

Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpB gene from *Neisseria meningitidis* (serogroup B)

AAGTGGGAATCTAAAAATGAAAAGCAACAGGAATTTATCGGAAATGACCGAACTGAACGGACTGGATTCCCGCTTTTCGC
 GGGAAATGACGGCGACAGGGTTGCTGTTATAGTGGATGAACAAAAACAGTACGTCGTTGCTTCGCTTAGCTCAAAGAGA
 ACGATTCTCTAAGGTGCTGAAGCACCAGTGAATCGGTTCCGTCTATTGTACTGCTCTGCGGCTTCGTGCGCTTGTCTCT
 40 GATTTCTGTTTCGTTTTTCGGTTATTTCCGATAAATTACCGCGGTTTCTCGTCATTCTTTAACCCTTCGTATTCGCCGCG
 AGGCGGGAATCTAGTTTTTTGAGTTCAGTTGTTTCTGATAAATCTTGACAGCTTTGAGTTCCTAGATTCCCACTTCG
 TGGGAATGACGGTGGAAAGTTGCCGTTATTCGGATAAATTTCTGTAACGCATAATTTCCGTTTTACCCGATAAATGCC
 CGCAATCTCAAATCCGCTATTCCCAAAAACAAAAATCAAAAACAGAAATATCGTCATTCCCGCGCAGGCGGGAATCT
 45 AGACCTTAGAACAACAGCAATATTCAAAGATTATCTGAAGTCCGAGATTCTAGATTCCCACTTTCTGTTGGGAATGACGAA
 TTTTAGGTTTTCTGTTTTTGGTTTTCTGTCTTGGCGGAATGATGAAATTTTAAGTTTTAGGAATTTATCGGAAAAACAG
 AAACCGCTCCGCGCTCATTCCCGCACAGGCTTCGTCTATCCCGCGCAGGCTTCGTCTATCCCGCATTGTTAATCCACTA
 TATCCCGCGGTTTTTTACATTTCGACAAAACCTGTCAACAAAAACAACTTCGCAAAATAAAACGATAATCAGCTT
 50 TGCAAAATCCCCCCCCCTGTTAATATAAATAAAATAATTAATTAATTTTCTTATCTGCAAACTTTAACGGT
 TTGGATTACTTCCCTTCATACACTCAAGAGGACGATTGA

SEQ. ID NO:9

Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpA gene from *Neisseria meningitidis* (serogroup B)

CTATAAAGATGTAATAAAAAATCTCGGTAACGGTAACACTTTGGCTCAGCAAGGCAGCTACACCAAAACAGACGGGTACAA
 CCGCAAAAATGGGGGATTTACTTTTAGCAGCCGACAACTGACACAGCCGTTTCACGAACAAAATGCTATCCATTAGCCAT
 GTTCGGGAAACACGATTTCCCGTTTGTGTTTAGGCTGTCTAAACAAATAACCATAAATGTATATCATTATTTAAATAAA
 60 ATAAAAGTATTTAAGTATTATGACGAAATTTAGAGAAAGAGTAGACTGTGCGATTAAATGACAAACAAATAGTGAGAAAG
 GAAATATTTACTATCCGAGCAGAGCATATTTAGGTAGCCTGTAAGTGTTCCTGCTGGCGGAAGAGGATGAAGGTGGA
 CTTACCCGAGAAATATGCTGTTGTGTGATATGGATGCCATGCCGGAAGCAATTGATGCAATCACGGCAGTCCCTACT
 TGAATGAAACCTGTGCTGTCAGAAATTTGAAAACGCTATTTTAAAGAAAGGATAAAGGAGAAAGATTTTGGTTTTTAA
 GCTGCATGAAACCGTGTGGAATAAATGCACACCTACGATAAATTAATAATTTTCGTTTTTATTCTACAAGCTATTATA
 65 TATGATTGCTAAAGTTTTATTTTATAGTGCCAAAAATATATTTATATACCTTCATATTGTTTATATGCTTTATTTGA
 ATATATCTTACGATGGGGAATATTTATATATTTTATAATAAATTTTACTCATTGCTAATATGTCATGGAATATTACTT
 GTATTTTGTAGAATTTTCCATATGAAATATTCCATTACTATTTTCTGAACTTTATTAGTTTTTAAATATTTT

ACCTCTTATATTTACCATAAGAGAGCTAATTGATTATATTATATTGAGTCGATAATTAATTTATTCTTAATTTTAATTC
CTCACGTTATTTTTTAAATTTACTTGAAAGGAAAGCAGAT

SEQ. ID NO:10

- 5 Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpC gene
from *Neisseria meningitidis* (serogroup B)

GGAAACAGAGAAAAAGTTTCTCTTCTATCTTGGATAAATATATTTACCCCTCAGTTTAGTTAAGTATTGGAATTTATACC
TAAGTAGTAAAAAGTTAGTAAATATTTTTTAACTAAAGAGTTAGTATCTACCATAATATATTTCTTAACTAATTTCTAGGC
TTGAAATATGAGACCATATGCTACTACCATTATCAACTTTTTATTTTGTATTATGGGAGTGTCTTACTATGACCTCA
10 TGGAACCTGTGAATGAAAAGACAGATCAAAAAGCAGTAAGTGCGCAACAGGCTAAAGAACAAACAGTTTCAACAATCC
CGAGCCAATGACAGGATTTGAACATACGGTTACATTGATTTTCAGGGCACCAAAATGGTTATCCCTATGGCTATCTTG
CACGGTATACGCAAGACAATGCCACAAATGGCTTTCCGACACGCCCGGGCAGGATGCTTACTCCATTAATTTGATAGAG
ATTAGCGTCTATTACAAAAAACCCGACCAAGGCTGGGTTCTTGAGCCATACACCAGCAAAACAAAGCACACTTTATCCA
15 ATTTCTACGCGACGGTTTGGATAGCGTGGACGATATTGTTATCCGAAAAGATGCGTGTAGTTTAAGTACGACTATGGGAG
AAAGATTGCTTACTTACGGGGTTAAAAAATGCCATCTGCCTATCTGAAATACGAGGCTTATGAAGATAAAAGACATATT
CCTGAAAATCCATATTTTTCATGAATTTTACTATATTAAAAAGGAGAAAAATCCGGCGATTATTACTCATCGGAATAATCG
AATAAACCAACTGAAGAAGATAGTTATAGCACTAGCGTAGGTTCTGTATTAAACGGTTTCACGGTACAGTATTACCCGT
TTATTCGGGAAAGCAGCAGCTCACACAGCAGGAGTTGGTAGGTTATCACCACAAAGTAGAGCAATTGGTACAGAGTTT
20 GTAAACAATTCAAATAAAAAATAATTTAAAGGATCTTATT

SEQ. ID NO:11

- 25 Nucleotide sequence of DNA region (1000 bp) up-stream from the Omp85 gene
from *Neisseria meningitidis* (serogroup B)

ACGTCGGAACCGTGATTCCGCAACGCCGCGCCCAAAACCAAGCCCAAGCCAAAATGCCGATATAGTTGGCATTGGCAAT
CGCGTTAATCGGGTTGGCGACCAAGTTTCATCAGCAGCGATTTCACACTTCCACAATGCCGGAAGGCGCGCGCGGACA
30 CATCGCCCGCGCCGCCAAACAATGTGCGTGGGAAACCATACCGGCGATGACGGCGGTGAGGGTTCGGGAAACGTA
CCAAATGAGGTAAGGATGATAATCGGCCTGATATGCGCCTTGTTCCTTTTGGTGCTGCGCGATTGTGGCCGCCACCAA
AATAAATACCAAAACCGCGCGACCGCTTTGAGCGCGCGGACAAACAGGCTGCCGAACAAGCCTGCCGCCAAGCCAGTT
GCGGGGAAACCGAACCATTACGATGCCAACGCCAAACCGCGGCAATCTGCCTGACCAAGGCTGACGCGGCCGATCGCA
35 TGAATTAAGGATTTGCCGAACGCCATAATCTTCTCTATGTTGTGATATGTTAAAAATGTTGATTTTAAAGAAAACT
CATTCTCTGTGTTTTTTTTTATTTTTCGGCTGTGTTTAAAGGTTGCGTTGATTGCCCCATGCGAGTGCCGGACAGGCTTTG
CTTTATCATTCCGGCGCAACGGTTTAAATTTATGAAACGAAAAATAATTTATTTAATCCTGCCTATTTTCCGGCACTATTC
GAAACGCGAGCCTGTTTTCCATATGCGGATTGGAACAAAAATACCTTAAACAAGCAGATACATTTCCGGCGGGCGCAAC
40 CTCGAAATACCGGCGCGCAGTATGCCGTCTGAAGTGTCCCGCCCGTCCGAACAACACAAAAACAGCCGTTTCAAAACCCCT
GTCCGAACAGTGTTAGAATCGAAATCTGCCACACCGATGCACGACACCCGTACCATGATGATCAAACCGACCCGCTGCT
CCTGCCGGCTTATTTTTCTTTCCGCACGCATACGCGCCT

SEQ. ID NO:12

- 40 Nucleotide sequence of DNA region (772 bp) up-stream from the PilQ gene from
Neisseria meningitidis (serogroup B) (ATCC13090)

GCGATGTCGGGAAGCCTTCTCCGAATCATTACCCCTTGAGTCGCTGAAAATCGCCCAATCTCCGGAACCGCGGCAAT
CATGACGGCAAGAGCAGCATCCTGAACCTCAGTGCCATTGCCACCACCTACCAAGCAAAATCCGTAGAAGAGCTTGC CGC
AGAAGCGGCACAAAATGCCGAGCAAAAATACTACGTTAGGGAAACCATGAAACACTATGCCTTACTCATCAGCTTTCT
45 GGCTCTCTCCCGGTGTTTCCCAAGGTTCTGAGGACCTAAACGAATGGATGGCACAAACGCGACGCCAAGCCAAAGCAGAAA
TCATACCTTTCCCAAGCACTACCCCTGCCGCTTGGCGCGGTATACAGCCCGCCGAGCTTACAGGGCCGAACGCATTCCGAC
TTCCGCCGATGGAACCGACAAAAAGGGGAAAAATGCCCCGACACCAAGCGTATTAAAGAAACGCTGGAAAAATTCAG
TTTGAAAAATATGCGTTATGTCGGCATTGTTGAAGTCTGGACAGAAAGTCTCCGGCTTATCGAGGCTGAAGGTTATGTCT
ACACTGTGCGGTGTCGGCAACTATTTGGGACAAAACACGGTAGAATCGAAAGCATTACCGACGACAGCATCGTCTGAAAC
GAGCTGATAGAAGACAGCACGGGCAACTGGGTTTCCCGTAAAGCAGAACTGCTGTTGAATTTCTCCGACAAAAACACCGA
50 ACAAGCGGCAGCACCTGCCGACAGAAAAATTAAGAAGAGGATTACTCCATT

SEQ. ID NO:13

- 55 Nucleotide sequence of DNA region (1000 bp) up-stream from the Hsf-like gene
from *Neisseria meningitidis* (serogroup B)

TTTGTTTTTCTTTGGTTTGTGTAATGGTTAAATCGGGGTTTGGGGCGGATGGTGCGGCATCCGCCCGGTTTTTGGG
GGTTGGGGGTTTTCTGATAAATCCCCCAACTTAAATCTCGTCATTCCCGCGAAGGCGGGAACTCTGGGACGTGGAATCT
AAGGAAACTGTTTTATCCGGTAAGTTCCGTGCCGACGGGTCTGGATTCCCGCTTTTGGCGGAATGACGGCGGTGGGGTT
TCTGTTTTTTCCGATAAATCCTGTTGCGTTGCGTTTTTGGATTCCAGCTTTTGGCGGAATGACGGTCCGTTGGGGTTCT
60 GTTTTTTCCGATAAAGTCTGCCGCGTTGTGTTCTGGATTCCCGCCTGCCGCGGAATGACGGTCCGTTGGGGTTCTGT
TTTTGCTGATAGATTCTGTGGTTTTTCGGTTGCTGGATTCCCGCTTTTGGCGGAATGACGGTCCGTTGGGGTTCTGT
TTCCGATAAATCCTGTTGCGTTGCGTTTTTGGATTCCCACTTTTGGCGGAATGACGGTCCGTTGGGGTTCTGT
CGATAAATCCTGTTGCGTTGCGTTTTTGGATTCCCACTTTTGGCGGAATGACGGTCCGTTGGGGTTCTGT
TAAAGTCTGCCGCGTTGTGTTCTGGATTCCCGCCTGCCGCGGAATGACGGTGGGGTTTCTGT
70 TTCCGTGGTTTTTCTATGATTCAATCATTCTGATAAATCCATAATCTAAATCTCGTCATTCCCGCGAAAGCGGG
AATCTAGGACGTGGAATCTAAGGAACTGTTTTATCCGGTAAGTTTCCGTGCCGACGGGTCTGGATTCCCGCTTTTGGCGG
GAATGACGGCGGTGGGGTTTCTGTTTTTCTGATAAAGTCTGCCGCGTTGTGTTCTAGATTCCCGCTTTTGGCGGAAT
GACGGCGGTGAGGTTCTGTTTTTCCGATAAATCCTGT

SEQ. ID NO:14

- 70 Nucleotide sequence of DNA region (1000 bp) up-stream from the Hap gene
from *Neisseria meningitidis* (serogroup B)

AATCAGCATAGGTTGCCACGCGCGCTTGGCGTTTCCACACAAAGCCTCTGCCATCGGCAGCAGGTTTTCCCGAT
 ATGCGTATCAGCCCCACGCCCGCGCGCGGTGCGGTAGCGACTGCCGCAATCGTTGGAACGTTATCCGACATAAAACC
 CCGAAAAATCAAAACAGCCGCGATTATAGCAATGCCGTCTGAAGTCCGACGGTTTGGCTTTCAGACGGCATAAACCG
 CAAAAATGCTTGATAAATCCGTCCGCTGACCTAATATAACCATATGGAAAAACGAAACACATACGCCCTTCTGCTCGGT
 5 ATAGGCTCGTGTGGGTCTGTTCCATCCCGCAAAACCGCCATCCGCCCCAATCCGCGGACGATCTCAAAACATCGG
 CGGCGATTTTCAACGCGCCATAGAGAAAGCGCGAAATGACCGAAACGCACAGGACAAGCGCGGCGAGGCTGTGAAAC
 CGTCGTCAAATCCCCGGAGCTTGTGAGCAAACTCTGTCCGACGAGTACGTGCAAAATATGATAGCCCGCGCTTCCATT
 CGGGATCGTGTCCGCGCGCTCCGACTTGGCGCAATACACGACATTATCAGCAACGGGGCAGACCGCATTATGGCAATG
 GCGGAAAAAGAACAGCCGTCCGGCAGCAAAACCATACGGCAAGACCAACCTTCAACAGGCGCGGGCAACTGTACGGCTT
 10 CATCAGCGTCATCCTGATACTGCTTTTTGCGCTCTTCCCTCGTATGGAGCGGCTACCCCGCAACCGCGCTCCCTTGGCG
 GCGGCACAGTGGTTGCTTGGCGGTGCTTTCGTGATTGGAAGAAGCCGAGACCAAGGCAAAATTAATTGCAAACTCTA
 GGGCGTGTCTCATATCCGCCGAACGCCGACATATAGGCACATCCCGCGCGCGCGGGAAGCGGAAGCCGCGCC
 CTCCTCAACAAACCGGAATCCCGTCAGATAAGGAAAAATA

15 SEQ. ID NO:15
 Nucleotide sequence of DNA region (1000 bp) up-stream from the LbpA gene
 from *Neisseria meningitidis* (serogroup B)

GATTTTGGTTCATCCCGACAGCTTCTTGTGCAAGGGCGTGAAATTCCTTTGGTTAGCCAAGAGAAAACCATCAAGCTTGC
 CGATGGCAGGGAAATGACCGTCCGTGCTTGTGCGACTTTTGGACCTATGTGAAACTCGGACGGATAAAACCGAACGCC
 20 CGGCAAGTAAACCAAGCGGAAGATAAAAGGGAGGATGAAGAGAGTGCAGGCGTTGGTAACGTCGAAGAAGCGAAGGC
 GAAGTTTCCGAAGATGAAGCGGAAGACCGGAAGAAATCGTCGAAGAAGAACCCGAAGAAGAGCTGAAGAGGAAGAAGC
 TGAACCCAAAGAAGTTGAAGAAACCGAAGAAAAATCGCCGACAGAAGAAAGCGGCGAGCGTTCAAACGCCATCTGCCTG
 CCTCGGAAGCCTCTAAAGGCAGGACATCGACCTTTTCTGAAAGGTATCCGCACGGCGGAAGCGGACATTCCAAGAACCC
 GGAAAGCAGCACTATACCGGCACTTGGGAAGCGGTATCGGCACACCCATTCAATGGGACAATCAGGCGGATAAAGAAC
 25 GGCAAAAGCAGCAATTACCGTTAATTTCCGGCAGAAATCGATTTCGGGAACGCTGACGGAGAAAAACGGGTGTACAACCTG
 CTTTCTATATTGAAAACGGCAAGATTGAGGGCAACGGTTTCCACGCAACAGCAGCAGCTCGTGAGAACGGCATCAATCTT
 TCGGGAAATGGTTCGACCAACCCAGAACCTTCCAAGCTAGTGATCTTCTGTAGAGGAGGATTTTACGGCCCGCAGCG
 GAGGAATTTGGCGGTTATTTTCAATAAGGATGGGAATCTCTTGGTATACTGAAGGTACTGAAAAATAAGTTGAAGT
 30 TGAAGCTGAAGTTGAAGTTGAAGCTGAACTGGTGTGTCGACAGTTAGAACCTGATGAAGTTAAACCCCAATTTCGGCG
 TGGTATTCGGTGCAGGAAGAGATAATAAGAGGTGAAAAA

SEQ. ID NO:16
 Nucleotide sequence of DNA region (1000 bp) up-stream from the LbpB gene
 from *Neisseria meningitidis* (serogroup A)

35 CGGCGTTAGAGTTTAGGGCAGTAAGGGCGCGTCCGCCCTTAGATCTGTAAGTTACGATTCCGTTAAATAACTTTTACTGA
 CTTTGAGTTTTTTGACCTAAGGGTGAAAGCACCTTACTGCTTAAAGTCCAACGCAAAACCAAAAGACAAAAACACTT
 TTATTACCCTAAAATCGAACACCCATAAATGACCTTTTGTCTTTGGCGAGGCGCGAGTAAGGGCGCGTCCGCCCTTAG
 ATCTGTAAGTTATGATTCCGTTAAATAGCCTTTACTGACTTTGAGTTTTTTGACCTAAGGGCGGACGCGCCTTACTGCT
 40 TCACCTTCAATGGGCTTTGAAATTTGTTGCGCTTGGCTTGCTTGACCTAAGGGTGAAAGCACCTTACTGCGCCCTCGCC
 AAAGACGAAAAGGGTTATTTACGGGGGTGGATTTTAGGCGAGTAAGGGCGCGTCCGCCCTTAGATCTGTAAGTTATGATT
 CCGTTAAATAGCCTTTACTGACTTTGAGTTTTTTGACCTAAGGGTGAAAGCACCTTACTGCTTCACTTCAATGGGCTT
 TGAATTTTGTGCTTTGGCTTGCTTGAATCTAAGGGTGAAAGCACCTTACTGCGCTCTCGCCGAAGACAACGAGGGCTA
 TTTACGGCGTTAGAGTTTAGGGCAGTAAGGGCGCGTCCGCCCTTAGATCCAGACAGTCACGCTTTGAATAGTCCATTTT
 45 GCCAAAGAATCTAAACGCAGGACCTAAGGGTGAAAGCACCTTACTGCTTACATCCAAGCACCTTACTGCACCAGC
 TCCACGCACCTTACTGCCCTACGTCCACGCACCTTACTGCCCTACATCCAAGCACCTTACTGCTTACATAGACATG
 ACAGACCGCGAGCAGCGGAACAGGACTAAAAACAATTAAGTGATATTTTCCCACTATAATAGACATGTATAATTATA
 TTACTATTAATAATAATAGTTTATCCTCCTTTTCACTCC

SEQ. ID NO:17
 50 Nucleotide sequence of DNA region (731 bp) up-stream from the TbpA gene
 from *Neisseria meningitidis* (serogroup B) (ATCC13090)

TATGAAGTCGAAGTCTGCTGTTCCACCTTCAATTATCTGAATTACGGAATGTTGACGCGC
 AAAACAGCAAGTCCGCGATGCAGGCAGGAGAAAGCAGTAGTCAAGCTGATGCTAAAACG
 GAACAAGTTGGCAAGATATGTTCTCCAAAGGCGAGCGCACCGATGAAAAGAGATTCCA
 55 AACGACCAAAACCTCGTTTATCGGGGTCTTGGTACGGGCATATTGCCAACGGCACAAGC
 TGGAGCGGCAATGCTTCCGATAAAGAGGGCGGCAACAGGGCGGACTTACTGTGAATTTT
 GGTACGAAAAAATTAACGGCACGTTAACCGCTGACAAACAGGCAGGCGGCAACCTTTACC
 ATTGTGGCGGATATTGAGGGCAACGGTTTTTCCGGTACGGCGAAAACTGCTGACTCAGGT
 TTTGATCTCGATCAAAGCAATAACCCGCACGCTAAGGCATATATCACAACGCCAAG
 60 GTGCAGGGCGGTTTTTACGGGCCCAAGCCGAAGAGTTGGGCGGATGGTTTGCCTATTCCG
 GACGATAAAACAAACGAAAAATGCAACAGATGATCCGGCAATGGAATTCAGCAAGCAGT
 GCAACTGTGCTATTCGGTGCGAACGCCAAAAGCCTGTGCAATAAGCACGGTTGCCGAAC
 AATCAAGAATAAGGCCTCAGACGGCACCGCTCCTCCGATACCGCTGAAAGCGAAGAGT
 65 AGGGAACACAT

SEQ. ID NO:18
 Nucleotide sequence of DNA region (373 bp) up-stream from the OmpA gene
 from *Neisseria meningitidis* (serogroup B) (ATCC13090)

70 CGTACCGCATTCGCACTGCAGTGAAGAAAGTATTGAAGCAGTGAAGCAGGCGATAAAGTGCCTGCACAAAGCGGTTTA
 CCAAGAGTCCGTCAAAGTCATCGACCGCATCGCCGACAAAGGCGGTTCATATAAAACAAAGCGGCTCGCCACAAAACCC
 GTTTGTCTCAAAAGTAAACCTTGGCTTGATTTTGAACCACTGCAATCCGGTTTTTCATCGTCGATTCCGAAAACCCC
 TGAAGCCCGACGGTTTTTGGGGTTTTCTGTATTGCGGGGACAAAATCCCGAAATGGCGGAAAGGTTGCGGTTTTTATCCG
 AATCCGCTATAAAATGCCGCTGAAAAACAATATGCCGACAATGGGGTGGAG

SEQ. ID NO:19

Nucleotide sequence of DNA region (1000 bp) up-stream from the Pla1 gene from *Neisseria meningitidis* (serogroup B)

5 TTTTGGCTTCCAGCGTTTCATTGTTTTCTGACAAAGTCGTAAGTCAGCTTCAGATTGTTGG
CTTTTAAAGTCTTCGACCGTACTCTCATCAACATAGTTTCGACCGATTGTAGATGTTCA
GAGTATCGGTGGCAGCGGCTTCGGCATTGGCAGCAGACGCGCTCTGCTTGAGGTTGCA
CGGCGTTTTTTTCGCTGCCGCGCAGGCTGCCAGAGACAGCGCGCCAAAACGGCTAATA
CGGATTTTTTCATACGGGCGAGATTCTCTGATGAAAGAGGTTGGAAAAAAGAAATCCCGCG
10 GCCCCATCGTTACCCCGCGCAAGGTTTGGGCATTGTAAAGTAAATTTGTGCAAACTCAA
AGCGATATTGGACTGATTTTCTTAAAAAATTATCTGTTTCCAAAAGGGGAGAAAAACGT
CCGCCGATTTTGGCGTTTTTTTGGCGCTGTGAGGGTGTCCGACGGCGGATAGAGAGAA
AGGCTTGCATATAATGTAACCCCTTTAAAAATTGCGCGTTTACAGAATTTATTTTCTT
CCAGGAGATTCCAATATGGCAACAGCGCACAAAGCAGCAACGTGCCCGCCAGTCCGTC
15 AAACACGCGCCCAATGCTAGCCTGCGTACCGCATTCGCGACCGCAGTGAAAAAGTA
TTGAAAGCAGTCGAAGCAGGCGATAAAGCTGCCGACAAGCGGTTTACCAAGAGTCCGTC
AAAGTCATCGACCGCATCGCCGACAAAGGCGTGTCCACAAAAACAAAGCGGCACGCCAC
AAAAGCCGCTCTGTCTGCAAAAGTAAAAGCCTTGGCTTGATTTTGTCAAAACCGCCAAGGC
GGTTGATACGCGATAAGCGGAAACCCCTGAAGCCCGACGGTTTGGGGTTTTCTGTATTG
20 CGGGGGCAAAATCCGAAATGGCGAAAGGTCGATTTTTATCCGAATCCGCTATAAA
ATGCCGTTTGAAAACCAATATGCCGACAATGGGGGCGGAG

SEQ. ID NO:20

Nucleotide sequence of DNA region (1000 bp) up-stream from the FhaB gene from *Neisseria meningitidis* (serogroup B)

25 TACGGAAATGCAAGCGGATCCAGAAGTTACAGCGTGCATTATTCGGTGCCCGTAAAAAATGGCTGTTTTCTTTAATC
ACAAATGGACATCGTTACACGAAAGCAACCGAAGGCTATTCCGTCATACGATTACACCGCAACAAATATCAGAGCAGC
CTGGCCCGCAGCGCATGCTTTGGCGTAACAGACTTCATAAACTTCAGTCGGAATGAAATTTATGGACACGCCAAACCTA
TAAATACATCGACGATGCCGAATCGAAGTGAACGCGCCGCTCTGCGAGCTGGGAAGCCGAATTCGCCACCGTGT
30 ACCTCAACCGTTGGCAGTTGACGGCAAGTTGTCTTACAAACGCGGACCGCATGCGCCAAAGTATGCCTGCACCGGAA
GAAAACGCGCGCATATTCTTCCAGGTACATCTCGTATGAAATCATTACTGCCGGTTTGGACGCGAGCCGCCATTAT
TTTAGGCAACAGCAGTTTTCTACGCAACCGCCATTCAAGCTCAATGGAACAAACCGCGTTGGTTGCCCAAGATAAAT
TGTCATTCGCGAGCGCTACACCGTTCCGCGATTGTATGGGAGCAGAGTCTTTTCGGAGAGCGAGGTTCTACTGGCAG
35 AATACCTTAACTTGGTATTTTCATCCGAACCATCAGTTCTATCTCGGTGCGGACTATGGCCGCGTATTGGCGAAAGTGC
ACAATATGTATCGGGCAAGCAGCTGATGGGTGCAAGTGGTCCGCTTCAGAGGAGGGCATAAAGTAGGCGGTATGTTGCTT
ATGATCTGTTTGGCGGCAAGCCGCTTATAAACCCAAAGGCTTCAGACGACCAACACCGTTACGGCTTCAACTGAAT
TACAGTTTCTAACCTCTGAATTTTTTACTGATATTTAGACGGTCTTCTTATCTCAGACCGTCAAACTTACCTACGT
ACTTGCGCGCAGTACGTTTCATCTTCAAAATGGAATAGAC

40 SEQ. ID NO:21

Nucleotide sequence of DNA region (1000 bp) up-stream from the Lipo02 gene from *Neisseria meningitidis* (serogroup B)

TTATCTTGGTGCAAACTTTGTGGGGTCCGACTGGCTACGGCTTTGGGTTTGGACCCGCTCATCGGTCTGATTACCGGT
TCGGTGTCTGCTGACGGGCGGACACGGTACGTAGGTGCGTGGGACCTAATTTTGAACGCAATACGGCTTGGTCCGGCGC
45 AACCGTTTGGGTATTGCATCGGCTACTTTCCGGCTGGTGTTCGGCGGCTGATCGGCGGGCCGGTTGCGCGCCGCTGA
TCAACAAAATGGGCGCAAAACCGTTGAAAACAAAAACAGGATCAGGACGACAACCGGACGACGTGTTCCGAGCAGGCA
AAACGCAACCCGCTGATTACGGCGGAATCTGCCGTGAAACGCTTGGCATGTTTGGCCGCTGTTTGGCGTTTGGCGAGAT
TTGAGACGGCTTCCGCAAGAAATATCTGTTGCACTGCCCAAATTCGTGTGGTGTCTGTTTGGCGGCGTGGTCATCCGCA
50 ACATCTCTACTGCCGATTCAGGTCAATATGTTGCAACCGGCCATCGATGTGTTCCGCAATGCTTCGCTTTTCCGCTTTT
TTGGCAATGGCGTTGCTGAATTTGAACTGTGGGAGCTGACCGGTTTGGCGGGGCGTGAACCGTGATTCTTGGCGTACA
AACCGTGGTGATGTTTTGTACGCGACTTTTGTACCTATGTCTTATGGGGCGGACTATGATCGGCGAGTATTGGCTG
CCGGCCATTGCGGTTTCCGCTTGGGTGCAACGCCGACGGCGGTGGCAAAATATGAGTCCGTACGCATATTTCCGGCGG
TGCATAAGGCGTTTTTGTATTGTCCTATGTTGCGCGCTTCTCGTCGATTTGATTAATGCCGCGATTCTCACCGGTTT
55 TGTGAATTTCTTTAAAGGCTGATTTCCGCCCTTCCGACAAAGCACCTGCAAGGTTTACCGCTGCAGGTGCTTTTGCTA
TGATAGCCGCTATCGGTCTGCACCGTTTGAAGGAACATC

SEQ. ID NO:22

Nucleotide sequence of DNA region (1000 bp) up-stream from the Tbp2 gene from *Neisseria meningitidis* (serogroup B)

60 CCTACTCCACCGATTCCAAATATGCTCGGCGGACCCAGCAAGCCTGGAATTTTTGAACTCGGGCATCAAAATC
GTCAAAACCATATGGGCGTTGCCTTTTGGGACGAAAACGTTGAAGTCAGCCCCGAAGAGTCAGCGTGCCTTTGAAGA
AGGCGTGCCGGTTGCATGAACGGCAAGAAATACGCCGACCCCGTCGAATCTTCTCGAAGCCAAACCGCATCGGCGGGC
GCCACGGCTTGGGTATGAGCGACCAAAATCGAAAACCGCATCATCGAAGCCAAATCGCGCGGCATCTACGAAGCCCGGGT
65 ATGGCGTTGTTCCACATCGCCTACGAACGCTTGGTGACCGGCATCCACAACGAAGACACCATCGAACAATACCGCATCAA
CGCCCTGCGCCTCGGCCGTTTGTCTACCAAGGCCGCTGGTTCGACAGCCAAGCCTTGATGTTGCGGCAACCGCCCAAC
GCTGGTTCGCGCAAGCCGTTACCGCGAAGTTACCTCGAACTGCGGCGCGGCAACGACTACTCGATTCTGAACACCGAA
TCGCCCAACCTGACCTACCAACCCGAACGCTGAGTATGGAAGAGTCGAAGTGGCGGCTTACCCCGCTCGACCGCAT
CGGACAGCTCACGATGCGCAACCTCGACATCACCGACCCGCGCCAACTGGGCATCTACTCGCAAGCGGTTTGGCTGT
70 CGCTGGGCGCAAGGTCGTTTACCGCAGTTGGGCAATAAGAAATAAGGTTTGTCTTTTGCATCATTAGCAACTTAAGG
GGTCGTCTGAAAAGATGATCCCTTATGTTAAAGGAATCCTATGAAAGATACAAAGTCGTATTTATCAGGAAGCCAG
TTGTCCAGCCTGTTTTTCCGGCGGGCAAGGTTCAACCCGTCATTTTACGCGGTTCTCAACAAACAAACCCCGCAAG

GCTGGCGGGTCGAGACCTTTGCAATAACATAGGTTACTAA

5 SEQ. ID NO:23

Nucleotide sequence of DNA region (1000 bp) up-stream from the PorA gene from *Neisseria meningitidis* (serogroup B)

GAATGACAATTGATAAGTTTCCCGAAATTCACACATAACCGAAACCTGACAATAACCGTAGCAACTGAACCGTCATTCCCG
 10 GCAAAAGCGGGAATCCAGTCCGTTTCAGTTTCGGTCATTTCCGATAAATGCCTGTTGCTTTTCATTTCTAGATTCCCACTT
 TCGTGGGAATGACGGCGGAAGGGTTTTGGTTTTTCCGATAAATCTTGAGGCATTGAAATTCCAAATTCGCGCTGCGC
 GGAATGACGGCTGCAGATGCCGACGGTCTTTATAGTGGATTAACAAAAATCAGGACAAGGCGACGAGCTGCAGACAGT
 ACAGATAGTACGGAACCGATTCACTTAGTGCTTCAGTATCTTAGAGAATCGTTCTCTTTGAGCTAAGGCGAGGCAACGTC
 GTACTGGTTTTTGTTCATCCACTATATATGACACGGAAACGCGCGCTCCAAACCATGCCGTCTGAAGAAAACTACACA
 15 GATACGCGCGTTATATTACAAATCGCGCCCGCTGGTTCGAAAAACCTCCACACTAAAAAACTAAGGAAACCCCTATGTCC
 CGCAACACGAAGAGCTGCAAGGTATCTCGCTTTTGGGTAAATCAAAAAACCAATATCCGCGCAATACGCGCCCGAAAT
 TTTGGAAGCGTTCGACAACAAACATCCCGACAACGACTATTTTCGTCAAATTCGCTGCCCCAGAGTTCCACGAGCTCTGCG
 CCATGACCGGGCAGCCGACTTCGCGACCATCGTCATCCGCTACATTCCGCACATCAAAATGGTGGAAAGCAAAATCCCTG
 AACTCTACCTCTTACGCTTCGCGAACACGCGGATTTTCATGAAGACTGCGTCAACATCATCATGAAAGACCTCATTGC
 20 CCGTATGGATCCGAAATACATCGAAGTATTCGGCGAGTTCACACCGCGCGCGGCATCGCCATTTCATCTTTCCGCAATT
 ACGGCAAGCAGGCACCGAGTTTGAAGCATTGGCGCGTAA

SEQ. ID NO:24

***Neisseria meningitidis* (serogroup B) PorA Promoter Region**

25 GATATCGAGGTTCGCGCTTGAATTGTGTTGTAGAAACACAACGTTTTTGAAAAATAAGCTATTGTTTTATATCAAAATA
 TAATCATTTTTAAATAAAGTTGCGGCATTTATCAGATATTTGTTCTGAAAAATGGTTTTTTCGGGGGGGGGTATA
 ATTGAAGACGTATCGGGTGTGTTGCCGATGTTTTTAGGTTTTATCAAAATTACAAAAGGAAGCCCAT

SEQ. ID NO:25

Nucleotide sequence of DNA region (1000 bp) up-stream from the PorB gene from *Neisseria meningitidis* (serogroup A)

30 gttttctgtttttgaggggaatgacgggatgtaggttcgtaagaatgacgggatataggtttccgtgcggtgaggttcgtc
 attcccgcgaggcggaatctagaacgtggaatctaaagaaacggttttatccgataagtttccgtgcggaacaagtttgg
 attcccgccctgcgcggaatgacgggatgtaggtttctaattttgggttttctgtttttgaggggaatgacgggatgtagg
 ttcgtaggaatgacgggatgtaggtttccgtgcggtgtaggttcgtcattcccgcgaggcggaatctagacctagaac
 35 aacagcaatattcaagattatctgaaagtcgagattctagattcccgccctgagcggaatgacgaaaagtggcgga
 tgacggttagcgttgcctcgccttagctcaaaagagaacgattctcctaaagtgctgaagcacaagtgaaatcggttccgta
 ctatttgcactgtctgcggttcgtcgccttgctcgtatttttggtaatacactatctcctgcccagggcggttttg
 catccgcccgttccgaaagaaacccggtgtgctgtttttgcccgtctttataacccccggtttgcaatgcccctcaatacc
 ctcccgagtaagtggtgtaaaatgcaaatcttaaaaaatttaataaacatattgttataaaacaaaaatacccataat
 40 attcttatccgtccttcaaaatgcacatcgaaattccacacaaaaacaggcagaagtttggtttttccagacaggaacatct
 atagtttccagacatgtaatcgccgagccctcggcggttaaatgcaaaagctaagcggttggaagccccggcgtgcttaaa
 tttcttaacaaaaaggaatacagcaatgaaaaatccctgattgcccgtgactttggcagcccttccctgttcagcaat
 ggctgacgttaccctgtacggcaccatcaaaacccggcgta

45 SEQ. ID NO:26

***Neisseria meningitidis* (serogroup B) PorB Promoter Region**

GTTTTCTGTTTTGAGGGAATGACGGGATGATAGGTTTCGTAAGAATGACGGGATATAGGTTTCGTCGCGGATGGATTCGTC
 ATTCGCGCGAGCGGGAATCTAGAACGTGGAATCTAAGAAACCGTTTTATCCGATAAGTTTTCCGTGCGGGAAGTTTG
 50 GATTCCCGCCTGCGCGGGAATGACGGGATTTAGGTTTCTAATTTTGGTTTTCTGTTTTGAGGGAATGACGGGATGTAG
 GTTCGTAGGAATGACGGGATATAGGTTTCGTCGCGGATGGATTCGTCATTCCCGCGCAGGCGGGAATCCAGACCTTAGAA
 CAACAGCAATATTCAAAGATTATCTGAAAGTCCGAGATTCTAGATTCCCGCCTGAGCGGGAATGACGAAAAGTGGCGGGA
 ATGACGGTTAGCGTTGCCTCGCCTTAGCTCAAAGAGAACGATTCTTAAGGTGCTGAAGCACTAAGTGAATCGGTTCCGT
 ACTATTTGTACTGTCTGCGGCTTCGTCGCTTGTCTGATTGTTTGTAAATCCACTAT

55 SEQ. ID NO:27

Nucleotide sequence of DNA region (1000 bp) up-stream from the siaABC gene from *Neisseria meningitidis* (serogroup B)

ATACGGCCAATGGCTTCAGAAAGCGATAAGCCTCTGGCTGAAAAACCGATTCTTGTGTCTCCCCACCGCACCCATAGA
 60 CGTAAAGGTATAGGATTTGGTAATCATGGTAACACATCACCGCGACGCGAGCAAAATATTTTGTGCGGATTTGCAACTA
 AATCTTCAAGGCAACAGTTCGTACTACATTGCCAGGTGTAGCTGCACATTCTGATCTCGTACATTGCGGTTGAACCA
 CCTACCGCAGCCACCGCATCCAACACACGCTCACCGGCTGCGCTGACGGGCATACGCACACTATTCCGAGCACGAATCAC
 CGACACATTGCGCGCATTTCTGCACCAACGACCATCACTTGTGGCTGATTGGCCATTTTTTTCAGGCGGCGCTTAA
 TAATTTCTGAACCTGACAGGCGTTTTACCGACCGCAAAATATCGCCAAACAGGCGACAGAAACCGTACCACTGCC
 65 GTGACCAACTGCTCTGGCAACTTAGTTTGTATGCGCACTACCGAGGCCATCGAAGAAAGGCCACCACCAAACTACTGC
 CGCGGCGCTTCCAAATCATAATATCAATACATCACTAATATTAGCGTACCAGCCGAAGCATAACTCGCCAACT
 GAGTGAATGACTGATTTATCTGAGCCTTATATAAATGAGCAACCGATGATTACATCAATCAGCTCCACTTCAGGA
 ATTTGAACCTCAGATTGTTGCCCTAAAGAGACAATTTTTTTCGCGTGGGCGCTGATGAAGGAATCGCAGAGCATCTAC
 AATTAACCTCCACACAATAATACTGCGTGACGAATATAAAATTCACCTTTAAACACAAGCCAAATCCTAATATAAT
 70 TATAAATGGCCTAATATAGCACTTAATCGAAATAAATTTATGAGTACGTAGATATAATTAGTATTCTTCTTTCCAAC
 TCCTTACTTATATATATATATCTTATAGATTCTAAATC

SEQ. ID NO:28

GCCAAAGCATTGGGCGCGGATGCCGCCGTGCGGAACGCGCGCGCGTCTTGCCAAAGCCGACTTGGTAACCGAAATGGT
CGGCGAGTTCCCGCAAGTCCGAAGGACAGTAGGGCAAACTACTATGCTGTTTGGACGGCGCAACCGAAGAAATTGCCGAAG
CCGTGCAGCAGCACTATCAGCGCGGTTTTCGCGGCGCAAGCTGCCGCAAGCAAAATTGCCCGCCGCTGGCACTGGCC
GACAAACTAGAAACCTTGGTCGGCATTTGGGGCATCGTCTGATTCCGACCGGGCAAGAACGCCCTACGCCCTGCGCG
CGCTGCTTTGGGTATTTTGGCGTATGCTATGCAGTAGTATGGTTGGACGTGAACGAATGATTACAGCGGCATTGCACAGCT
TCCCAAAAGGTTTGTCTCAACGAAAAACGCGCGTCTGAACACCGCGACTTTATGCAGCGCGCTTGGCGGTGTGTGCAA
AACGATTATCCGCAAGACATCGTTGCCGCCGTACTCGCCAAACAGCGCGCGGTTTGGACGATTGACCGCCAACTGCA
GGCCGTTGCCCGGTTTCAACAACTGCCCGAAGCGCGCGCGTTCGCCCGCGCCAAACAAACCGCGTGCAAAACCTGTGAAA
AAGCGATGCGGAGTTGGGCGCGGTTTCAAGAAAGCTGTGTGAACAGGACGAAGAAAAGCCCTCTTTGCCCGCGCGCAA
GGCTTGCAGCCGAAATTCGCCCGCGCGCTGCGCGGAAGGCAATTTCCAAACCGCCTTGTCCGAATGGCTTCGTCAAC
GCAAGTGTGCATGCTTTTGAAGCGGTGATGGTAAATGGCGGAAGATGCCGCCGTAAACAAACCGCCTGAACCTGCTGA
ACCGCTTGGCAGAGCAATGAACGCGGTATCGGCACATCGCGCTTTTGGGCGAGTAACCGTTGTACAGTCCAAATGCGCTC
TGAAGCCTTCAGACGGCATCGTGCCATATCGGCGAAGATAAA

Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpB gene from *Neisseria meningitidis* (strain MC58)

Nucleotide sequence of DNA region (1000 bp) up-stream from the *opc* gene from *Neisseria meningitidis* (serogroup A)

Nucleotide sequence of DNA region (1000 bp) up-stream from the siaD gene from *Neisseria meningitidis* (serogroup B)

Nucleotide sequence of DNA region (1000 bp) up-stream from the ctrA gene from *Neisseria meningitidis* (serogroup B)

84

5 TGCAATAAAATTTATCAGATTGATATTTTAATAATCTTGCAGAATACTTTCAGTGCCAGTGTCATTATTAGGGTAGATGC
 TAATGATATTTTGGCCACTTAATTCTAATGCTTTGAAATATTGGGCCGCATATTGTGGCATTAAATGTGCTTCTGTAGTC
 ACGGGGTGAAACATAGAAATACCATAATTTTCGTATGGTAAACCGTAATATTCTTTGACTTCTTCTAAGGATGGGAGGGT
 10 GGAAGAGGGCATAACATCTAAATCGGGGGAGCCGATGATGTGAATATGCTTTCTTTTCTCCCATTTGCAGTACGGCCGAG
 TGACAGCTTTGTTTCAATTTGCTACCAAGTGGATATGAGAAAGTTTACTAATAGAATGACGAATGGAGTCATCTACTGTACCA
 GATAGTTCACCACCTTCGATATGGCAAACCTAAACGGCTGCTTAATGCACCTACAGCTGCGCCTGCTAGTGTCTTCTAAACG
 GTCGCGGTGAATCATGACCATATCAGGTTCAATTTTCATCAGATAGACGAGAGATAAACGTAATGGTATTGCCTAAAACGG
 CACCCATTGGTTTACCTTGGATTTGATTTGAAACAGATATGTATGTTGATAGTTTCTCGAGTTACTTCTTGTAGGTT
 15 CTGCCATATGTTTTATCATATGCATACCAAGTTACAATCAAATGCAATTCAGGCTCTGGGTGATTTTCAATATAGGCTAA
 TAAAGGTTTTAGCTTGGCAGGTCGGCTCTGGTACCTGTAATGCAAGAATCTTTTCATGATTTTAGAATCTATAAGTA
 TATATATATAAGTATAAGGAAGTTGGAAGAAGAATACTAATTATACTCTACGTACTCATAAATTTATTTTCGATTAAAGT
 CTATAATTAGGCCATTATAATTATATTAGGATTGGCTT

SEQ. ID NO:33

15 **Nucleotide sequence of DNA region (1000 bp) up-stream from the lgtF gene from *Neisseria meningitidis* (serogroup A)**

20 TCTTTTTCGGACTGAAAGGACGCATCATCCGACATCGAGCGCGTGTCTCGCCGACGCCAAGGCATAGGTTATGCCTAC
 GAAGCCATCAAAATACGGTCTGACCGATATGATGCTGGCGGGCGGAGGCGAAGAATTTTCCCGTCCGAAGTGATGTTTT
 CGACTCGCTTTATGCCGCCAGCCGCGCGCAACGGCGAAGCGGAAAAACCCCGGCCCATACGACGCGAACCAGCGACGGGC
 25 TGGTCATCGGCGAAGGCGCGGGGATTTTCGTGCTGGAAGAATTGGAACACGCCAAACGGCGCGGTGCGATAATTTACGCC
 GAACTCGTCGGCTACGGAGCCAACAGCGATGCTTACCATATTTCCACGCCCCGCCCCGACGCGCAAGGCGCAATCCTTGC
 CTTTCAGACGGCATTGCAACACGAGACCTTGCGCCGGAAGACATCGGCTGGATTAATCTGCACGGCACCAGGAGCGCACC
 ACAACGACAGTATGGAAGACCGCGCGGTTGCAAGCGGTTTCGGCAACAATACGCCCTGCACGTCCACCAAGCGCAAAAC
 30 GGACACACGCTGGGGCGGGCGGGGCAATCGAAGCCGCTTCGCGTGGGGCATTGCTGACCGGAAAAGCAATCCCGAAGG
 GAACTTCCGCCCGAGCTTTGGGAGCGGCGAAGCATCCCGACCTTCCCGCCATCAACCTGACCGGCAGCGGCGAGCCGCT
 GGGAAACCGAAAAACGCATTGCGGCCAGCTCGTCTGTTGCTTCGGAGGAAGCAACTGCGTTTTACTCATCGGATGAAAT
 AAGTTTGTCAATCCACCGCTATGCTATACAATACGCGCTACTCTTGATGGGTCTGTAGCTCAGGGGTTAGAGCAGGGG
 35 ACTCATAATCCCTTGGTCTGGGTCGAGCCCCACCGGACCCACCAATTCCCAAGCCCGGACGTATGTTGGGCTTTTTT
 GCCGCCCTGTGAACCAAAATGCTTTGAGAAACCTTGATA

SEQ. ID NO:34

30 **Nucleotide sequence of DNA region (1000 bp) up-stream from the lgtB gene from *Neisseria meningitidis* (serogroup B)**

35 TAGAAAAATATTTCCGCCAATCATTAGCCGCGCTCGTGAATCAGACTTGGCGCAACTTGGAGATTTGATTGTGATGAC
 GGCTCGACAGACGGTACGCTTGCCATTGCGCAAGGATTTCAAAGCGGGACAGCCGTATCAAAATCCTTGCAACAGCTCA
 AAATTCGGCCCTGATTCCCTCTTTAAACATCGGGCTGGACGAATTGGCAAAGTCAGGAATGGGGGAATATATTGCACGCA
 CCGATGCCGACGATATTGCCGCCCCGACTGGATTGAGAAAAATCGTGGGCGAGATGGAAAAAGACCGCAGCATCATCGCG
 40 ATGGCGCGCTGGCTGGAAGTTTGTGCGAAGAAAAGGACGGCAACCGGCTGGCGCGGCATCACAGGCAGCGCAAAATTTG
 GAAAAAGCGACCCGGCAGGAATATTGCCGACTTTTCCCTTTCGGCAACCCCATACACAACAACACGATGATTATGA
 GGCGCAGCGTCATTGACGGCGGTTTGCCTTACAACACCGAGCGGGATTGGGCGGAAGATTACCAATTTTGGTACGATGTC
 AGCAAAATTTGGCAGGCTGGCTTATTATCCCGAAGCCTTGGTCAAATACCGCCTTCACGCCAATCAGGTTTCATCCAAATA
 45 CAGCATCCGGCAACAGGAATCGCGCAAGGCATCCAAAAACCGCCAGAAACGATTTTTCAGTCTATGGGTTTTAAAA
 CCCGGTTTCGACAGCCTTGAATACCGCCAAATAAAGCAGTAGCGTATGAATTGCTGGAGAAACATTTGCCGGAAGAAGAT
 TTTGAACGCGCCCGCGGTTTTTGTACCAATGCTTCAAACGACGAGACACGCTGCCCGCGCGCTGGCTGGATTTTGC
 50 GGCAGACGGCAGGATGCGGCGGCTGTTTACCTTGAGGCAATACTTCGGCATTTTGCACCGATTGCTGAAAAACCGTTGAA
 AACGCGCGCTTTATCCAACAGACAAAAACAGGATAAAT

SEQ. ID NO:35

50 **Nucleotide sequence of DNA region (1000 bp) up-stream from the lgt gene from *Neisseria meningitidis* (serogroup B)**

55 GCGCACGGCTTTTCTTCATCGGTTTGAAGGTGCGCAGGATAATCGGGGACGGCAAGCCTTTAGACTGCAATTCCTTAA
 TCGCGCGCGGTGAGTTGAGGTACGGATGCGCTGATGTTGCGGAGTTTGATTACGTTTGCATCGGGCTGTTTACACAGTTTCG
 CCAATTCGGCAAGCGCGTGGGTACGCGCTGCGCTTCGGTCAGATATTGCGGGAATGCGGCCAAAAATACGGCCGGACAG
 60 GGAATGTGCGCAGTTTTCATCAATATCGGCGTGGCGGGCAACGCCTGCACAATCGGCAGCAGCGATTGGGTGCGCA
 GCGCGGGGGCTTCGTGGTATGGGTATAAACAATGGTGATTTTTCAGTATAGGATTATCTCTTGTAGGTTGGTTTTT
 TCTTTTGGAAACATTTGCGCGGGGAATGTGCGCGGCTATTATGGCATATTTTGGCGGCTTTGTTGCGCTTTGTTGATC
 TTGGCGTGTGTAACGCGGCGAGCGTGAAGGAAGGGGAATGGTTTTCCCGCGTTTGGCGGCGGTGTCGGAGGTGCTGT
 65 GCCTGATGTGCGGCGGCATATTTTCGGTGAAATGATTTTATAGTGGTTTAAATTTAAACAGTACAGCGTTGCCTCGCC
 TTGTCGTACTATCTGTACTGTCTGCGGCTTCGTTGCTTGTGCTGATTTAAATTTAAACCACTATAATATTCCGTTAATC
 70 TCGGAATATCTGCTAAAATTCGCAATTTTCCGCCCTCGGACACTCGGGGCGTATGTTAATTTGTCGAATGGAGTTT
 AGGAT

SEQ. ID NO:36

65 **Nucleotide sequence of DNA region (1000 bp) up-stream from the msbB gene from *Neisseria meningitidis* (serogroup B)**

70 GCCGACGGCGAAGACAGACGCTCGTGAATCAACCGCTTGGACAGTACGGCGGCGCAATACGACATGCTTGACGTTATC
 TTGAAAGACTTGGCGGAAAAACCGACGCTTGGCGTGGCGCTACCGCCAAATGCGCTCTGAACACCCGATTATCCTTTT
 GAAAGCGGATATGCCCCATACCTTCCCGATATTTCCCAATGTATCAGACAAAATTTGGAACAATATTTCAAAGACCT
 GAACGGTACCGAACCTTGGCGGCTGACGATATGGTCTGCATCAGGTGGAAAAACCGCTGCTGGTGTGCGTGGGAAC
 AATGCGGCGGCAACAGTCCAAAGCCTCCGTCATGTTGGGACTGAACCGCAATACTTTGCGTAAAAAATGATTCAACAC
 GGTTCGTGTAATATGTGCGCAACCGTCCGATCTTTGGGATTTGACCCGGGCGAGTCGCGTAACGGGTTTCGGTGTCTC
 75 GATGTGAGGGGCGGATCATTTTACGTCGCTCCGGCTGCATCAAAACGCTGCCGATGCGCCTCTGGCAGACAGGAT
 TGCCGTGATTGTGCGGCATATCGCGAAGTCTTACCGTTTACAAGCCTCAACAGGCGGCGAGTGGAAACAGGTGTTCTGTC

5 ACGTCAATCCGGCATCGACGCTGATGCTCGGTGAGGCTAGGGGCGCGGCATTGGCGGCATTGGTCAGCCATAAGCTGCCC
GTTTCGGGAATACAGGCCTTGACGGTCAAACAGGCGGTAGTCGGCAAGGGCAAGGCGGCAAAAGAACAGGTGCAGCATAT
GGTGGTGCAGATGCTGGGGCTTTCCGGAAACGCCGAGGANTGGCGGCGGACGGTCTTGCCGTGCGCTGACCCACGCCTT
ACGCAACCACGGGCTTGCCGCCAAACTCAATCCTTCGGGGATGCAGGTCAAGCGCGCAGGTTTCAATAGTTTCAGACGG
CATTTGTATTTTGCCGTCTGAAAAGAAAATGTGTATCGAG

SEQ. ID NO:37

Nucleotide sequence of DNA region (1000 bp) up-stream from the htrB gene
from *Neisseria meningitidis* (serogroup B)

10 CCGCCAAGCGTTTCCCCCTTTGTGCGGGCTTAACATTTGCTTTGTACGGCAGACTTTTCCCTTCATAACGCCGCTTTCC
GAAAAGACGATGGTAGGCGCGACGTAATTCTCAACCTTAAGGTACGGTTGGACGAAAAGTTTCCCTTTTCATTCCACCT
GCCAATTTTTCGGGTACACCGAGTGGTCTCGTTAGGTTTGGGCGAACTACGCCCTTAAAAAACGGACATTCTTTGCATG
CCCCTCTCTAAGGTTTCACGGTAAGTTTACCCTTATAAAGAGTTGACTTACCATACTTATCCCTTTAAACGATATAAAG
15 GCGACAGCTGTAATACAAGTATGTTGTACGGCAGACTTCTTCTACCAACAAAAAGTTCCTTTTAGAGTTACTCGCTTA
TAGACAAATGAAGGCTTAGCCATAGGCTTCCGGTAGGCTATTCAACGGCTGGTTACAGGGCTACGCTAAAACCTACGG
TAGAACCCGCTTCTGGGGTTTCGCGCACAGCGCGCTCTTTGGAACAGTTGTGTCGGAACACGCATAACCGCCCGCTTTA
ATGCGGTGGCGGTTACCTGATGTAGTTTCAGCGTGGCTTGGTAGTTTGGCTAGCCGATGTTGAGGAGGCTCGACC
CGAACTACGGTTGCCAGCGCCAGCGCACATGATGCTGGTTCGTTAGAGGCTGTAGCGGGTCCGCACTTGCTTCGG
20 CTTCCGTAATGAACTTGGTTCCGCGACCGCTGGTTCCAAATACAAGCCGATACGGACGCTGCTTTGGGGCTGGGACTA
CGGCAACCGGTAGATAATGTCGGTGGCGGACTACGTCGCAGTTTCGCTTAATGCGTTTCTGCGGAGGACGGAACCGACG
CAGGGCTGCGTTTTCGGGTGACTGGCACCATAATGCTATCGCTTAGGCCGTTTCATTTTGGCTAACTATGGCAGCAGGAG
AGATACGTTGTGCTGGGCTTTAGCCAATACTTCTCACT

SEQ. ID NO:38

25 Nucleotide sequence of DNA region (1000 bp) up-stream from the MltA gene
from *Neisseria meningitidis* (serogroup B)

30 CACAAAAACCAAGTTATGACGGGAATAAGGTACAGCAGCCAAACCAAGGCTCGGCCCTGC
GTCGGATGGTTCGGTATAGCCGAAAAATCCGCCGAGCAGCAGCCCAACGGCTGTCTTCG
TGCAATATTTTGATGAGTCGAACACAATGTCTGAAGCGGCTTCCAAATGCCTGCTTCG
TGCAGCGCAGCAGCGAACCGGCAAGCAGACCAGCGGCAACGATAATCAGAAACGCCCT
GTCCAAACGAAAACTTCGCCAGATTACGGCGCATCCACCTGATAAATCAACCGGCCA
ATCAGGGCGGCAGCAAAACCCCGCTACCGCACCGCGCGGATCTGCCACGTGCGGCTC
TGTTTGAATACGGCAAGCAGGAAAAAACGCTCTCCAAACCTTCGCGCGCCACGGCAAGA
AACCCCATACCGACCAAGGCCCATCCTTGACCGCTGCCACGGTTCAAAGCCGCTGCACA
35 GAATCTGAAGCTGCCGCTTCATCGAACGGGCGGCTTTTTTTCATCCATAAAATCATATAA
GTCAGCATCGCGACAGCAACCAACCGATAATGCCGACGACGAACTCCTGCTGCTTCTGG
GGAATCTCGCCGTTGCGGAATGGATTCCGTACCCAGCCCCAAACACATCAAAGAAGCA
AGAACAACCCCGCATCAGCTTAGGCATCAGTTTGAATGTCCGGACTGTTTCAGAAAA
CCGGCAACGATGCCGACGATGAGCGCGGCTTCGATACCTCGCGCAACATAATTAAAAAA
40 GCGACCAGCATAAACGCGAACGAACAAGGATGATGAATAATATATTCGGAATATTTTC
ATTGCTTGTAAATACAAATGCAAGTTATTTTTATCTGCAGTACCGCGCGGCGGAAAGTTC
CGCAGCTGCAGCTGCGCCCTGTGTTAAATCCCTCTCCACGGCTGCGCGCAACGCGGCC
GAAACCATCTTTCTTATTACTGCCGGCAACATTGTCCATT

45 SEQ. ID NO:39

Nucleotide sequence of DNA region (1000 bp) up-stream from the ompCD gene
from *Moraxella catarrhalis*

50 GCTGATTTGTGAGCAAGCGGGCGCATCAGGGATTACCTTGCAATTTGCGAGAAGATCGTCG
ACATATTCAAGATGAAGATGTTTATGAATTGATTGGGCAATTGACAACACGCATGAATCT
TGAGATGGCAGTCACTGATGAGATGCTAAATATTGCCCTAAAGGTACGACCAGCATGGGT
GTGTTTAGTACCAGAAAAACGCCAAGAGCTGACTACAGAAGTGGGCTTGATATCGCCAA
TTTATCAAATATTCAAGCATTTATACACAGTCTTCAGCAGGCGGATATTAAAGTTTCTTT
ATTCATCGATCCAGATCCGCATCAAATTGATGCTGCAATTGCTTTGGGTGCTGATGCGAT
55 TGAGCTGCATACGGGAGCTTATGCTCAAGCGACTTTACAAAAATAATCAAAAGCTTGTTGA
TAAAGAGCTTGACCGTATTCAAAAAGCCGTTGCAATGGCACAAAAAAATCATCATTATT
GATTAAATGCAAGGTCAATGTTTACGCGTGATAATGTTGCAGCGATTGCCCAAATTGATGG
TATTCAGCTGAAATATCGGGCATGCATTGATTTACAGATGCGATATTTATGGGCTTGA
TAATGCAGTCAAGGCAATGAAAATGGCTTTTATTCAAGATAAACGACCAATCATTGATG
CGTTAGAAAGAAAAATCGTAAATAATGATGACTATTGTGAATATTATGATTTTTGTTCA
60 AAAAAAGGTTGTAAAAAAATTCATTTACCATTAAAGCTAAGCCCAAGCCACAATGAATA
CCTATTGGTTTACTCATTAGTCACTAAGAATCTGCAAAATTTGTAACAGATTATTGGC
AGGTCTTGGATCGCTATGCTAAATAGGTGCGGTAATCTTGA AAAACCAACCATTCCTTG
GAGGAATTTATGAAAAAGGATATAAACGCTCTTGCGGTCACTCGCAGCGTTGACGCTCC
AGTTGCAGCTCCAGTTGCTGCTCAAGCTGGTGTGACAGTC

65

SEQ. ID NO:40

Nucleotide sequence of DNA region (1000 bp) up-stream from the copB gene
from *Moraxella catarrhalis*

70 GATGCTGTAAAGTGGGTATTGGTCTGTTCTATTTGTACAACCCGATTGTTGCAGGC
ATTGGCGTCCCGCAGATAAGTGCCATTGATAGTGTGGCAAGTGCGTTAAAGATCGCATT

SEQ. ID NO:41

20 AAAAAGCTGGTGATGTCTTCACTGCTATTTCATGGTGAACCAATCAATGATTGGCTAAGTGCC
ACCAAGATTATTTCAGGCAAAATCCAGAAACCATCGTGTGATGTGACAGCTATGCGTCAAGGT
AAGCAGGGTGTATTAAAAATTAATGCCCCGTGGTGTAAAGACACAACACGGCGTAGTCGGT
CAACTGGGTATTGCCCCAGATTGATATCGATACGCTCATTCTGATGAATATCGTATG
25 ACGATTCAATATGATGTGCGTGAGGCACTTACTCAAGCCATCCGACGAATCTATGATTTA
TCATAATGACCTTAGATCGGTAGGTAAGATGATTACAGGATGATTGGCATGAAAAAT
CTATCAGGTCCCATTGCCATTGCCGATGTTTCTAAGACCAAGTTTGGATTGGGATTTCAA
GAAGTGTTATCGACAGCCCAATCATCAGTTTAAAGCTGGCAGTCACTGAATCTTTTACCC
ATTACAGTTGTAGATGGCGGGCACTTTGGTATTTTATACCTATGAATGGATTATGGGCAAA
TCTATGAATGAAGCGGTCGAGATGGCAGCAATTTAAAGCGGGTGCCTTATGCTTTTGT
30 TCTATGTACTTGCAATCGACTAAGCAATCATGCGATTTTGGCTAAGTTCTGATTAT
CGTACCATTAAACAAAATTTTGGCTTTTTTAAGCTGAAATACCTGGCCAAATTAACATTTT
TGGCTTACCTTTACACAAATAAAATTTGGGTGTAGAAATTTTGGATACATTTTATACCA
TTATTTTGTAGAAATTTTAAAAATTAAGTTTGGATGAAGCTATGCGCTAATCATATTTAA
AGGTTTTCAGGTCAAGTGCATGACAATGGCTGTCATGATGGTAAATGTCAACTCATGCACA
35 AGCGCGGGATTTTATGGCAAAATGACATGGCCATCAGGAGCTACAGCGAGTGACCATTTGA
AAGCTTACAAAGCGGTGCTGCCGTTTTCGCTTGGGTCAAGTG

ACTTGGCGAAAATACCATTATATCGATTGTGATGTTATACAGGCAGATGGCGGTACACG
CACAGCCAGTATCATGTGGCTGCTGGCGGTGGCACTATTGATGCTTTAGAACAATTCGACGCT
TCGTA AAAAGCTTACCCAGATCCGCTTTTGGGGCTGGTGGCAGCGGTTCTGTGGGTGT
TAATCAAGGCCGTGTATTGCTTGATTGGATTATGCTGAAGATTCAACTTGTGATACCGA
45 TTTAAATGTTGTCATGACGCGAGCGAGGTGGGTTTATTGAGATTCAAGGCCACAGAGA
AAAGCCATTATTCTCGTGGTGAAGTAAATGGGATGCTTATTGGCAGAGCTGGGAATTGG
GCAGATTATCGAAGCCAAAAGCAAGTATTAGGCTGGTGATATGCTAATCGTTGAAGATA
ATGGCGTGATCATCATTAATGGGACAAAGTAAAGACCCATTATTTGGTGGTGCATGA
TATTGCTGCTGCTGGGTGCTTGGTGGCAATCACTTTGGTTGATGACCCGTTTTTATG
50 CAATCGGTGGCTTGGCTTTATTGTGAGTTGTGGTATTGTGTTTAAATATCAAAGGCCAA
AAGCCAAAATGTCATATGTTTTCACAAGGTGCGTTGAAGATTACGTCCAAACGCTTTG
AGATTCAATAACAATCACTAACCCTTACAGCATCGGCAACAATATCTGCTAAAGATAACA
AAATGAATATGTTGATCGGGCGAGTGAATATCACTTTACAGGTTTGTGATGACGGTGT
AAATTGAATATGGCAAAACAGGTACTTTTGGGAAGTCAATCAAACCAATGTCGGTGGCGG
55 TAACATTGGCTAAGTAGTTGTTGTGATACAGACAGGTGGATGGTCTTTAACTCCACCCA
CCTAACCTTTCTTTGTTGGATTAAAGAGTATGTTATGATGGGCAGGATTTATTTTAA
GTCACTATTATGATGCAATTGCTGCAGAGATGCGGTTTC

65 GTGATCGGCAACACCCACCATTTCAGGAGCAACCAAAATTGCCCGTGCCTTGCCGTGCTT
GGTGGTATCATTTGGCAGGGCAATGTGGCTAAGTAGTGGTGTCATCAGGTGCGGGTGGT
GGTGAGTGTACGATTCTGTATTGTCTATAAAATTATCTTTGGGTGGATGATATCAATG
AAATACCTCTACGGTTGTATGGAATTTTATCCATTGTACACCGGATATGGTCTTTTAAAT
TAACAAGCAGCTTCTAGCAAGTCAAAGTTTTTATGCCTATTTTTTCAGATTTTAAAGGTAC
ATAAAGCCCAATTGTTAATAATATGGTATTGTCTATGATTATGATGAATTGCGACCAAAAT
TTTTGGGAAAATTAATCCCTAGATGCGGTTAACAAGATGCTGAATGGGAAGCATATGTGAC
70 GGATGTGGCGCGTGTGTTTGGTGAATTTCTTGATGATGACAATGTTAAATTGACCGAA
TATACCGATGTGCGTCGCAGCTATTGGATTGCTCAACAGGATTTGGCCAAACTGATGCC
AAGCGTCAAACGATTGTGCGAGATTGTATTGCGTTAAACCTGATATGCTGCTGATATG
CTGTGGTTGCCACGCCATTGTGCTTTAAAGCGGTTGTATCTGGGCAAAATCTGCCAGCA
TGGCACAGGCTCATTAATCATAGCCAAACCATGGTGCGAGGATTTGCGAAATCTTCAACT
GCTGGGCGATGTGTGAGTGAGCTTGGTATGAGTGAATGAAGACATAGAAGGCGAGTGGTG

AAATGGGTTAAACCTTGACATGATTGTTGACATGATTGACAGACAATAAAATTTGGCAAA
 TTTGATAAAATTTGGTGTATGTGTGATTTTATCAAAAGCACTTGAATAAAACCGAGTGA
 TACGCTAAATTTGTAGCAAAACCAATCAATTCATCATAATTTTAAATGAACACGAGGTTAAAT
 TATACTGTCTATGTCTGATGACAATTCAGCACTTGGTCTG

5

SEQ. ID NO:44

Nucleotide sequence of DNA region (1000 bp) up-stream from the lbpA gene
 from *Moraxella catarrhalis*

10 TAACAAAGGCAACCAACACGCGAGTTATTTGTGCAAGGCGGTCAAGCGGATGTCAGTAC
 TCAGCTGCCAGTGCAGGTAAATTCACCTATAATGGTCTTTGGGCGAGGCTACCTGACCCA
 GAAAAAGACAAAGGTTATAGCAAGATGAGGATACCATCAAGCAAAAGGCTCTTAAAGA
 TTATATATTGACCAAGACTTTATCCACAAGATGACGATGACGATGACGATGACGATGAC
 TTTGACCGCATCTGATGATTACACAAGATGATAATACACATGGCGATGATGATTGATTGC
 15 ATCTGATGATTACAAAGATGATGACGCGAGATGGCGATGACGATTCAGATGATTGGGTGA
 TGGTGCAGATGATGACGCCGAGGCAAGTGTATCATGCAGGTAATATTGCCCTGAATT
 TGAAAAACAATACTTGGCCATTAAATGAGCCTACTCATGAAAAAACCTTTGCCCTAGATGG
 TAAAAATAAGGCTAAGTTTGTATGTAATTTGACACCAACAGCCCTAAGTGGTAAATTTAA
 CGATGAGAGAGGTGATATCGTCTTTGATATCAAAATGGCAAAATTTGATGGCACAGGATT
 20 TACCGCCAAAGCCGATGTGCCAACTATCGTGAAGAAGTGGGTAACAACCAAGGTGGCGG
 TTTCTTATACAACATCAAGATATTGATGTTAAGGGGCAATTTTGGCACAAATGGCGA
 AGAGTTGGCAGGACGGTTACATCATGACAAAGGCGATGGCATCACTGACACCGCCGAAAA
 AGCAGGGGCTGTCTTTGGGGCTGTAAAGATAAATAAAGCCCCCTCATCATCGTTTAST
 CGCTTGACCGACAGTTGATGACGCCCTTGGCAATGTCTTAAACAGCACTTTGAACAGT
 25 GCCTTGGGCGAATTTCTTGATAAATGACCCAGATTTGCCTCGGGCTAATATCTTGATAAA
 ACATCGCCATAAAATAGAAAATAAAGTTTAGGATTTTTTT

SEQ. ID NO:45

Nucleotide sequence of DNA region (1000 bp) up-stream from the lbpB gene
 from *Moraxella catarrhalis*

30 CAGCTTGACCATTTGGTGAATATATACCATTTGGTGGTTTGTGGATATTTACCAGGG
 CTTGAGGGGTGTCGTAGCCTAAGCCGTGGCGATGATAAGCAACCACCGCTCAAATTTGGGC
 GGCGGCGTGGCGCATACGATTGGTGGCGCAATTTGTTATGAGGTGGCATATCCTGAGACG
 ACGCGTAAAAATGCCTTGGCAGTAATTTTTTATTAACCGTCTCAAACGATGCTTGGTTT
 35 GGTACAACAGCAGGTCTTTGCAGCATTACAAATGGTGCAAAATGCGAAGCTTGGAGACG
 GGGCGATGGTTTGTGCGGTGCAACAAACAACGGAGTGACTGCATTAATTGACCATCAAGGA
 CGGATTATCAAGCAGATACCGCAGTTTCAGCGAGATATTTGCGAGGTGATGATCCCAAT
 TATGTTGGACACACGCCCTTATATGTTTGGGGGCATTATCCCATGTTGGGGTTTTCTTTG
 GTGCTGATTTTTCTTAGTATCATGGCAAAGAAAATGAAAAATACCACCGCCAAACGAGAA
 AAATTTTATACCGCTGATGGTGTGGTAGACCGCTGAATTTGTCCACTTTGGGCGTTAGAG
 40 CATGAGCAAGATTAGGCGTTGGGTGAGCTTTGGTTGTATTACTCATCAGCCTACCCGAAA
 CCTGCCAAACATCACCGCCCAAAACCTAAACATACAATGGCTAAAAATATCAGAAAATAA
 CTTGCTGTATTGTAATTTCTTATGTTATCATGTGATAATAATTATCATTAGTACCAAGAT
 ATCCATTACTAAACTTCAATCCCCCATCTTAACAGTTACCAAGCGGTGAGCGGATTATCCG
 45 ATTGACAGCAAGCTTAGCATGATGGCATCGGCTGATTGTCTTTTGCCTTGTGTGTGTT
 TGTGGGAGTTGATTGTACTTACCTTAGTGGTGGATGCTTGGGCTGATTTAATTAATTTG
 ATCAAAGCGGTCTTCAACACACCAACGAGATATCACC

SEQ. ID NO:46

Nucleotide sequence of DNA region (1000 bp) up-stream from the tbpB gene
 from *Moraxella catarrhalis*

50 AGTTTGGCCTGATTTTGAGAGCCACTGCCATCATGAATTTGTTGGCGTAAACACCACTCG
 TATCTTCTTCGGTTTCCCTTTCCATGCAAAACACAGGGATACCAGCGGCCCGCATGGCA
 GCGGCGGCGTGGTCTTGGGTGCTAAAAATATTGCATGATGTCCAGCGAACTTCTGCACCC
 AAGGCAACCAAGCTCAATCAGCACCGCTGTTTGAATGGTCATGTGGATACAGCCTAGG
 55 ATTTTAGCACCCCTTAAGTGGTTGCTGGTCTTGATAGCGTTTCTTAACCCCATCAGGGCT
 GGCATCTCAGCTTCTGCCAAGGCAATCTCACGGCGACCATAATCGGCTAAACGGATATCA
 GCGACTTTATAATCGGTGAAGTTTGGGTGGTACTTGGATTGATTGAGGTAGGCATATCT
 TTATTCCTAAGCTATTTTAAAGTATTTTAAACAATAATTTTATGAATTTGAGATAATTG
 ATGCTAAAGGTTGAATGACCAACCATCGCTAACCAATCAAGAAAAGACATTTTAAGCAT
 60 AAAAAAGCAATGTGTCTTGATGGCTTATTATAACAGTTATTATGATAAATTTGGGTAGAA
 AGTTAAATGGATCGTTGGGTAAAGTTTGTGGCTATCCTTAATTAATTTTAAATTTTAAAT
 AATGCTTTTACTTTTATTTTAAAAATAGAGTAAAAATGGTTGGCTTTGGGTTTTTATCTC
 ACTATGGTAGATAAAATGATACAAAATGGTTTGTATTATCACTTGTATTGTATTATAA
 TTTTACTTATTTTACAAACTATACATAAAATCAAAATTAATCACTTTGGTTGGGTGG
 65 TTTTAGCAAGCAAAATGGTTATTTTGGTAAACAATTAAGTTCTTAAAAACGATACACGCTC
 ATAAACAGATGGTTTTTGGCATCTGCAATTTGATGCCTGCCTTGTGATTGGTTGGGTTGT
 ATCGGTGATCAAAGTGCAAAAGCCAACAGGTGGTCATTG

SEQ. ID NO:47

Nucleotide sequence of DNA region (1000 bp) up-stream from the *tbpA* gene from *Moraxella catarrhalis*

5 TTGGGGGCGGATAAAAAAGTGGTCTTTGCCCAAAGGGGCATATGTGGGAGCGAACACCCAA
ATCTATGGCAACATCATCAAAATCACAAAAATACAACGACCATTTGGGGCAGACTGGGG
GCAAATTTGGGCTTTTGGCTGATGCCAAAAAGACCTTAGCATTGAGACCTATGGTGAAAAA
AGATTTTATGGGCATGAGCGTTATACCGACACCATCGGCATACGCATGTCGGTTGATTAT
10 AGAATCAACCCAAATTTCAAAGCCTAAACGCCATAGACATATCAGCCTAACCAACCAT
CGGACGCCAGGGCTGACAGTAATAACACTTTATACAGCACATCATTGATTTATTACCCA
AATGCCACACGCTATTATCTTTGGGGGCAGACTTTATGATGAAAAAGTGCCACAAGAC
CCATCTGACAGCTATGAGCGTCGTGGCATAAGCACAGCGTGGGGGCAAGAATGGGCGGGT
GGTCTTTCAAGCCGTGCCCAAATCAGCATCAACAAACGCCATTACCAAGGGGCAACCTA
15 ACCAGTGGCGGACAAATTCGCCATGATAAACAGATGCAAGCGTCTTTATCGCTTTGGCAC
AGAGACATTACAAATGGGGCATCACGCCACGGCTGACCATCAGTACAAACATCAATAAA
AGCAATGACATCAAGGCAAAATATCACAATAATCAAATGTTTGTGAGTTAGTCGCATT
TTTTGATGGGATAAGCACGCCCTACTTTTGTGTTTGTGAAAAAATGTGCCATCATAGACA
ATATCAAGAAAAATCAAGAAAAAAGATTACAAATTAATGATAATTGTTATTGTTTAT
GTTATTATTATCAATGATAATTTGCCGTATTTGTCCATCACAACGCCATTATCATCA
ATGCCACAGACAAATACGCCAAATGCACATTGTCAACATGCCAAATAGGCATTACAGAC
20 TTTTATGATAATACCATCAACCATCAGAGGATTATTTT

SEQ. ID NO:48

Nucleotide sequence of DNA region (1000 bp) up-stream from the *ompE* gene from *Moraxella catarrhalis*

25 AAAGACATTACACATCATTTCAAACGCCCAACCATGTACCTCTGCCCCGTGGTCGCAC
GCCAACGCTTTTTGATGCGGTGCGTTGGGTTTCAAGTGGCTTGCAATCATTGGTTTTAT
TAAAATTCATACCTTTGGTAGTTTGGCTTTACCTGATATGTCATTTGATTATCGAAACAA
TACGCAGTTGACCAAAACATCAATTTTAGCCATTGCCAAGCACTCAATATTACCGCTCA
TACGACCATGCTTGGTATTAAATCATCACATAAAGATACTTTACATCCATTTGAATTGAC
30 ATTACCCAAATCAGGGCATGCTCCTAAATATGATGATGAATTGGTGCAAAACATCCATT
GGCTTATTTTTCATCAACTGTCTGCCGTCTGCCGATATTTTATACCAACCGTTTGTAT
TGTTGGCGGTGAAAGCTCAGGGAAACTACCTTGGTGCAAAACTTGCCAATTATTATGG
TGCCAGCATCGCACCTGAAATGGGTGCGATTATACACACACTCCCATCTCGGGCGTAGCGA
ACTTGGCCTTCAATACAGCGACTACGCATCCATTGCCATCAATCAGGCCAACGCTATCGA
35 AACCGCTCGTACCACTGCCAGCTCTGCTGTACACTGATTGATACTGATTTTGGGACAAC
GCAAGCATTTTGTGAAATTTATGAAGGGCGAACGCATCGCTTGTGCGAGAATTTGCTAA
ACAAATGCGATTGGATTTTACGATTATTAGATAAATGTTGCTTGGGTGCTGATGG
CATGCGTAGGCTTGGTGATGATCATCAACGCAGTTTGTTCGCCAATAAATTGCTTGAGAT
TTTGGCACGATATGATATTAGTTATCATATCATTATGACACCGACTACCACAAACGCTA
40 TCTACAAGCATTAAAGCTTGATAGACAATCATATTTTAAATCATTTTACAAAAATTCATGA
CAATTAATTAGGAAAAATCTGATGAAAATTGATATTTTAG

SEQ. ID NO:49

Nucleotide sequence of DNA region (1000 bp) up-stream from the *uspa1* gene from *Moraxella catarrhalis*

45 GGATGTGGCATATCTGCCATCGACCCAATACACATCGGTGCGAGGCTATCAAGATGTGGT
ACGAATTAATAGCCAGTCAGGTAAGGGCGGTGCTGCGTATATCTTGACGCGGCATTTTGG
TTTTAATTTACCACGCTGGACACAGATTGATTTTGTCTGCTGTTGGTACAGGCTTATGCAGA
AAGTATGGCGCGTGAACATAAACTGATGAGCTGCTTGAATTTTACCCAAGCGTATCT
50 TAAGCAAGATAAATTCGCCCTAAGTGACTATACCATCAGCAATAAAGGCGATGCTGTCAG
CTTCCAAGGCCAAGTAGCGACACCCAAAGCGGTGTTGAGGTGATTGGTCAAGGCAATGG
TGCGTTATCTGCGTTTCATTGATGGCTTGGTGAAATCCACAGGCGAGACAGATTGATGTCAC
CAATTACGCCCGAACACGCGATCGATAACAAAACCCATCAAAAAACCGATACGGATAACCA
55 AACCGATGCCGCGGTGCCGCTTATATCCAGCTGTGCGTAGAGGGGCGAGTTTATTCAGGC
ATCGCCACTTGCCATAGCACCGTATCCGCCATGCTAAAAGGTGCATTATCCGCTTTGGCA
CAGGCGTGGTAATCTGACCCAATCAAAATCCTGCATGATGGCAGGATTTTATTATTAGT
GGGCTGCCCAACAATGATGATCATCAGCATGTGAGCAATGACTGGCGTAAATGACTGAT
GAGTGTCTATTATGAAAGATATCAATATATAAAGTTGACTATAGCGATGCAATACAG
TAAAATTTGTTACGGTTAAACATAACGACGGTCCAAAGATGGCGGATATCGCCATTTACCA
60 ACCTGATAATCAGTTTGATAGCCATTAGCGATGGCATCAAGTTGTGTTGTTGATTGTCA
TATAAACGGTAAATTTGGTTTGGTGGATGCCCATCTGATTTACCGTCCCCCTAATAAGT
GAGGGGGGGGAGACCCAGTCATTATTAGGAGACTAAG

SEQ. ID NO:50

Nucleotide sequence of DNA region (1000 bp) up-stream from the *uspa2* gene from *Moraxella catarrhalis*

65 CCCCCAGCTTTCCGTTTGTGTGCTGCTGGTGTGCGGGCGGTACATACCATGCTGGTGGCTG
CCGATGGTGTGGGCGTGTATCAGATATTGGGCGAGTCTATCGATGATGCGGTTGGGTGAAT
GCTTTGATAAAACGGGCAAAAAATGCTCAAACTGCCCTATCCTGGTGGCCCAAAATATCGAAA
AATTAGCCAAAAACGGCAACCCACACGCCCTATGAGCTGCCAAGACCCATGCAGCATAAAG
70 GGCTGGATTTTTCGTTTCAAGTGGCATGAAACCGCCATTCAATCTCATCAAGACACAC
CAAAACGCCCAAGCCCGCCACACGAGCAGACATGCCCGCAAGCTTTGAGTATGCGG
TGTTGGATACTTTGGTCAAAAAATGCACCAAGCACTACAGATGACAGGCATTGCCGAC

5 TGGTGGTCGCGAGGGGGCGTCTCTGCCAATCAGATGCTACGCCGCACCTGACCGAGACGC
 TCCGCCAAATCGATGCGTGGGTACTATGCCCGACCGAGCTATGCACGGATAATGGTG
 CGATGATCGCCTATGCTGGCTTTTGTGGGCTCAGCTGTGGACAGTCGGATGACTTGGCGG
 TTCGGTGTATCCCCGATGGGATATGACGACGCTTGGCGTATCGGGTCATAGATAGCCAC
 ATCAATCATACCAACCAATCGTACAAACGGTTGATACATGCCAAAAATACCATATTGAA
 AGTAGGGTTTGGGTATTATTTATGTAACCTTATATCTAATTTGGTGTGATCTTTGATAA
 AGCCTTGCTATACGTGAACCTAAATGGATATGATAGAGATTTTCCATTTATGCCAGCAA
 AAGAGATAGATAGATAGATAGATAGATACTCTGTCTTTTATCTGTCCGCTGATGCTT
 TCTGCCTGCCACCGATGATATCATTTATCTGCTTTTAGGCATCAGTTATTTACCGTGA
 10 TGACTGATGTGATGACTTAACCACCAAAAGAGAGTGCTAA

SEQ. ID NO:51

Nucleotide sequence of DNA region (1000 bp) up-stream from the omp21 gene
 from *Moraxella catarrhalis*

15 GAGTGAACTTTATTGTAAATATGATTCAATTAAGTATCAAAATCATCAACGCAGCATC
 AGGGTTTGTAAATCAATTTTTTACCATAATTATAGCCATAACGCACAGCAAGCGTAGT
 TATGCCAGCGGCTTGCCCTGATAAAATATCATTTTGGAAATCAACCAACATAATGGCATC
 AGTCGGTGGATGCCCCAGTGATTGACACAGGTATAATAAGGCGTTGGGTGGGGCTTTT
 GACGCTGAGCGTATCACCGCCAATCACTTGGTCAAACAGTGTGAGCCATCCAAATGTGA
 20 TAAATTTTAGGCAAAATAACGCTCAGGCTTATTGGTACAAATTGCCAAATAAAACCCGC
 TGCTTTTAATCGTTCAAGCCCTTGATAACCCCTGCATAGCTTTGCGTATTTTCAATTGT
 TTTATGGGCATATTCTGCCAAAAATAACTCATGGGCATGGTGAATCATAGTCGTATCATA
 GATATGATGTGCTTGCATTGCTCGCTCAACCAATTTAGCGAACCATTTGCCACCCAGCT
 TTTGATGATATCAATTGGCATAGCGGTAAGTTAAGCTTGGCATACATGCCATTGACCGC
 25 CGCCGCCAAATCAGGGGCACTATCGATAAGCGTACCATCCAAATCAAAATATAATCAGTTT
 TTGGCAGCTGACAGTGTGTTGTCATGCTTTTCCCTTATTCTTAAATTTGGCGGCTGTT
 TGGTATTTTTTAAATCAGTCAATTTTACCATTGTGCATATAATGACAAAGTACAAATT
 AGCAATATTTAGTGCATTTTGGCGAAGTTTATGAAAACCTGGTCATTGGTTGCAAAA
 30 CTTTACACAGTACCTATAAACTTGCACAGTTAATAAGAAATATTGTTACTATAGGGG
 CGTCATTTGGAACAGACAGTTATTTGTAATAGTTATTTGCAAAAGACGGCTAAAAGAC
 AGAACAGCGTTTGTTCAGTGATTAAC TAGGAGAAAAACA

SEQ. ID NO:52

Nucleotide sequence of DNA region (1000 bp) up-stream from the omp106 gene
 from *Moraxella catarrhalis*

35 TTGATCGGTTTTGCCCCACTGTTTCATGATTTACTCAAAACAGGCGGCTTGATCGTGCTG
 GCAGGTCTGACCCAAAACCAACCCAAAGCGGTCATCGATGCCCTACTCGCCTTATGTTACG
 CTTGATACGCCATTTTGTATGCAGATGCCCAAGACTGCCATTGGCAACGCCCTAAGCGGC
 ATCAAACCTACCAACCCATAAGCGATATGCCATGAGCCACAAACCTAAGCCAAACCCGCT
 40 ATATCAACAAGTTGAGCAGACCCGCAAGCGTTATTTGAGACATTGGGCGATGCTCATAC
 TCATGATGTCTATGCCACTTTTTTGGCCGAATTTGAAAAACCGCTGCTCATCGCCGCACT
 CAATCACACGCGACGGCAATCAGTCAAAAACCGCCCAATCCTTGGTATCAATCGTGGCAC
 ATTACGCACCAAAATGAAACCCATCACTTACTTTAGACCGCCAGTTATCGCCATGGATA
 TGGGCGAGTGTGCTCGCCTGCCGTATGATGGCGATGACACCCCATTTGCCCATATCTGC
 45 ACGATTTGACATGATTTAACATGTGATATGATTTAACATGTGACATGATTTAACATTGTT
 TAATACTGTTGCCATCATTACCATAATTTAGTAACGCATTGTGAAAAATCATTGCCCCCT
 TTTTTATGTGTATCATATGAATAGAAATATTATGATTGTATCTGATTATTGTATCAGAAT
 GGTGATGCCCTACGATTTGATTTGGGTTAATCACTCTATTATTGATATGTTTTGAAACTA
 ATCTATTGACTTAAATCACCATATGGTTATAATTTAGCATATGGTAGGCTTTTGTAAA
 50 AATCACATCGCAATATTGTTCTACTGTTACCACCATGCTTGAATGACGATCCAAATCACC
 AGATTCATTCAAGTATGTTGTTGATACGCACCATTTACCCTAATTATTTCATCAAT
 GCCTATGTCAGCATGTATCATTTTTTTAAGGTAAACCAAC

SEQ. ID NO:53

55 Nucleotide sequence of DNA region (1000 bp) up-stream from the HtrB gene
 from *Moraxella catarrhalis*

ACTATTCTGCTTTTTGTTTTTCACGAATGCGAATGCCAACTCAGCCAACTGGCGATTAT
 CAACTTCAGCAGGTGCTTCGGTCAATGGGCAATCTGCCGTCTTGGTTTTTGGGAAGGCGA
 TCACATCAGCGATTGAGCTGGCACCACCATCAGCATATCAGGCGATCTAGACCAATG
 60 CCAAACCAACCGTGGCGGCTGCACCAAAACGCAATGCATCCATCAAAAACCTTAACTTAA
 GCTCTGCTTCTTCTTAGAAATACCAAGGCATCAATACCGCCTCTTGATGTCAACCG
 TATTAATACGCAGCGAACC GCCACCAATTTCTGTGCCATTAGTACCATGTCTAGGCAA
 TGGATAGGGCGGTTTCGGGACTTTGTTGAGTTCCCAACCGAGCCTTTTGGGCGTGTA
 AAGGATGATGAACATGATGCCACTTACCATCATCAGTTTCTCAACATTTGGAAAATCAA
 65 CGACCCAAAGCGGTGCCCATTCACAGTAAATAAATTTAAATCAGTACCGATTTTAAACAC
 GCAATGCAACCATATGATCATTTGACGATTTTGGCTTTATCGGCACCAAGAAATGATAT
 CGCCAGTTTGGGCATCGGTACGCTCAATCAGCTCAATCAAAACCTCATCGGTATATTT
 TAATGATGGGTGATTGTATCTGATTCTTTTCAACGCCATTATGATATTGCTTGGCT
 CATTGACCTTAATATATGCCAATCCACGAGCGCCATAAATACCAACAAATTTGGTGTA
 70 CATCAATCTGCTTGGGACTCATGTTACCGCCATTTGGAATGCGTAAGGCAACACACGGC
 CTTTAGGATCTTTGGGCGGGCCCTGAAAATACTTTAAATTAACATGTTGCATGATGTCAG
 CAACATCAATAAGTTTTAAGGGAATGCCGTAATCAGGCTTATCTGAGGCATAATCAGCA

TGGCATCTGCGTAAGTCATGCGGGGAAGGTATCAAACCTCA

SEQ. ID NO:54

5 Nucleotide sequence of DNA region (1000 bp) up-stream from the MsbB gene
from *Moraxella catarrhalis*

10 TGGATCATATTCTTTATTAATGGTACTGTTTAAACCTGTATTTTAAAGTTTATTGGGTCA
TATTTTCAAGCTCATCCCATCGCTCAAGCTTCATCATCAAAGCTCATCAATCTCTACCA
ATCGCTCACCAGCCTTCGTTGCTGCGCCAAATCGGTATTAACCATGAACCATCTTCAA
15 TCTTTTGGCAAGCTGTGCGCTGCTCTTGTTCAGTGCAGCAATTCATTAGGCAAACTCT
CAAGTTCACGCTGCTCTTTATAGCTGAGTTTGGCTTTTGGGCAACGCGTGATTGAGGTG
GTTTGATTTGGATGGGTTTTCAGCGGGTTTGTGCGCTTAGGTTTATTGTCTGTGGCGTGAT
GAGCAAGCCATCTTTTCATGCTGTTGTACATAGTCTTCATAACCGCCAACATATCCAAAA
CGATACCGTCGCGCTACTTATCAGTATCAAATACCAAGTTTGGGTAACAACATTATCCA
20 TAAAAGCAGCGTCATGGCTGATGAGTAATACCGTGCGCTTAAATTTGACCACAAAATCTT
CTAAAAGCTCAAGTGTGCCATATCCAAATCATTGGTAGGCTCATCAAGCACCACAAAAT
TGGCAGGTTTTAGCAATAATTGGCCAATAAAACGCGTGCTTTTTCACGCGCTGATAGTG
CTTTAACAGGTGTGCGAGCAGGATTGGCGTGAATAAAAAATCTTGCAATAGCTTAAAA
TGTGCGTAGTTTTTCACCACACATCGACATGGTCAGAGCCTTCTGAAACATTATCTGCGA
25 TAGATTTTTCAGGGTCTAGGTCGTCTTTGAGTTGGTCAAAAAAGCAATATTTAGATTGG
TGCCAAGCTTAACTGAACCTGACTGAATCGCTGAATCATCAAACCCAAAATGCTTTTAA
TTAAGGTTGTTTTACCAACGCCATTTTGCCAATGATACCAACTTTATCACCACGAACAA
GCAGCGTTGAAAAATCCTTAACTAAGGTTTATTGTCTGAT

25 SEQ. ID NO:55

Nucleotide sequence of DNA region (1000 bp) up-stream from the PilQ gene
from *Moraxella catarrhalis*

30 CAACTTGAAAATCAGCTCAATGCTCTGCCACGCACAGCACCAGTGAAGCGAGATTATCGGA
ATGATAAATACCAAGCACAAGCGGTTAATGTGCAGGTGGTGAGTGCATCAGTTCAAGCA
GGTCGTGAACAGGATTATTATACCGAACGCCCTATCGCAGTGAGTGCGACAGGGGATTAT
CATGCTTTGGGTGCGATGGTTACTTGAGTTGTGAGAGGCTAACCATTTGCTGACAGTGCAT
GATTTTGATCTGAAGGCTGGTTTGAACATCAGCTGATGATGATTGTTGAGATGAAAAT
35 TATCAAGCGAACAACGCCCCAAAACAGTTGCTCAGCAGGTGCGCTGATGTTCAATGAATA
TTATCGGTGGGGCATTTTGGGTGCTTGGATTGGGTTGGGATTGGATGTGCTGATAGCAC
CAGTCAAGTTTGTGATGATTAAGCTTGACATATTAACCATGAAGAGCGTATGGCGATCAG
TGAGCCTGTGCGGATACCTTATCTGTGCGGATGATATATCAGCAAGGCAAGATCCTTT
TATCAATCCTTATAGAAATGTTGAGGTTCTTGATACCAATCATGCCGCTGATCAGCAAGA
40 TGAGCCAAAACCGGAATCTACCAAGCTTGGCCTATGGCAGACACTATGCCATCTCAGCC
ATCTGATACTCATCAGTCTGCCAAGGCTCAGGCACAAGTCTTCAAAGCGATCCGATAGT
CATTGATACCAACCGTGTTCGAGAGCCTTTAGAAAGCTATGAGTTATCAAGCCTACGCTA
TCATGGTCGTATTTTGTAGTATGTTAGACTTGTGGCACTCATTATGAGTCCTGATGGCAT
CGTTTCATCGTGTGAGTACTGGACAATATCTTGGTAAAAATCAGGAAAAATTACCATAT
45 TGACAGTCGTACGATACATCTGATTGAAGCGGTGCTGATACACAAGGTGGCTATTATCG
CGTGATGTAAACATTCAATTTATTTCATAAGCAATGACAC

45 SEQ. ID NO:56

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipo18 gene
from *Moraxella catarrhalis*

50 TTCATGCAACAGCGACCATCTTGGCCGATGATACCATCCTGCTCACCTAAGAAAAATCAG
TTTATCAGCTTTCAGGGCAATGGCTGTGGTCAGTGCTACATCTTCTGCCAATAGATTAA
AATTCGCCCCGTAACCGAAAAACCTGTCGGTCTTAGTAGGACAATATGGTCATTATCCAA
ATTATGGCGAATGGCATCGACATCAATTGAGCGTACCTCACCTGTCTGATAATCCAT
55 ACCATCTCTGATGCCGTAAGGGCGAGCGGTGACAAAATTACCGAAATGGCATCAATACG
AGATCCGTACATTGGGGAGTTAGCAAGCCCATCGACAGCCGAGCTTCGATTGTAGACG
AATTGAGCCGACTGCCCTCAAGATGGCAGGCATAGATTATACGGTGTTACACGCACATT
CTCATGTAGGTTTATATCAGCTTGGGATTTTGTAAATTTTTTCCACTTGTGGGCGTAC
ACCATGCACAAGCACCATTGATGCCAAGCTGTGTAGCAGTGCAAAATCATGAATCAG
60 CGTACTAAAATTGTGACGAGCGACCGCCTCATCACCACATAACCAAGGTTTGGC
ACGATGGGTGTTAATGTACGGGGCAGAATTACGAAACCAATGCACAGGTGTGAGTGCAGG
AGTGTCTGATAGGTGCTGACAGAATTCATGAATGCTCCAAAGAGTCAATGGCTGGTAAA
ATAAGAAATGGCGAATATATGGCGAGAGCGTCTGATGTTGGTCAAAATGTCCCAATTA
AACTATCAAGATACCATATACCATAGCAAAAGTTTGGGCAGATGCCAAGCGAATTTATC
65 AGCTTGATAAGGTTGGCATATGATAAAATCTACCATCATCGTCGCCAGTTTGGAGCATGT
GTAAGTAGTTACCATATTAACAGTCAAGAAATTCACACCGTCAATCAGCTGTGCTATG
CTTATGGGCACATAAACTTGACCAACACAGGATAAATTTA

SEQ. ID NO:57

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipo11 gene
from *Moraxella catarrhalis*

70 GGCATACTTTGGCCATGCTTTATTTTGGCATAACTGCTATAAGCCCATGCTACTTTTAA
TCATTTATCCATATGTCCAATAATGTGCTTTATGTAATTTAGGCACACTATTAACCTCGTG
CCACTGTTAACATTACGATATAAAATCTTAACAATGAATCAAAGCATCGTATTGGCTGTT
AAATGATAAGCTTATATTTATTTAAATTCAGACTAAATGATTGTAATATGGACATATCAA

GGTTGAAATCAAAAATTTTGGAGAGTTATGTACGATAATGATAAAAAATTGACCACCATC
 TAGGGGTGGTTGTATACGGTGTCTTATATTGCCATATGGTTGGTCAGTGGCTATATTTTA
 TGGGGCTGGATTGGTGTGACAGGATTACTCGTGCAGTACTTTGGCTGATCGCTTGGATG
 ATTGTGGGTACGATTGCTGATAGAATTCTGATACCGATTATTTTGACCGTCGTGGTTGGG
 5 TATTTTCTATCTTTTGAAGGCGATAATTGGTTATTTTTCACAAAAATCAT
 GATTTTGTAACTATCTAAAAATCAATTATGTTATATTATGTGATAAAAGATGGG
 CATGCTTAAGTTTGGATTGCAAAAATCCTAATATCATCACTGACCAAAGCTGTGATGAT
 ATCAAACTTTATCAAAGTTCTTAGGGTATTATCAAGATATCATACCAAATGAATACTTA
 10 CCCAACTTACTATAAAAAATCAATGATATGACTGTGATTTTATTATCATAGATACAAAA
 TCAAAACGCATGAGCCAAAGGTATGATGAATGAATACAAAATTTGCGACACATTATGACA
 ATCTAAATGTGCGCCAGAAACGCTGACATTGCGGTGATTGGTGGGATAGGGGTCAAGCCA
 GTGCGATTAAAGCTAAATTTTATGTGGCAATCGCTGACTTTATTTTATTTGTGCCAGTT
 GGAACAATTCGTGGTCTAATGTATTATTTAAGGAGATAA

SEQ. ID NO:58

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol0 gene
 from *Moraxella catarrhalis*

TCTGGTCTACATCCCAAATTTTACACAAGAACTAAAGACAGTGGAGCAGATGACG
 CTCAAAAGGCATCTTATAGTAATTTGACAGTTAATTTTCGTCAAGTGCTTGTACAAAA
 20 TACACCATCGTGAAGAAGTTTGTACCAATTTAAGCACAATCATTTTGGCAGCACTGTC
 AAGCAATGCTTCAGGCAAAATAGCTGCTGGTAAAGATACTTGGGTCAATGCAATCGCA
 TCACCCCTTCTGTGCTGGTGAAGCGATAAGTTTGGCATCTTGCCAAAATTGACCATGGT
 TTAGACCTTGGCGTGGCTTGTGGTATCGCTCCACATGTCGTAGAGTAGATATTCGGTCA
 25 TATCAAAAGGGCGATGGAAATGTATGGAATGGTCAATACTAGCCATTTGTAGACCTTGTG
 TCATCAGGCTTAGCCATGACTCATTAACCTGTGCTGACCAATAAATACAGACACAA
 ACGCAAGTAGTGCTTGAATGGAATGGCACTGGCTGCTCCCAATATCAGCGATACGCACCC
 AATTGGCTTGGCGTGGACGCTCAGGCTTGGGTGTCACAGGGTCTCGTGGTGTGACGGGGC
 GGATTTGACATGACGCTGACGCATAAATCTTGTCTTGTAGTGGTTCGGGAATTTATGTA
 30 AATAATCCGCTTTGAGTTCTTGTCTGGTTTTTAGGCTTTCAGGGGGTGGATAATCAGGCA
 TGGTTTCTTGGTAATCAAGCCCGCTTCCATGGGTGAAAATGAGGCAATCATCGAAAAAA
 TGACCTGTTTATGGTGTATGATTACCGTTTTTGTGCGGTGGTGGCAGATTTGACCCG
 CAATGACTTCTGAGCTGATAAACTGCGTCCATCACGTAAGCGGCGTACTTGATAGATGA
 CTGGTAGACGAATATCGCCACCTCGTAAAAAATAACCATGTAGGCTATGACAAGGTTTAT
 35 CAATCGTTAATGTGTTAGCACCAGCAAGCAGCGCTTGGGCA

SEQ. ID NO:59

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol2 gene
 from *Moraxella catarrhalis*

TAAAATGACCTTACAAAATAAAATTATATGTTCAAAAATCGCTTAAGTATTGAAAAAGC
 40 TATAAAATCTTATCTATAAAGCATAAAGATATTAAGCATAAAGACGAGAAAAAGAGC
 AAGCGTCAATGATGATATTATATATAAAATCTTATGAAATTTTCAATTTTTATCGATT
 GATTCAGCTTGGCTATCGGTGGTCAACTTTGGCTGCCAAGACATCGCCGGCTTTTTGAAA
 AATCATCACAAATGGCAACATGATGATGGTTGAAATCCACTTGACATATACCATGTTGCG
 ATGCTCACCATAGTTAATCGCAAGGCTTCCCAAGCCACCACCGCCACACCATGCCAT
 45 TGCAGAATAACCAATCAAAGACACCAAGGTCAATGTGACCGCATTAATCAAAATGGGCAG
 GCTTTCAGCAAAATAGTATTGTGACAACTTGCCAAATGCGTGTGACCCATAGATTGGC
 AGCTTCGGTCAGTCTGTGGTACTTCTAATAAAGCATTGGCACTCAAGCGTGCAAAAAA
 TGGAATTGCTGCCACACTCAAAGGGACGATGGCGGCTGTTGTGCCAAGGGTGTCTCCAC
 CAAAAATCGTGTGACTGGCATGAGAATAATGAGCAAAATAATAAAGGAACGGAGCGACC
 50 AATATTAAATAAACATCCAAAATTACAATACACTGCGATTTCAGGATACGCCCTTT
 ATCGGTTAAAAATGCCAAAAACCTATCGGTAGCCCAACCAAAACAGCGATGGCAGTGGC
 AGCAAGCCCCATATAGATGGTTTCCCAAGTGGATTGGGCAACCATCTCCACATTCTTGG
 GTGCAATTCCTGACGATTTTGTGACGATTTCATTCCACATAGCCGATAATCTCAATAT
 55 TGACCGATGGGTGGTTAAAAATCTATTGCTTGCATGACCGAGGTGCCTTACCGGATAA
 GCTCAGCAATGGTAAAGCCAAATTTATATCACCTGCATAA

SEQ. ID NO:60

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol7 gene
 from *Moraxella catarrhalis*

AGTAAACAATGGTAACAAATACAGCAGTGTGCGACAGTCTCAGTACGATGATTCTGAAT
 60 TTGAATATGCAGGATTTGGATACGATTTGTGGCATGTCTTGTGCGATAATTTAATTGTTA
 TGATTATATATGACCGCTATTGGTTTTATAATTATCAGCAAAATGATGGCCATGCCTGCTG
 ACCAAATACCGTTTTATAGTGTGGGGATGCCATCCTTTATAGTGTGGGGATGCTATCC
 TAACTTAGTGATGGCGGGCGGGTGTGGTTTGGGTAAAAAAGGTGCAACACCAG
 65 GTAAATGCTCTTTGGGCTGCAAGTCCGTGATGCCAAACAGGGCAATTTATCAGTGTGC
 CAAGGGCATATTGCGATATTTAGTTATCTGATTTCATCCGTGATTCTTTGTTTGGGAC
 TTATTTGGGTGGTTTTGATAAGAAAAACAAGGCTGGCATGATAAATTGCCAAAATCTG
 TTTGGTAAAAACGCATTCGCTGATGGGTGCGCCAGTTAAACAATAAAACCATCAACGCAA
 GCAGGGCGATGTGTTTGTGAGCAGTTGCGCGTAGATAAGCTAAACAAGCAGGCTATGAAAT
 70 TATTTTAACTTATCCACCCCATTTGTTGGTGAGATTGATATTATCGCCAGACAGCC
 TTTGGAGCAATCGACCGTTTGGTGCAGCCAGATTTTGTACGGTATTTGTTGAAGTGGC
 TAGCCGAACAAGTTCTGTGATGGTACAGCGCTTGTGAGTGTACCTCAAAAAAGCAGGC
 AAAAAATCTACCGAACAGCAGAACGATTTTAAATCAATTATCCCAATATATTGATGATGC
 75 ATACCGTTTTGATGTCATGGTTTTGATTGGTTGATGGATTGATTGAACATGAATGGAT
 AAAAAATGCGTTTTGATTGGTCAATGGTGTGAATTAATCAATCAAGCAATCCGTAG

CTTTACTATAAGATATATCCCAGTAATATGGAACATAGCA

SEQ. ID NO:61

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipof gene
from *Moraxella catarrhalis*

5 CGTTTAGCTTCATACGCAGACCTTGTGCACCTTCGGGCAACCGAAGCATCAGCCAGCAT
CACGCATCCGCACAAAACCCATCATGCCATCAATTCGCTGCTGATATGATATACCCCA
CCAAAGTAAACCGCTTAAATCGTGGATAACGCCTGCTGCTGAGGGTGAAGCTTCAGGCA
10 AAACCAAGGTAACTTATCCCCCACTTAAGTCCCATGTGAGAGACAATGGACTCACCTA
ATATAATACCAACTCGCCGATATGTAATCATCCAAATTGCCTGCGGTATATGCTCAT
CAATGATAGAACTTGTCTTTTCGTAATCAGGCTCAATGCCAGAAACACGATTCCAGTCA
CCTGACCTTCAGCGGTTAACATACCTTGTAGTTGAATATAAGGGGCAACTGCTTGCACCTT
CTGGATTTGCAATTTTGTATTTTCGCGCAAGTTCTTGCCAAATTTGTCAAATTTCTGTTG
15 AGGTAAGTGAAGCTTGAGGCACCATGCCAAGAATCGTGATTTAATTTACGGTCAAAGC
CATTATGACCGACAAAACCGTGATAAGCACTGCAACCCCAAGCGTAAGCCCAATGGTTG
AGATAAAGAAATAAAGGAAATAAGCCATTTTACGCTTAGCTTTGGTATATCTAAGCC
CAATAAATAACGCGCAAGGGACGAAACATAAGCTGTGTTCCAAACGACCCAAACCGTGCTAG
TTTAGCACTTTTTTGGACAAATACCAACATCACATAACAAATGAATCATCAGTTGGTT
20 TTGTTGCGCTTGTGATCTGTATGATAAGTTCTTGCTAAAACAGCTTTTATATGTCAGA
ATACAGAAAAGGTATATCTTATATTTTAACTTTAAATAGATCTGCTTTTTATACCGA
TGATTTGGCATGAAGTTTATCGGCTGTATGCTGGATATAAGTTTATCGGCTTGATATA
AATTTTAATTAATCATCAAATTTTAAAGGAATTTATCATT

SEQ. ID NO:62

25 Nucleotide sequence of DNA region (1000 bp) up-stream from the P6 gene from
Moraxella catarrhalis

TAAGGATACCAGATTTTGGCTTGTCAATCGTTGTGTTAATCATTGTAACGGTTTATAGTG
ATTGTCAATTAATAAGGGTAAAAAGTATTATCAAGTAATAATCTTCTTATATGTGAA
TATAATGACAAATTTATCACATTTTACAAGGATTTTTATCAAGATTAGGATATGTTCC
30 AGCTTAATTATTAGTGATGAGCGTGTGATTATTTGGCATCGTTAAATTTATGAGTGCTAA
AATTGCCAAATGATTAATTTTGTCAACATGATAGCCCTTTGGTAGGCTTATTTGGT
ATTGATGAGCAATAATAATATACCGAGTTAATGGATTAACTTAACATACGCCAAAACT
TAACAACGAAAAGTAGATGATTATGACAGATACAGTACAAAAGATACAGCACAGTCCCC
CAAAAAGTTTATCTAAAAGACTACACGCCCGCCAGTATATGCAAGTTAATAAGTGGATTT
35 GGATATCCGCTTGTGTTGATGATGATGCTGTGCTGTTGGTGCCAACTTAAATGACACGAGC
ACACGCGAGGCGAGCTTCGGCTTCTTGGGCGAGATTTAAAGCTTAAAGCATTCACCTAAA
TGGTCAGGAATTAGAGTCGACGGCGTATCATCTTGATAAGGAAGGCTTAACAATTTTAGA
TGCACCATGATGTCGAGTGATTGAGACATTGGTTGAGATTTACCACAAACCAACACAAC
ACTTGAAGGGCTATATCAAGCAGGAACAGGTGATGATAAGATGTTTGTGACACAATGCGA
40 ACCTGAGGTTTTTCGAAAATCACCTTTTTCCCTGACCGCCCTGATGTTTGTGACAGAATA
CACCACAGCCCTAGAGCAGCAACCAAGCATTTTAAACCTTGCTTGCCAATGGTAATTTGGT
TGAGTCAGGAGATGTTGATGAAATCGCCATTATACCATTTGGCATGATCTACCAAAAA
ACCCAGCTATCTATTCGCCGCTGTCTATTGCCAATCTAGAAG

45 SEQ. ID NO:63

Nucleotide sequence of DNA region (1000 bp) up-stream from the MshB gene
from *Haemophilus influenzae* (HiRd)

AAATCAAGCGCCTGTGCCTGCTGGTGATGGTTGTGGAGACGAATTATATTCTTGGTTTGA
ACCGCCAAAACAGGCACTTCAGTGAGCAAACTAAAGTTACACCGCTGAGCCGTTTTT
50 GTGCCAACAGATTTTGAATCACCAGATCGGAGAGAATGGTTAGATAGCATTGAGGTAA
ATCAATATGGATATCGGCATTGATCTTTTAGCAATATTGTTTGTGTTGGTTTTGTGCGA
TCATTATCGATGCAATGCTGGCGGTGGTGGATTAATCACCATTCCAGCGTTACTCATG
ACAGGTATGCCACCAAGCAATGGCGTTAGGCACCAACAAATTGCAAGCTATGGGCGGTGCA
55 TTATCCGCAAGCCTTTATTTCTTGCAGAAAAGAGCGGTCAATTTACGCGATATTGGTTT
ATTTTGATTTGGGTTTTCTTAGGTTCTGCCCTAGGTACATTATTAATCAATCAATTGAC
GTGGCGATTTTCAAAAAATGCTTCTTTTGGATTAGCCATTGGTCTATATTTTTTA
TTTACTCCTAAATAGGTGATGAAGATCGAAAACAACGATTAAAGTTATCTGTTATTTGGT
CTTTTAGTTAGCCCATTTTAGGTTTTATGATGGCTTCTTTGGGCCAGGGACTGGCTCA
60 ATCATGAGTTTAGCCTGTGTTACTTTGCTAGGATTTAATCTCCCGAAAGCGGCAGCACAT
GCAAAAGTGATGAATTCCTTCAACCTTGCTTCTTTGCACTTTCTTATTGGGCGGA
CAAAATCTTTGGAAAGTGGGTTTCGTGATGATGGCTGGGAGCATTTAGGTGCAAAATTA
GGTGCCAAAATGGTGATGACGAAAGGTAAACCTTGATTGACCGATGGTTGTTATCATG
TCTTTTATGATGACGGCTAAATGGTTTACGATCAGGGTGGTTTCATTTTAAATTCGGA
65 AAGCGCGCAAAAGTGGGTTAAATTAATTACATTTTATTA

SEQ. ID NO:64

Nucleotide sequence of DNA region (1000 bp) up-stream from the HtrB gene
from *Haemophilus influenzae* (HiRd)

70 TTGAAGTCCCAATTTACCACCAACATTCCTGCGGCAACATTGGCTAGGTAACAAGATT
CTTCGAAAGAACGTCATCTGTAATGTGGTTGCTAATACACTAATGACAGTGTACCCGG
CTCCGTCACATCAAACTTCTTTTGCAACGGTTGGCAATGATAAGGCTCTGATTTG
GGCGTAATAATGTCATGCTTTTTCAGAACGCGTCACCAAAAGTGGGTTAATTCATAT
CAGAAATTAATTTAAACCTTTCTTAATAATCTTCTCTGTATTACATTTACCTACAA

CGGCTTCAAATTCAGACATATTGGGTGTCAATAATGTAGCCCCACGATAACGTTCAAAT
 CAGTTCCTTTGGATCGATCAACACAGGCACATTCGCTTTGCGTGCAATTTGAATCATT
 TCTGAACATCTTTAAGCGTGCCCTTTGCCGTAATCAGAAAGAATCAAAGCACCGTAATTTT
 TCACCGCACTTTCTAATCTCGCTAATAAATCCTTGCAATCTACATTATTGAAATCTTCTT
 5 CAAAATCAAGCGGAGCAGCTGTTGATGACGAGATAAAATACGTAATTTAGTAATGGTTG
 GATGGGTTTCTAATGCAACAAAATTACAATCAATCTTTTGTTTTCTAATAAGTGGGAAA
 GTGCAGAACCTGTCTCATCTTGTCCAATCAATCCCATTAACCTGAACGGGTACATTGAGTG
 AAGCAATATTCATCGCCACATTTGCAGCACC GCCCGCGCTTCTTCATTTCTTGTACGC
 10 GAATCTAGGCACTGGTGCTTCTGGTGAAATACGGTTGGTTGCACCGAACCAATAACGAT
 CAAGCATCACATCGCTAATACAAGTACTTTTGTCTGCTTAAATCTGCTGAATATTGAG
 CCATTTTAAATCTCTCTATTTTGAATAACCAAATTTGTGGCGATTTTACCACAACCTCAA
 TTTACGATAAATACGCCCTTAACCTACGTGGAAAGAACA

SEQ. ID NO:65

15 Nucleotide sequence of DNA region (1000 bp) up-stream from the protein D
 gene from *Haemophilus influenzae* (HiRd)

AGCAATAATTATAGCTGGAATATTCTTTAAAGATGAAAGAGATCGTATAAGACAAAAAGA
 ATTTTATATTGGAGAATTATTAGCAATTATTGGTTTCGCTAATATTCTGTAATAAATAGTTC
 AAATATATGATGGAATACAGACTTTTTCTTGGGGCAATATTCTTTTACAGCTATTTT
 20 TATTCAATCTGTACAGAATTTAATTGTAAAAAAGTAGCCAAAAAGATAAATGCTGTTGT
 AATAAGTGCATCGACAGCAACAAATTTCAAGGATATTATTTTATGTTTACGCTTTAATAC
 TAAACAAATATATTATTACAAGATGTTGGCATTGGAATGTTGATAGGTTTAGTTGCGC
 TGGCTTTTATGGGATGCTAACAGGGATGTTGATGGCTTTTATATTGTTCAAAAACAGGG
 AATCACTGTTTTTAACTTTTGAATTTTGAATTTTCACTTTTCACTGCGATAATAGGTTA
 25 CTTAACATTAGATGAAAGAATAAATATCTATCAGGGAATTAGCGGTATTATTGTAATTAT
 TGGTTGTGATTGGCATTAAAAAGAAAAACAAGGAGTGTGATATATAAAGTAGATGAT
 GTTGGTGGAAATAGGTATAGTTAAATATCTGGTTCAATTGGTTTTATTAAGGGCGTTAGCA
 ATTCTCCATTTAAGTTTATGTTTGAATTAGATATTTGGGAAAAGATGGAAGAATAAAGC
 TGTAAATATGCTGAAACATATGAACTATACCAATCTCAAAATAAAATTAATCTGCTG
 30 GAAATGATTATAAATCTCTAATCTAATCTGATAGAGAGGATAATGACTATCAATCAGAAA
 GAATGATTAAAGCCATTAAAAATATTATTCATTGTATGACTAATAATCATCAACCTATT
 CAAGTGCTGAAACATCTTTAGAACTATTAAATATTACGGAATAAATTAATCTGTTA
 AAATAGGTAATGATCTTAACAATATATAAGGAGAATAAGT

35 SEQ. ID NO:66

Nucleotide sequence of DNA region (1000 bp) up-stream from the *Hin47* gene
 from *Haemophilus influenzae* (HiRd)

TAAATACTCCAAATAAATTCAGATAACGTGGTCTGTAAGACAAAAAATAAAAAAAT
 GTTCAATAAGAGGAGAGCAAAATATCTTGTTTAAAGGAAATCGGAGCAGTACAAAAACG
 40 GTCTTACAAGTAGCAAAATCTATAAATTTATGTTCTAATACGCGCAATTTCTAGTCAAT
 AAAAAGGTCAAAAAATGAGCTGGATTAAACCGAATTTTAGTAAAAGTCCTTCTCTCCA
 TCGAAAAAGCCAAATGTGCCAGAGGCGTATGGACAAAATGTACTGCTTGTGAACAAGTAC
 TTTATAGTGAAGAACTCAAACGTAATCTGTATGTTTCCCGAAATGTGGTCATCATATGC
 GTATTGATGCTCGTGAGCGTTTTATTAATTTATTGGACGAAGATTCAAGCCAAAGAAATG
 45 CGGCAGATTTAGAACCAAAAGATATTTTAAATTCAAAGATTAAAGAAATATAAAGATC
 GTATCAATGCGGCCCAAAAAGAACGGGCGAGAAAGATGCGCTAATTACTATGACAGGTA
 CACTTTATAATATGCCAATCGTTGTGGCTGCATCGAATTTGCTTTTATGGGCGGTTCAA
 TGGGTTCTGTAGTTGGTGCAAAATTTGTTAAAGCGGCTGAAAAGCGATGGAATGAATT
 GTCCATTTGTGTGTTCTCTGCGAGTGGTGGTCTGCTATGACGGAAGCATTATCTCTT
 50 TAATGCAAAATGGCAAAACTAGTGCCGTAATGCTCAAAATCGGTGAAAAGGGGTGCCAT
 TTATTTCAATTAACGGATCCGACTTTAGGCGGCGTATCAGCCAGTTTTCGATGTTAG
 GGGATTTAAATATTGCCGAGCCAAAAGCCTTAATGGTTTTGCAGGGCCACGCGTTATTG
 AACAACTGTGCGTGAAAATTTGCCAGAAAGTTTCCAACGTAGTGAGTTTCTACTTGAGA
 55 AAGGGCAATTGATATGATCGTGAACGTTTCAGAAATGCGT

SEQ. ID NO:67

Nucleotide sequence of DNA region (1000 bp) up-stream from the P5 gene from
Haemophilus influenzae (HiRd)

TCACCTAATTCAAGCGCATCAATGTTTTCTAAAACATCAACAGAATTGACCGCACTTGTA
 60 TCTAAAATTTCCCATTTTATTAAGACTGCGCGTAATGCCAAAACATGATTAGAGGTTTTA
 CCATATTGCAATGAGCCTTGCCCGAGGCATCGGTGTTAATCATTCCACCTAAAGTCGCT
 CGATTGCTGGTGGACAGTTCTGGGGCAAGAACAACCATGTGGTTTTAAAAATTGATTA
 AGTTGATCTTTTACTACGCTGCTTGACTCGAACCAACGTTCTTTTACATTGAGTTCT
 AAGATGGCTGTCATATGACGAGAAAGATCCACTATTATATTGTTATTGATGGATTGCCCA
 65 TTTGTGCCAGTGCCCTCCACCGCGAGGCGTAAAGCTGATTGATTGATATTCAAGTAAATTT
 GCCAATTTTGTATCCGCACTATATCAGCAACCGTTTTCCGAAAAAGAATTGCTTGTGGA
 AGTTGTTGGTAAACGCTGTTATCCGTAGCCAGACTTAATCTATCTGCATAGTTTGTGCGA
 ATATCCCCCTCAAAATGTTGGCATTGAAGATCATCAAGATAATCAAGTACATATTGTTCA
 ACTTGAGGAATGCGATTTAGATTTGGCAACATAGTATTGACCCATTTAAACATATCAGA
 70 TGGAGGCTTTGATAATATCCTAAGGCTAGAATAATGTCGATTAGGAAAGAGAGAGGAGAA
 AGTAAAAAGTCTGTTTAAAGAAAGTGTATTTTGGATAAAAACTAAACAAAAAATCAAAA
 GAATTTGATCTTTTCAATTTTATAGGATAATAAGCGCACTTTTGACGTTCTTTGGGG
 TAAACATAAGCAAAGGAATTGAATTTGTCAAAAGGTAATAAAGTAGGGCAAAATCAAAAC
 75 CCTAGTTAAGTGACTGTTTATAATGTAGCTTTAATTAAAGTTTCAGTATAAACAAAGGACA
 CTTTTTATTACTATTTCGATCACTAAATAGAGGACATCAAAA

SEQ. ID NO:68

Nucleotide sequence of DNA region (1000 bp) up-stream from the D15 gene from *Haemophilus influenzae* (HiRd)

5 TCGATTGTATCCTATATAAATTATAGACGTAAAAATCATTAAATAATGCAAAACACCGTT
AAGCTTAATAACAGTGCTGCCCAATTTCGATAACAGATGCTTGCACCCGCTCAGAAACA
GGTTTTCCCTTTAACAGCTTCCATTGTTAAAAAACTAAATGACCGCCATCTAATACTGGT
AATGGAATAAATTCATAATCCCTAAATTTACACTAATCAATGCCATAAACTTAAAAAA
10 TACACCAATCCAATATTGCTGATGCGCCAGCACCTTTTGCAATAGAAATGGCCCACTT
AAATTATTAAATGACAAATCGCCAGTAAGTAATTTCCCTAATATTTTCAAGGTTAAAGG
GAAAGCTGTCTGTTTTTCAATGCCTTTTGTAAAGATTCAAGAATACCATATTTTAAAT
TCAGTACGGTATTCATCCGCTAATTTTGTTAAGGCTGGGCTAACCCCAACAAACCATTTG
CCATTTTGATTACGCACTGGAGTTAGGACTTTGTCAAATGTTTCTCCATTACGTTCAACT
TTAATAGAAAAGATTGCGCTTGTTCGACCTGTTTTATAAAATCTTGCCAAGGAAGTCGG
15 GTTAAATTTTCTTTTAAATTTTATCACCAGTTTGTAAACCAGCTTCTCAGCGGGAGAA
TTTTGAACAACCTTAGAAAGCACCATTTCATTTTAGGACGCATAGGCATAATCCCTAAT
GCCTCAAAGCACTTTCTTTTTCAGGATCGAATGTCCAATTTGTAAAGTTTAAAGTCCGT
TGTGTTTCAATATTAGAATTGAAAGGAGAAAGGCTAATCTCAACATTAGGCTCCCCATT
TTTGTGGCAAGTAGCATATTGATGGTTTCCCAATCTTGAGTTTCTTCGCCATCAATTGTA
20 AGAATTTGCGTATTGGGTTCATGTGGGCTTGTGCTGCGATTGAGTTTGGTGTATTGAT
TCAATCACTGGTTTAAACCGTTGGCATTCATAAAGGTAAT

SEQ. ID NO:69

Nucleotide sequence of DNA region (1000 bp) up-stream from the Omp26 gene from *Haemophilus influenzae* (HiRd)

25 TTTGATAAATACCTTAATTAATGATGGGTTTAATATTTCTCTGCCCAATTAAATTAG
TCAGAGAACGTGTGTTTGAAGTTCTGATGAAGAAAAAGTTCAATTTATTAGAAGAACC
TCCAATACTAATTTGGAACCTGTTGACATCATCTTTTCATATTTTAAATTGGTTTGGC
ATAAGAGAATACCAATGGCCCAATAGGAGATTGCCATTGGAATCCGACACCTGTAGAGGC
30 CGGAATACGGCTTGATTGCTTATGCCATAATCGGGTAAGCTTTTAAATACATTGTTATCAACC
ACTCTTATCCGATTTCCTACTTAGTATTCCAAACACTTGCCGCATCAACAAATAGGGAGGT
TCGGACTGTATTTGGCTTTTATCACTCACAACCGGTGTTGGTACAATAAGTTCTGCACT
CGCAGTTGTGATTGCTATACCAATCAGATCAGAACTTATCTCTTAAAGTACCATT
ACCATTACCATGTTCTGCATAAATTGCGTTAGGTCCAATACTACCATAAGCAAACACG
35 TAATGAACCGATGCCACCCGCTGTATAAGTTTGATAGAAGCGTAAACGCTTGTTCCTAAA
ACCATTGCAATATCCTGCAGATGCTTTTGCAGATACAACCCAGAGGTGATCTCTGTCTAA
TGGGTAGAAACCTGTACGCTGCACTTAGTTGTAGTATTGTTATCAGAACCTGGAAT
AGTAACTCGTCCCAAGACTTGCTTTAACCCCTTGAAGTTGGGAAATAGCCTCTATTAG
40 GCTGTTTGCATATCCTGCAGATGCTTTTGCAGATACAACCCAGAGGTGATCTCTGTCTAA
AAATTTCAATTGATTGAATATATAAATTACGGTTATATTCTAGAGCAAAGTTACTAATTT
ATTATAGGTATGGCTTAATCTACATAATAGGAGTTATTTTCATTTACAGGGAAACCTAA
AGTAACATTACTTCCATAAGTCGTACGCTTATAGTTAGAGG

SEQ. ID NO:70

Nucleotide sequence of DNA region (1000 bp) up-stream from the P6 gene from *Haemophilus influenzae* (HiRd)

45 TTAGATTTCCTCTAAATGAGTTTTTTATTTAGTTAAGTATGGAGACCAAGCTGGAAATTT
AAGTTGACCATCACTTCTGGAAGGCTCGCCTTAAAGCGACCATCTGCGGAAACCAATTG
TAGCACCTTTCTTAAGCCCTGTGTAGAACTATAAATAATCATAATTCATTGGAGAGAG
50 GCTTGGGCTTTGCGCTAGAAAAGATGTACTAAGTACCTCTGAAACGCCGTTGTGAGATC
TTGTTTAACTACATTATTGTTACCATTAACTCATCACAAGTGTTTTCCATCTGCACTAAT
TTGTGCGCTACCGGACCACTGCTGTTGCACTACCAACCGCTTGCATCCATTGCGATA
AAGTTGTGGCGAACCACTTCTATCGGATGTAAATAAATTTGAATTTCCGTCTGGCGACCA
55 CGCTGGTTCAGTATTATTACCGCACCACTCGTCAATTGAGTAGGTGTACCGCAATTGTC
TCCCATACGTAATATTTCAGAACACCATCACGAGAAGCAAAAGCTAAACGAGAACC
ATCTGGCGAAAAGGCTGGTGCCTTATGCCCTTGAAAAGATGCCACTACTTTACGTGC
GCCAGAAATTTAAATCCTGTACAAACAAGTTGTGATTTTTTATTTTCAAACGATACATAAGC
CAAACGCTGGCCGCTCTGGAGACCAAGCTGGAGACATAATTGGTTGGGCACTACGATTGAC
60 GATAAATTGATTATAGCCATCATAATCTGCTACACGAACCTTCAAGGTTGCGAACCGCC
ATTTTTTTGCACAACATAAGCGATACGAGTTCTAAAGGCACCAAGGATCGCAGTTAATTT
TTCAAAACCTTCATCGCTCACAGTATGCGCGCCATAGCGTAACCAATTTATTTGTTACTGT
ATAGCTATTTTGCATTAATACAGTCCCTGGCGTACCTGATGCACCAACCGTATCAATTAA
TTGATAAGTAATACTATAACCATTACCCGATGGAACCACTT

SEQ. ID NO:71

Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpA gene from *Haemophilus influenzae* (non-typeable)

70 GGCGATAACCGAGTTTTTGGGGTATTTAGTGCCAAAGAAGACCCACAAAACCCAAATTA
TCCAGAGAAACCTTAATTGATGGCAAGCTAATACTTTTTTAAAGAAGTGTGCAAAAACCC
AATACAACAGCCGATACAAACAACCAATAAAACAACCAATGCAATAACCGATGAAAAAAC
TTTAAGACCGGAAGATATACTAAGTTTTTGGTGAAGCTGATTATCTTTTAATTGACAATCAG
CCTGTTCCGCTTTTACCTGAAAAAAATACTGATGTTTCATAAGTAGTAGGCATCATACT
GTAGGAAATAAACGCTATAAAGTGGAAGCATGTTGCAAGAATCTAAGCTATGTAAATTT

GGTATGTATTATGAAGACCCACTTAAAGAAGAAGAAAAAGAAAAAGAAAAAGAC
 CAAGAAAAAGAAAAAGAAAAACAAACGACGACAACATCTATCGAGACTTATTATCAA
 TTCTTTAGGTGCACCGTACTGCCAAGGCCGACATACCTGCAACGGGAAACGTGAAATAT
 5 CGCGGTAATTGGTTTGGTTATATTGGTGATGACACGACATCTTACTCCACTACTGGAGAT
 AAAAATGCTCTCGCCGAGTTTGATGTAAATTTTGGCGATAAAAAGCTAACAGCGGAATTA
 AACGACACGATAATGGAAATACCGTATTTAAATTTACTGCAGACCTTCAAAGTGGTAAG
 AATGACTTCACTGGTACAGCAACCGCAACAAATTTTGTAAATAGATGGTAACAATAGTCAA
 ACTGGAAATACCCAAATTAATATTAACCTGAAGTAAATGGGGCACTTTTATGGACCTAAG
 10 GCTACAGAATTAGGCGGTTATTTACCTATAACGGAAATCTACAGCTAAAAATTCCTCA
 ACCGTACCTTCACCACCAATTCACCAATGCAAGAGCTGCAGTTGTGTTTGGAGCTAAA
 AAACAACAAGTAGAACAACCAAGTAATGGAATACTAAAA

SEQ. ID NO:72

Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpB gene
 from *Haemophilus influenzae* (HiRd)

15 TAGAATTATATTCTTATACAAAATTGATAATTGTTTCGCTTATCATTTTTTTTTTTGTAAT
 AATGTCAACTTATAATTTTTTAAGTTCATGGATAAAATATGAAAATGGCGTAAAAACAC
 TTTTCTCTTATCATTAAATAGGCTTATCATTACGAATGTAGCTTGGGCAGAAGTTGCAC
 20 GTCCTAAAAATGATACAAATGACAAATACGATTCAAAGTGGGAATTAACCACTCCTCTT
 TTTCTCTATGCTAAGAAAGAAATACCAATAGGCATATTATTTCTCTTTCCAAAAGCC
 AATTAGCGCACCATCCAAGGCTTGTTCGCTGGGTAAATCCTGCTTTATATCAAAAATA
 ACACTCAGGCAGTTCAACTGTTATTACCACTATATAACAATTTCTCAACAAGATAATT
 TCTTACTAATTGGGCAAGGCTATTGAAGCTCGTGAACAAGGTGATTAACTCAATCTA
 25 TTGCTTATTATCGTGAATTTATTCGCTCGAGACGATCTTACTACCTTACGTTATTAAT
 TAGCTCAAGCTCTATTTTTAACTATGAAAATGAAGCTGCCAAAATTCATTTGAAAAAT
 TACGTACAGAGGTAGATGATGAAAAATTTTAGGTGTTATTGATCAGTATCTTTTAACAC
 TAAATCAGCGGAATCAATGGATATGGCAAGTAGGATTAATTTTTAAATGATGATAATT
 TGAATAACGCTCCAAAAGTGGCACAAAATTTGGTAGTTGGACCGCTTGGGAAAAAGAAA
 30 GTGGGCGAGGGGTAGGGTATTCTTTATCAGTAGAAAAAAATGGCCATGGGCAGATCAT
 TTTTATGATAAACTATGTTTAAATGGGAATGGAAAATATTATTGGGATAATAAAAAATACA
 ATGAGGCTACTGTGCGTATAGGTGGTGTAGGCTATCAAACTGCCTCAGTTGAAGTCT
 CGTTGTTTCTTTTCAAGAAAAACGCTGGTATGCAGGCGGT

SEQ. ID NO:73

Nucleotide sequence of DNA region (1000 bp) up-stream from the HifA (pilin)
 gene from *Haemophilus influenzae* (LKP serotype 1 genome)

35 TAATAAATTGCTCCATAAAGAGGTTTGTGCCTTATAAATAAGGCAATAAAGATTAATATA
 AACCGTTTATTAATAAGCCAAAGGCTTAATAAACAGCAAACTTTGTTTCCCAAAAAAAG
 TAAAAAAGCTCTCCATTATATATATATATATATAATTAAGGCCCTTTTGAATAATTT
 40 CATATTTTTTTGAATTAATTCGCTGTAGGTTGGGTTTTTGGCCACATGGAGACATATAAA
 AAAGATTTGTAGGGTGGGCGTAAGCCACGCGGAACATCATCAACAACCTGTAATGTTGT
 ATTAGGCACGCTGGGCTTATGCTTCGCTACGGGGAATGAATAAGGATAAATATGGGCT
 TAGCCAGTTTATGGATTAATATGTTGAAATGGGGAACAATGTTTAAAAAACACT
 45 TTTATTTTTTACCGCACTATTTTTTGGCGCACTTTGTGCATTTTACGCAATGCAGATGT
 GATTATCACTGGCAGCAGAGTGATTATCCCGCTGGGCAAAAAATGTTATCGTGAAGTT
 AGAAAAAATGATGATTGCGCAGCATTGGTGCAAGCCTGGATTGATAATGGCAATCCAAA
 TGCCGATCCAAAATACACCAAAACCCCTTTTGTGATTACCCCGCCTGTTGCTCGAGTGGA
 AGCGAAATCAGGGCAAGTTTGGCGATTACGTTACAGGCGAGCGAGCCTTTACCTGATGA
 50 TCGCGAAAGCCTCTTTTATTTAATTTGTTAGATATTCGCGCGAAACCTGATGCGGCATT
 TCTGGCAAAACACGGCAGCTTTATGCAAATGCCATTGCTCAGCTTTGAAGTTGTTTTA
 TCGCCCTGCGAACTCTCGATGGATTCTCGTGATGCAATGAAAAAGTAGTGTTTAAAGC
 CACACCTGAAGGGGTGTTGGTGGATAATCAAAACCCCTTATTATGAACTACATTGGTTT
 GTTACATCAAAATAAACCTGCGAAAAATGTCAAATGGTTG

SEQ. ID NO:73

Nucleotide sequence of DNA region (1000 bp) up-stream from the HifE (tip
 pilin) gene from *Haemophilus influenzae* (LKP serotype 1 genome)

55 TAGTAGATTTCCGACAGGGCAAAAATACAATGGTGTTATTTAACCTCACTTTGCCAAATG
 GCGAGCCAGTGCCAAATGGCATCCACCGCACAAGATAGCGAAGGGGCATTGTGGGCGATG
 60 TGGTGCAAGGTGGTGTGCTTTTCGCTAATAAACTTACCCAGCCAAAAGGCGAGTTAATCG
 TCAAATGGGTGAGCGAGAAAGCGAACAATGCCGTTTCCAATATCAAGTTGATTGGATA
 ACGCACAAATACAAAGTCACGATATTCAATGCAAAACCGCAAAATAAATTAATGAAGAGG
 ATTTATGCAAAAAACACCCAAAAAATTAACCGCGCTTTCCATCAAAAAATCCACTGCTAC
 65 TTGTAGTGGAGCAAAATATAGTGGAGCAAAATATAGTGGCTCAAAATGCTTTAGGTTTCA
 TCGCTGCGCTCTGCTTGCTTGCTGGCTCTGCTTGATTGCATTGTGGCACTGCCTGCTTA
 TGCTTAGCATGGCAGAGTGACCTTTCAAGGGGAGATTTAAGTGATGGCACTTGTAATAAT
 TGAACACAGACAGCCAAAATCGCACGGTTACCTGCCAACAGTGGGAAAAGCTAATTTAAG
 70 CCACGCAGGGCAAAACCGCGCCCTGTGCTTTTCCATCACGTTAAAGAATGCAATGC
 AGATGATGCTATGAAGCTAATCTGCTATTTAAAGGGGAGACAACACAACAGGGCAATC
 TTATCTTTCCAATAAGGCAAGGCAAGGCAAGCCCAACGTTGGGCATTCAAATTTGTCAA
 AGCCGATGGCATAGGCAGCCTATCAAGGTGGACGGCACCAGCAACAGCGCAAAAAGC
 CCCCACACAGGTAAGCGCAAAACGGCACAGTTATTAACCCCGTTTTGGCTACTTTGG
 75 CTCGTTATTACGCCACAGGTGAAGCCACCGCAGGCGAGCTTGAAGCCACTGCAACTTTTG
 AAGTGCAGTATAACTAAAAATTTATTATCCAGTGAAAAAA

SEQ. ID NO:75

Nucleotide sequence of DNA region (1000 bp) up-stream from the P2 gene from *Haemophilus influenzae* (HiRd)

```

1  TTATCCGCTA ACATTTTCATC AGTAATTCCA TGAACCTTAA TCGCATCAGG
5  51 ATCANCAGGGG CGATCTGGCT TAATATAAAT ATGAYAATTA TTACCTGTGT
101 AACGACGATT TATTAATTCA ACTGCACCAA TTTCAATAAT GCAGTGTCCCT
151 TCATAATGGG CGCCAAGCTG ATTCATACCT GTAGTTTCAG TATCTAATAC
201 AATTTGGCGA TTGGGATTAA TCATTGTGTC AACCTATCTC TTTCCATTAA
251 AATACTTGCC ATTCTACACA ACAACCTTTT TGTATGCGCK AAACAGATTG
10  301 AAATTTTAC TGATGGATCT TGCTTAGGTA ATCCAGGGGC GGGCGGAATT
351 GGTGCCGTAT TGCGTTATAA ACAACATGAA AAAACACTCT CCAAAGGCTA
401 TTTCCAAACC ACCAATAATC GAATGGAATT ACGCGCTGTC ATTGAAGCAT
451 TAAATACATT AAAAGAACCT TGCTTGATCA CGCTTTATAG TGATAGCCAA
501 TATATGAAAA ATGGCATAAC CAAATGGATC TTTAACTGGA AAAAAAATAA
15  551 TTGGAAAGCA AGTTCTGGAA AGCCTGTAAA AAACCAAGAT TTATGGATAG
601 CCTTAGATGA ATCCATCCAA CGTCATAAAA TTAATTGGCA ATGGGTAAAA
651 GGCCATGCTG GACACAGAGA AAATGAAATT TGGCATGAAT TAGCAAAAAA
701 AGGGGCAGAA AATCCGACAT TGGAAGATAT GGGGTACATA GAAGAATAAT
751 ACAACTGATA TACCGTCATA TTTTTCGATA CCTAAAAATA TTTAATACTT
20  801 AAACCTAAAA CAGAATAAAA AATAATCAAA TTCATTTAAA AAATGTGATC
851 TCGATCAGAT TTCAAGAAAA TTTAAATTTT GGAGTATTGA CATCAAAAAA
901 TTTTTCGTGA AAGATGCAGC TCGTCCGTTT TGGCGATTGG ACAATTCAT
951 TGGAGAAAAA TTCAATCATA GATAGTAAAC AACCATAAGG AATACAAATT
1001 A
25

```

SEQ. ID NO:76

Nucleotide sequence of DNA coding region (partial) of the *Moraxella Catarrhalis* HtrB gene

```

1  TCAGTGCTTG GTTTTTTAAG ATATGTACCG CTGTCAGTCC TGCATGGATT
30  51 GGGCGCGTGT GCGTCTTATA TTTCTATCA TTGCAGGCTT AGTATTTATC
101 GCAGCATCCA AGCCAAATTTA ATCTTGTTTC ACCCCAAGAT GCCAGACGCA
151 CAGCGGCAAA AACTCGCCAA ACAATCCTA AAAAATCAGC TCATCAGTGC
201 AGTCGACAGT CTTAAAACCTT GGGCAATGCC ACCAAAATGG TCTATCGCAC
251 AAATTAAAC GGTTCATCAT GAAGATATCC TAATCAAAGC ACTTGCCAAT
35  301 CCAAGTGTTG TGCTTGCCAT TGTGCTCAT ATCGGCACTT GGGAGATGAT
351 GAATGCTTGG CTCATACCT TTGGCTCCCC TACTATCATG TATAAGCCCA
401 TCAAAAATGC GCGGTTAGAT CGCTTTGTTT TACAGGGGCG TGAAAGACTA
451 AATGCCAGCC TTGTACCCAC AGATGCTAGT GGTGTTAAGG CAATTTTTAA
501 AACACTCAAA GCAGGTGGAT TTAGTATCAT ACTGCCCGAC CATGTACCTG
40  551 ATCCATCAGG TGGTGAGATT GCTCCTTTTT TTGGTATTAA AACCTTAACC
601 AGTACGCTGG CGTCAAAGCT TGCTGCAAAA ACTGGTTGTG CTCTTGTTGG
651 CTTAAGCTGT ATTCGGCGTG AAGATGGCGA TGGTTTTGAA ATTTTGTGTT
701 ATGAATTAAT TGATGAACAA CTTTATTCAA AAAATACCAA AATTGCAACC
751 ACTGCTTTAA ATGGTGCGAT GGAACAAATG ATTTATCCAC ATTTTGTGCA
45  801 TTATATGTGG AGCTATCGTC GGTTCAGCA TACACCACTA TTAATAATC
851 CTTATTTACT TAATGAAAT GAGCTAAAAA AAATAGCCAT AAAGCTTCAA
901 GCCATGTCAA AGGATAGTTA TGAG

```

Protein Seq: 25% identity and 35% similarity with HtrB from *E. coli*

```

50  1 SVLGFLRYVF LSVLHGLAAC ASYISYHCR LSIYRSIQANL ILVHPKMPDA
51 QRQKLAKQIL KNQLISAVDS LKTWAMPPKW SIAQIKTVHH EDILIKALAN
101 PSGMLAIVPH IGTWEMMNAW LNTFGSPTIM YKPIKNAVD REV LQGRERL
151 NASLVPTDAS GVKAIFFKTLK AGGFSIILPD HVPDPSSGGEI APFFGIKTLT
201 STLASKLAAK TGCALVGLSC IRREDGDGFE IFCYELNDEQ LYSKNTKIAT
55  251 TALNGAMEQM IYPHFLHYMW SYRRFKHTPL LNNPYLLNEN ELKKIAIKLQ
301 AMSKDSYE

```

SEQ. ID NO:77

Nucleotide sequence of DNA coding region of the *Neisseria* (meningococcus B) HtrB gene

```

1  ATGTTTCGTT TACAATTCGG GCTGTTTCCC CCTTTGCGAA CCGCCATGCA
51 CATCCTGTTG ACCGCCCTGC TCAAATGCCT CTCCTGTGTC CCACTTTCCT
101 GTCTGCACAC GCTGGGAAAC CGGCTCGGAC ATCTGGCGTT TTACCTTTTA
65  151 AAGGAAGACC GCGCGCGCAT CGTCGCCAAT ATGCGTCAGG CAGGCATGAA
201 TCCCGACCCC AAACAGTCA AAGCCGTTTT TGCGGAAACG GCAAAGGGCG
251 GTTTGGAATC TGCCCCCGCG TTTTTCAGAA AACCGGAAGA CATAGAAACA
301 ATGTTCAAAG CGGTACACGG CTGGGAACAT GTGCAGCAGG CTTTGGACAA
351 ACACGAAGGG CTGCTATTCA TCACGCCGCA CATCGGCAGC TACGATTGTT
70  401 GCGGACGCTA CATCAGCCAG CAGCTTCGTT TCCCGCTGAC CGCCATGTAC
451 AAACCGCCGA AAATCAAAGC GATAGACAAA ATCATGCAGG CGGGCAGGGT
501 TCGCGGCAAA GGAATAACCG CGCCTACCAG CATACAAGGG GTCAAACAAA
551 TCATCAAAGC CCTGCGTTCG GCGGAAGCAA CCATCGTCCT GCCCGACCAC

```

601 GTCCCTCCC CTCAAGAAGG CGGGGAAGGC GTATGGGTGG ATTTCTTCGG
 651 CAAACCTGCG TATACCATGA CGCTGGCGGC AAAATTGGCA CACGTCAAAG
 701 GCGTGAAGAC CCTGTTTTTC TGCTGCGAAC GCCTGCCTGG CGGACAAGGT
 751 TTCGATTGCG ACATCCGCCC CGTCCAAGGG GAATTGAACG GCGACAAAGC
 801 CCAATGATGCC GCCGTGTTCA ACCGCAATGC CGAATATTGG ATACGCCGTT
 851 TTCCGACGCA GTATCTGTTT ATGTACAACC GCTACAAAAT GCCG

Protein Sequence - 30% identity and 38% similarity with Htrb *E. coli*

1 MFRLLQFGLFP PLRTAMHILL TALLKCLSLP PLSCLHTLGN RLGLHAFYLL
 51 KEDRARIVAN MRQAGMNPDP KTVKAVFAET AKGGLELAPA FFRKPEDLET
 101 MFKAHVHWEH VQALDKHEG LLFITPHIGS YDLGGYISQ QLFFPLTAMY
 151 KPPKIKAIK IMQAGRVRGK GKTAPTSIQG VKQIKALRS GEATVLPDH
 201 VPSPEEGEG VWVDFGKPA YMTLAAKLA HVKGVKTLFF CCERLPGGQG
 251 FDLHIRPVQG ELNGDKAHD AVFNRNAEYW IRRFPTQYLF MYNRYKMP

SEQ. ID NO:78

Nucleotide sequence of DNA coding region of the *Haemophilus influenzae* (non-typeable) HtrB gene

1 ATGAAAAACG AAAAAGCTCC TCAATTTCAA CCGCACTTTT TAGCCCCAAA
 51 ATACTGGCTT TTTTGGCTAG GCGTGGCAAT TTGGCGAAGT ATTTTATGTC
 101 TTCCCTATCC TATTTTGGCG CATATTGGTC ATGGTTTCGG TTGGCTGTTT
 151 TCACATTAA AAGTGGGTAA ACCTCGAGCT GCCATTGCAC GCCGTAATCT
 201 TGAACCTTGT TTCCCTGATA TGCTGAAAA CGAACGTGAG ACGATTTTGC
 251 AAGAAAAATCT TCGTTTCAGTA GGCATGGCAA TTATCGAAAC TTGCATGGCT
 301 TGGTTTTGGT CGGATTCACG TATCAAAAAA TGGTCGAAAG TTGAAGGCTT
 351 ACATTATCTA AAAGAAAAATC AAAAAGATGG AATTGTTCTC GTCGGTGTTC
 401 ATTTCTTAAC GCTAGAACTT GGCGCACGCA TCATTGGTTT ACATCATCCT
 451 GGCATTGGTG TTTATCGTCC AAATGATAAT CCTTTGCTTG ATTGGCTACA
 501 AACACAAGGC CGTTTACGCT CCAATAAAGA TATGCTTGAT CGTAAAGATT
 551 TACGCGGAAT GATCAAAGCT TTACGCCACG AAGAAACCAT TTGGTATGCG
 601 CCTGATCAGC ATTACGGCAG AAAAAATGCC GTTTTGTGTC CTTTTTTTGC
 651 AGTACCTGAC ACTTGCCTA CTACTGGTAG TTATTATTTA TTGAAATCCT
 701 CGCAAAACAG CAAAGTGATT CCATTTCGCG CATTACGCAA TAAAGATGGT
 751 TCAGGCTATA CCGTGAGTAT TTCAGCGCCT GTTGATTTTA CGGATTTACA
 801 AGATGAAACG GCGATTGCTG CGCGAATGAA TCAAATCGTA GAAAGGAAA
 851 TCATGAAGGG CATATCACA TATATGTGGC TACATCGCCG TTTTAAACAA
 901 CGTCCAGATG AAAATACGCC TAGTTTATAC GATTAA

Protein Sequence - 57% identity and 66% similarity with HtrB *E. coli*

1 MKNEKLPGFQ PHFLAPKYWL FWLGVAIWS ILCLPYPILR HIGHGFGWLF
 51 SHLVGKRRRA AIARRNLELC FPDMPENERE TILQENLRV GMALLETGMA
 101 WFWSDSRKIK WSKVEGLHYL KENQKDGIVL VGVHFLTLEL GARIIGLHHP
 151 GIGVYRPNDN PLLDWLQIQG RLRSNKDMLD RKDLRGMKA LRHEETIWAY
 201 PDHGYGRKNA VVPPFAVDP TCTTTGSSYL LKSSONSKVI PFAPLRNKDG
 251 SGYTVSISAP VDFDLDQDET AIAARMNQIV EKEIMKGISQ YMWLHRRFKT
 301 RPDENTPSLY D*

SEQ. ID NO:79

Nucleotide sequence of DNA coding region of the *Haemophilus influenzae* (non-typeable) MsbB gene

1 ATGTCGGATA ATCAACAAAA TTTACGTTTG ACGGCGAGAG TGGGCTATGA
 51 AGCGCACTTT TCATGGTCGT ATTTAAAGCC TCAATATTGG GGGATTTGGC
 101 TTGGTATTTT CTTTTTATTG TTGTTAGCAT TTGTGCCTTT TCGTCTGCGC
 151 GATAAATTGA CGGGAATTT AGGTATTGTT ATTGGGCATA AAGCAAAGAA
 201 ACAGCGTACG CGTGCACAAA CTAACCTTGA ATATTGTTTC CCTCATTGGA
 251 CTGAACAACA ACGTGAGCAA GTGATTGATA AAATGTTTGC GGTGTGCGCT
 301 CAGGTTATGT TTGGTATTGG TGAGATTGCC ATCCGTTCAA AGAAACATTT
 351 GCAAAAACGC AGCGAATTTA TCGGTCTTGA ACATATCGAA CAGGCAAAAG
 401 CTGAAGGAAA GAATATTATT CTTATGGTGC CACATGGCTG GCGGATTGAT
 451 GCGTCTGGCA TTATTTTGCA CACTCAAGGC ATGCCAATGA CTTCTATGTA
 501 TAATCCACAC CGTAATCCAT TGGTGGATTG GCTTTGGACG ATTACACGCC
 551 AACGTTTTCG CGGAAAAATG CATGCACGCC AAAATGGTAT TAAACCTTTT
 601 TTAAGTCATG TTCGTAAGG CGAAATGGGT TATTACTTAC CCGATGAAGA
 651 TTTTGGGGCG GAACAAAGCG TATTGTGTTA TTTCTTTGGG ACTTATAAAG
 701 CGACATTACC AGGGTTAAAT AAAATGGCAA AACTTTCTAA AGCCGTTGTT
 751 ATTCCTGTTA TTCTCTGTTA TAACGCTGAA ACGGGCAAAT ATGAAATGGA
 801 AATTCATCCT GCAATGAATT TAAGTGATGA TCCTGAACAA TCAGCCCGAG
 851 CAATGAACGA AGAAATAGAA TCTTTTGTTA CGCCAGCGCC AGAGCAATAT
 901 GTTTGGATT TCGAATTATT GCGTACAAGG AAAGATGGCG AAGATCTTTA
 951 TGATTAA

Protein Sequence - 45% identity and 56% similarity with MsbB *E. coli*

1 MSDNQNLRL TARVGYEAF SWSYLKPQYW GIWLGIFLL LLAFFVFLRL
 51 DLTKGLGIW IGHKAKQRT RAQTNLQYCF PHWTEQQREQ VIDKMFVAVD
 101 QVMFIGEIA IRSKKHLQKR SEFIGLEHIE QAKAEGKNII LMVPHGWAI
 151 ASGIILHTQG MPMTSMYNPH RNPLVDLWLT ITRQRFEGKM HARQNGIKPF

201 LSHVRKGMG YYLPDEDFGA EQSVFVDFG TYKATLPGLN KMAKLSKAVV
 251 IMFPPRYNAE TGKYEMEIH PAMNLSDDPEQ SARAMEEIE SFVTPAPEQY
 301 VWILQLLRTR KDGEDLYD*

5 SEQ. ID NO:80

Nucleotide sequence of DNA coding region of the *Moraxella catarrhalis* MsbB gene

1 ATGAGTTGCC ATCATCAGCA TAAGCAGACA CCCAAACACG CCATATCCAT
 51 TAAGCATATG CCAAGCTTGA CAGATACTCA TAAACAAAGT AGCCAAGCTG
 101 AGCCAAAATC GTTTGAATGG GCGTTTTTAC ATCCCAAATA TTGGGGAGTT
 151 TGGCTGGCTT TTGCGTTGAT TTTACCGCTG ATTTTCTAC CGCTGCGTTG
 201 GCAGTTTGG ATCGGCAAGC GTCTTGGCAT TTTGGTACAT TACTTAGCTA
 251 AAAGCCGAGT TCAAGACACT CTAACCAACC TGCAGCTTAC CTTCCTCAAT
 301 CAACCAAAAT CAAAACACAA GGCCACCGCA CGGCAAGTAT TTATTAATCA
 15 351 AGGTATTGGT ATTTTGAAA GTTTATGTGC ATGGTTTCGC CCTAATGTCT
 401 TTAACGCAC TTTTAGCATT TCTGGTTTAC AGCATTGTAT TGATGCCCAA
 451 AAACAAAATA AAGCGGTGAT TTTACTTGGT GGACATCGCA CGACGCTTGA
 501 TTTGGGCGGT CGGTTATGTA CACAGTTTTT TCGGCGGGAC TCGGTGTATC
 551 GCCCACAAAA CAACCTTTTG CTTGAATGGT TTATCTATAA TGCACGCCGC
 20 601 TCCATCTTTG ATGAGCAAA CTCAAATCGT GATATGAAAA AACTCATCAC
 651 TCGGCTCAAA CAAGGTCGGA TAATTGGTA TTCACCTGAT CAAGATTTTG
 701 GTCTTGAGCA TGGCGTGATG GCGACCTTTT TTGGTGTGCC TGCAGCAACG
 751 ATTACCGCTC AGCGTCGTCT TATTAAGCTG GGTGATAAAG CCAATCCTCC
 801 TGTTCATCAT ATGATGGATA TGCTCAGACA AACGCCCGAT TATATCGCAA
 25 851 AAGGTCACCG TCCACATTAT CACATCAGCC TAAGCGCTGT GTTAAAAAAT
 901 TATCCAGCG ATGACGAAAC CGCCGATGCT GAACGCATCA ATCGACTGAT
 951 TGAGCAAAAT ATTCAAAAG ATTTAACCCA GTGGATGTGG TTTTCATCGCC
 1001 GCTTTAAAC TCAAGCCGAT GACACCAATT ACTATCAACA TTAATG

30 Protein Sequence - 28% identity and 37 similarity with MsbB of *E. coli*

1 MSCHHQHKQT PKHAISIKHM PSLTDTHKQS SQAEPKSEFW AFLHPKYWGV
 51 WLAFAFALPL IFPLRWQFW IGKRLGILVH YLAKSRVQDT LTNLQLTFPN
 101 QPKSKHKATA RQVFINGGIG IFESLCAWFR PNVEKRTFSI SGLQLHIDAQ
 151 KQNKAVILLG GHRTTLDLGG RLCTOFFAAD CVYRPQNNPL LEWFIYNARR
 35 201 CIFDEQISNR DMKKLITRLK QGRIIWYSPD QDFGLEHGMV ATFFGVPAAT
 251 ITAQRRLIKL GDKANPPVII MMDMLRQTPD YIAKGHRPHY HISLSAVLKN
 301 YPSDDETADA ERINRLIEQN IQKDLTQWMW FHRREKTQAD DTNYQQH*

SEQ. ID NO:81

40 Nucleotide sequence of DNA coding region of the *Neisseria* (meningococcus B) MsbB gene

1 ATGAAATTTA TATTTTTGT ACTGTATGTT TTGCAGTTTC TGCCGTTTGC
 51 GCTGCTGCAC AAACCTGCGG ACCTGACGGG TTTGCTCGCC TACCTTTTGG
 101 TCAAACCCCG CCGCCGTATC GGCGAAATCA ATTTGGCAA ATGCTTTCCC
 15 151 GAGTGGGACG GAAAAAGCG CGAAACCGTA TTGAAGCAGC ATTTCAAACA
 201 TATGGCGAAA CTGATGCTTG AATACGGCTT ATATTGGTAC GCGCCTGCCG
 251 GCGCTTTGAA ATCGCTGGTG CGTTACCGCA ATAAGCATT TTTGGACGAC
 301 GCGCTGGCGG CCGGGGAAAA AGTCATCATT CTGTACCGC ACTTCACCGC
 351 GTTCGAGATG GCGGTGTACG CGCTTAATCA GGATGTACCG CTGATCAGTA
 50 401 TGTATTCCCA CCAAAAAAC AAGATATTGG ACGCACAGAT TTTGAAAGGC
 451 CGCAACCGCT ACGACAATGT CTTCTTATC GGGCGCACCG AAGGCGTGCG
 501 CGCCCTCGTC AAACAGTTCC GCAAAAGCAG CGCGCCGTTT CTGTATCTGC
 551 CCGATCAGGA TTTCCGACGC AACGATTCGG TTTTGTGGA TTTTTCGGT
 601 ATTCAGACGG CAACGATTAC CGGCTTGAGC CGCATTGCCG CGCTTGCAAA
 55 651 TGCAAAAGT ATACCCGCCA TCCCGTCCG CGAGGCGGAC AATACGGTTA
 701 CATTGCATT CTACCCGGCT TGGGAATCCT TTCCGAGTGA AGATGCGCAG
 751 CCGGACGCGC AGGCGATGAA CCGTTTTATC GAGGAACCGT GCGCGAACAT
 801 CCCGAGCAGT ATTTTGGCT GCACAAGCGT TTCAAACCC GTCCGGAAGG
 851 CAGCCCCGAT TTTACTGAT ACGTAA

Protein Sequence - 25% identity and 36% identity with MsbB *E. coli*

1 MKFIFVLYV LQFLPFALLH KLADLTGLLA YLLVKPRRRI GEINLAKCFP
 51 EWDGKKRET V LKQHFHMAK LMLEYGLWY APAGRLKSLV RYRNKHYLDD
 101 ALAAGEKVII LYPHFTAFEM AVYALNQDVP LISMYSHQKN KILDAQILKG
 15 151 RNRYNVFLI GRTEGVRALV KQFRKSSAPF LYLDPQDFGR NDSVFVDFG
 201 IQTATTGLS RIAALANAKV IPAIPVREAD NTVTLHFYPA WESFPSEDAQ
 251 ADAQRMNRFI EEPKANIPSS IFGCTSVSKP VRKAAPITFD T*

We Claim:

1. A genetically-engineered bleb preparation from a Gram-negative bacterial strain characterized in that said preparation is obtainable by employing one or more processes selected from the following group:
- a) a process of reducing immunodominant variable or non-protective antigens within the bleb preparation comprising the steps of determining the identity of such antigen, engineering a bacterial strain to produce less or none of said antigen, and making blebs from said strain;
 - b) a process of upregulating expression of protective OMP antigens within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of a gene encoding said antigen such that said gene is expressed at a level higher than in the non-modified bleb, and making blebs from said strain;
 - c) a process of upregulating expression of conditionally-expressed, protective OMP antigens within the bleb preparation comprising the steps of identifying such an antigen, engineering a bacterial strain so as to remove the repressive control mechanisms of its expression, and making blebs from said strain;
 - d) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS toxic, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain;
 - e) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS less toxic, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of said gene such that said gene is expressed at a level higher than in the non-modified bleb, and making blebs from said strain;

- 5 f) a process of reducing lipid A toxicity within the bleb preparation and increasing the levels of protective antigens, comprising the steps of engineering the chromosome of a bacterial strain to incorporate a gene encoding a Polymyxin A peptide, or a derivative or analogue thereof, fused to a protective antigen, and making blebs from said strain; or
- g) a process of creating conserved OMP antigens on the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to delete variable regions of a gene encoding said antigen, and making blebs from said strain.

10

2. The genetically-engineered bleb preparation of Claim 1, wherein said preparation is obtainable by employing one or more further processes selected from the following group:

- 15 h) a process of reducing expression within the bleb preparation of an antigen which shares a structural similarity with a human structure and may be capable of inducing an auto-immune response in humans, comprising the steps of identifying a gene involved in the biosynthesis of the antigen, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain; or
- 20 i) a process of upregulating expression of protective OMP antigens within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce into the chromosome one or more further copies of a gene encoding said antigen controlled by a heterologous, stronger promoter sequence, and making blebs from said strain.
- 25

3. The bleb preparation of claim 1 or 2, wherein the engineering steps of processes a), b), c), d), e), h) and i) are carried out by homologous recombination events between a sequence of at least 30 nucleotides on the bacterial chromosome, and a sequence of at least 30 nucleotides on a vector transformed within the strain.

30

4. The bleb preparation of claim 3, wherein the engineering steps are carried out by double cross-over homologous recombination events between two sequences of at least 30 nucleotides on the bacterial chromosome separated by nucleotide sequence 'X', and two sequences of at least 30 nucleotides on a vector transformed within the strain separated by nucleotide sequence 'Y', wherein during the recombination event X and Y are interchanged.
5. The bleb preparation of claim 4, wherein the two nucleotide sequences are of approximately the same length, and wherein the vector is a linear DNA molecule.
6. The bleb preparation of claim 4 or 5, wherein the recombination events of processes a), b), c), d), e) and h) are carried out within the region of the chromosome 1000 bp upstream of the initiation codon of the gene of interest.
7. The bleb preparation of claim 6, wherein for process a), d) or h) nucleotide sequence X comprises part of the promoter region of the gene, and nucleotide sequence Y comprises either a weak promoter region, or no promoter region.
8. The bleb preparation of claim 6, wherein for process b) or e) nucleotide sequence Y comprises a strong promoter region for the bacterium.
9. The bleb preparation of claim 8, wherein nucleotide sequence Y is inserted 200-600 bp upstream of the initiation codon of the gene of interest.
10. The bleb preparation of claim 9, wherein nucleotide sequence Y is inserted approximately 400 bp upstream of the initiation codon of the gene of interest.
11. The bleb preparation of claim 6, wherein for process c) nucleotide sequence X comprises part of the repressive control sequence of the promoter, and nucleotide sequence Y comprises no such sequence.

12. The bleb preparation of claim 4 or 5, wherein the recombination events of processes a), d), and h) are carried out such that nucleotide sequence X comprises part of the coding sequence of the gene of interest.
- 5 13. The bleb preparation of claim 4 or 5, wherein the recombination events of process i) is carried out such that nucleotide sequence Y comprises the further copy of the gene within an expression cassette.
14. The bleb preparation of claims 1-13 isolated from a modified *Neisseria*
10 *meningitidis* B strain.
15. The bleb preparation of claim 14, obtainable by employing at least process b) and/or i), wherein one or more genes are upregulated from a list comprising: NspA, Hsf-like, Hap, PorA, PorB, OMP85, PilQ, PldA, FrpB, TbpA, TbpB, FrpA, FrpC,
15 LbpA, LbpB, FhaB, HasR, lipo02, Tbp2 (lipo28), MltA (lipo30), and ctrA.
16. The bleb preparation of claim 14 or 15, obtainable by employing at least process a), wherein one or more genes are downregulated from a list comprising: PorA, PorB, PilC, TbpA, TbpB, LbpA, LbpB, Opa, and Opc.
20
17. The bleb preparation of claims 14-16, obtainable by employing at least process d), wherein one or more genes are downregulated from a list comprising: htrB, msbB and lpxK.
- 25 18. The bleb preparation of claims 14-17, obtainable by employing at least process e), wherein one or more genes are upregulated from a list comprising: pmrA, pmrB, pmrE, and pmrF.
19. The bleb preparation of claims 14-18, obtainable by employing at least process
30 c), wherein the repressive control sequence of the promoter is the *fur* operator region of either or both of the TbpB or LbpB genes.

20. The bleb preparation of claims 14-19, obtainable by employing at least process h), wherein one or more genes are downregulated from a list comprising: galE, siaA, siaB, siaC, siaD, ctrA, ctrB, ctrC, and ctrD.
- 5 21. The bleb preparation of claims 1-13 isolated from a modified *Moraxella catarrhalis* strain.
22. The bleb preparation of claim 21, obtainable by employing at least process b) and/or i), wherein one or more genes are upregulated from a list comprising:
10 OMP106, HasR, PilQ, OMP85, lipo06, lipo10, lipo11, lipo18, P6, ompCD, CopB, D15, OmpLA1, Hly3, LbpA, LbpB, TbpA, TbpB, OmpE, UspA1, UspA2, and Omp21.
23. The bleb preparation of claim 21 or 22, obtainable by employing at least
15 process a), wherein one or more genes are downregulated from a list comprising: CopB, OMP106, OmpB1, TbpA, TbpB, LbpA, and LbpB.
24. The bleb preparation of claims 21-23, obtainable by employing at least process d), wherein one or more genes are downregulated from a list comprising:
20 and lpxK.
25. The bleb preparation of claims 21-24, obtainable by employing at least process e), wherein one or more genes are upregulated from a list comprising: pmrA, pmrB, pmrE, and pmrF.
25
26. The bleb preparation of claims 1-13 isolated from a modified *Haemophilus influenzae* strain.
27. The bleb preparation of claim 26, obtainable by employing at least process b) and/or i), wherein one or more genes are upregulated from a list comprising: D15, P6, TbpA, TbpB, P2, P5, OMP26, HMW1, HMW2, HMW3, HMW4, Hia, Hsf, Hap, Hin47, and Hif.
30

28. The bleb preparation of claim 26 or 27, obtainable by employing at least process a), wherein one or more genes are downregulated from a list comprising: P2, P5, Hif, IgA1-protease, HgpA, HgpB, HMW1, HMW2, Hxu, TbpA, and TbpB.

5

29. The bleb preparation of claims 26-28, obtainable by employing at least process d), wherein one or more genes are downregulated from a list comprising: htrB, msbB and lpxK.

10 30. The bleb preparation of claims 26-29, obtainable by employing at least process e), wherein one or more genes are upregulated from a list comprising: pmrA, pmrB, pmrE, and pmrF.

15 31. A genetically-engineered bleb preparation from a Gram-negative bacterial strain characterized in that said preparation is obtainable by employing a process comprising the steps of: introducing a heterologous gene, optionally controlled by a strong promoter sequence, into the chromosome by homologous recombination, and making blebs from said strain.

20 32. The bleb preparation of claim 31, wherein the Gram-negative strain is *Neisseria gonorrhoeae*, and the heterologous gene is a protective OMP from *C. trachomatis*.

25 33. The bleb preparation of claim 31, wherein the Gram-negative strain is *Haemophilus influenzae*, and the heterologous gene is a protective OMP from *Moraxella catarrhalis*.

30 34. The bleb preparation of claim 31, wherein the Gram-negative strain is *Moraxella catarrhalis*, and the heterologous gene is a protective OMP from *Haemophilus influenzae*.

35. A vaccine comprising the bleb preparation of claims 1-34 and a pharmaceutically acceptable excipient.
36. A meningococcal vaccine comprising the bleb preparation of claims 14-20 and
5 one or more plain or conjugated meningococcal capsular polysaccharides selected from the serotypes A, C, Y or W.
37. A meningitis vaccine comprising the preparation of claims 14-20 or the bleb preparation of claim 32, a conjugated *H. influenzae* b capsular polysaccharide, and
10 one or more plain or conjugated pneumococcal capsular polysaccharides.
38. An otitis media vaccine comprising the bleb preparation of claims 21-25 and one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against non-typeable *H. influenzae* infection.
15
39. An otitis media vaccine comprising the bleb preparation of claims 26-30 and one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against *Moraxella catarrhalis* infection.
40. The otitis media vaccine of claim 38 or 39, wherein the vaccine additionally comprises one or more protein antigens that can protect a host against *Streptococcus pneumoniae* and/or one or more antigens that can protect a host against RSV and/or one or more antigens that can protect a host against influenza virus.
20
41. A vector suitable for performing the recombination events described in claims 3-34.
25
42. A modified Gram-negative bacterial strain from which the bleb preparation of claims 1-34 is made.
30
43. An immuno-protective and non-toxic Gram-negative bleb, ghost, or killed whole cell vaccine suitable for paediatric use.

44. An isolated polynucleotide sequence which hybridises under highly stringent conditions to at least a 30 nucleotide portion of the nucleotides in SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof.

5

45. The isolated polynucleotide sequence of claim 44 comprising at least 30 nucleotides.

46. The isolated polynucleotide sequence of claim 45 comprising at least 30 nucleotides of SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof.

10

47. Use of the isolated polynucleotide sequence of claims 44-46 in performing a homologous recombination event within 1000 bp upstream of a Gram-negative bacterial chromosomal gene in order to either increase or decrease expression of said gene.

15

48. A process of genetically upregulating expression of a gene within a Gram-negative bacterial strain comprising the steps of: making a vector comprising a strong promoter sequence and a nucleotide sequence of at least 30 nucleotides which is capable of recombination with a sequence of at least 30 nucleotides in the 1000 bp upstream of the gene, transforming a bacterial strain with the vector, and performing a homologous recombination event between the chromosome and the vector to introduce the promoter within 1000 bp upstream of the initiation codon of the gene.

20

49. The process of claim 48, wherein the promoter is introduced within 200-600 bp upstream of the initiation codon of the gene.

25

50. The process of claim 49, wherein the promoter is introduced approximately 400 bp upstream of the initiation codon of the gene.

30

51. The process of claim 48-50, wherein the homologous recombination event is effected through a double cross-over between two sequences of at least 30 nucleotides

on the bacterial chromosome separated by nucleotide sequence 'X', and two sequences of at least 30 nucleotides on the vector separated by nucleotide sequence 'Y' which comprises the strong promoter sequence, wherein the vector is a linear nucleotide sequence, and wherein during the recombination event X and Y are
5 interchanged.

52. The process of claim 51, wherein the two nucleotide sequences are of approximately the same length.

10 53. A modified bacterial strain obtainable by the processes of claims 48-52.

54. A method to produce a genetically-engineered bleb preparation from a Gram-negative bacterial strain by employing one or more processes selected from the following group:

15 a) a process of reducing immunodominant variable or non-protective antigens within the bleb preparation comprising the steps of determining the identity of such antigen, engineering a bacterial strain to produce less or none of said antigen, and making blebs from said strain;

 b) a process of upregulating expression of protective OMP antigens
20 within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of a gene encoding said antigen such that said gene is expressed at a level higher than in the non-modified bleb, and making blebs from said strain;

 c) a process of upregulating expression of conditionally-expressed,
25 protective OMP antigens within the bleb preparation comprising the steps of identifying such an antigen, engineering a bacterial strain so as to remove the repressive control mechanisms of its expression, and making blebs from said strain;

 d) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid
30 A portion of LPS toxic, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain;

e) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS less toxic, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of said gene such that said gene is expressed at a level
5 higher than in the non-modified bleb, and making blebs from said strain;

f) a process of reducing lipid A toxicity within the bleb preparation and increasing the levels of protective antigens, comprising the steps of engineering the chromosome of a bacterial strain to incorporate a gene encoding a Polymyxin A peptide, or a derivative or analogue thereof, fused to a protective antigen, and making
10 blebs from said strain;

g) a process of creating conserved OMP antigens on the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to delete variable regions of a gene encoding said antigen, and making blebs from said strain;

h) a process of reducing expression within the bleb preparation of an antigen which shares a structural similarity with a human structure and may be capable of inducing an auto-immune response in humans, comprising the steps of identifying a gene involved in the biosynthesis of the antigen, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from
15 said strain; or

i) a process of upregulating expression of protective OMP antigens within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce into the chromosome one or more further copies of a gene encoding said antigen controlled by a heterologous, stronger
20 promoter sequence, and making blebs from said strain.

55. A method of immunizing a human host against a disease caused by infection of one or more of the following: *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Haemophilus influenza*, *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia pneumonia*, which method comprises administering to the
30 host an immunoprotective dose of the bleb preparation of claims 1-34.

56. Use of the bleb preparation of claims 1-34 in the manufacture of a medicament for immunizing a human host against a disease caused by infection of one or more of the following: *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Haemophilus influenza*, *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia pneumonia*.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9





Figure 10

A



B




Figure 11




Figure 12

Schematic representation of the recombinant PCR strategy used to delete the *lacO* in the chimeric *porA/lacO* promoter.




Figure 13




Figure 14




Figure 15

SDS Page 1




Figure 16




Figure 17

